



Forage grasses with lower uptake of caesium and strontium could provide 'safer' crops for radiologically contaminated areas

Beth Penrose, Nicholas A. Beresford, Neil M. J. Crout, J. Alan Lovatt, Russell Thomson, Martin R. Broadley

Published: May 1, 2017 • <https://doi.org/10.1371/journal.pone.0176040>

Abstract

Substitution of a species or cultivar with higher uptake of an element by one with lower uptake has been proposed as a remediation strategy following accidental releases of radioactivity. However, despite the importance of pasture systems for radiological dose, species/cultivar substitution has not been thoroughly investigated for forage grasses. 397 cultivars from four forage grass species; hybrid ryegrass (*Lolium perenne* L. x *Lolium multiflorum* Lam.), perennial ryegrass (*Lolium perenne* L.), Italian ryegrass (*Lolium multiflorum* Lam.) and tall fescue (*Festuca arundinacea* Shreb.); were sampled from 19 field-based breeding experiments in Aberystwyth and Edinburgh (UK) in spring 2013 and analysed for caesium (Cs) and strontium (Sr) concentrations. In order to calculate concentration ratios (CRs; the concentration of an element in a plant in relation to the concentration in the soil), soils from the experiments were also analysed to calculate extractable concentrations of Cs and Sr. To test if cultivars have consistently low Cs and Sr concentration ratios, 17 hybrid ryegrass cultivars were sampled from both sites again in summer 2013 and spring and summer 2014. Tall fescue cultivars had lower Cs and Sr CRs than the other species. Three of the selected 17 hybrid ryegrass cultivars had consistently low Cs CRs, two had consistently low Sr CRs and one had consistently low Cs and Sr CRs. Cultivar substitution could reduce Cs CRs by up to 14-fold and Sr CRs by 4-fold in hybrid ryegrass. The identification of species and cultivars with consistently low CRs suggests that species or cultivar substitution could be an effective remediation strategy for contaminated areas.

Citation: Penrose B, Beresford NA, Crout NMJ, Lovatt JA, Thomson R, Broadley MR (2017) Forage grasses with lower uptake of caesium and strontium could provide 'safer' crops for radiologically contaminated areas. PLoS ONE 12(5): e0176040. <https://doi.org/10.1371/journal.pone.0176040>

Editor: Ivan Baxter, United States Department of Agriculture, Agricultural Research Service, UNITED STATES

Received: August 23, 2016; **Accepted:** April 4, 2017; **Published:** May 1, 2017

Copyright: © 2017 Penrose et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability: All relevant data are within the paper and its Supporting Information files.

Funding: Funded by Natural Environment Research Council: <http://www.nerc.ac.uk/>. NE/K500951/1 received by BP. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Introduction

Radiocaesium (^{134}Cs and ^{137}Cs) and radiostrontium (primarily ^{90}Sr) are anthropogenic radionuclides that have been distributed in the environment due to nuclear weapons testing, nuclear power production and nuclear accidents [1]. Radioisotopes of Cs were the primary long-term contributors to human doses received after the Windscale [2](1957) and Chernobyl [3] (1986) accidents, and are now the principal radiological concern in areas affected by the accident at Fukushima in 2011 [4]. Strontium-90 is a major contributor to doses in areas contaminated by releases from the Mayak facility in the Russian Urals [5] and is present in soils of the Chernobyl Exclusion Zone at activity concentrations close to those of ^{137}Cs [6]. These radionuclides are of particular concern as they are readily incorporated into biological systems due to their chemical similarity to the biologically essential elements potassium (K; radiocaesium) and calcium (Ca; radiostrontium). In the long term, radiocaesium and radiostrontium enter the human food chain mainly via consumption of crop plants grown on contaminated soils or products from animals fed on contaminated fodder.

A number of remediation strategies that aim to reduce the transfer of radionuclides to crop/fodder plants are available, and include mechanical soil amelioration techniques, chemical amendments and plant based strategies. Mechanical techniques such as ploughing can reduce transfer of Cs and Sr by 2–10 fold and chemical amendments such as the addition of organic matter, liming or application of fertiliser can reduce transfer by 1.3–3.0 fold, 1.5–4.0 fold and 1.5–3.0 fold, respectively [7]. For radiocaesium there are also highly efficient binders which can be orally administered to livestock to reduce absorption in the gastrointestinal tract and hence transfer to milk or meat [8]; however, this is not the case for radiostrontium[9]. Alternatively, plant based strategies are available; Cs and Sr uptake from soil can vary by two orders of magnitude between crop plant species [7] thus species substitution -where a crop species with high uptake of an element is replaced with a crop species with low uptake of an element- has the potential to cause a significant reduction in soil-plant transfer of these radionuclides [10]. However, changing production from one crop to another can be challenging; agronomic knowledge and a viable market must be available for the new crop [11][7]. Conversely, cultivar substitution for a given crop is likely to be readily applicable, socially acceptable and require no new skills, knowledge or markets. However, not enough is known about its efficacy for it to be widely recommended [12][13].

Ohlenschlaeger and Gissel-Nielsen [14] [15] investigated variation in Cs uptake between two perennial ryegrass cultivars and found little variation (maximum 1.3) in radiocaesium soil-plant concentration ratios. It is likely that more variation would have been found if more cultivars were included in the study. Studies investigating inter-cultivar variation in uptake of chemical analogues of Cs and Sr (K, Ca and Mg) have shown that heritability of shoot concentrations of these elements is high in ryegrasses [16][17][18]; and fescues [16][17][19], and therefore breeding and selection of cultivars for low concentrations of these elements (and thus the analogous elements Cs and Sr) may be feasible [20].

Cultivars of Italian ryegrass [21] and tall fescue [22] with consistently high shoot Mg concentrations have been identified. However, other studies have found significant genotype x environment interactions for both ryegrasses [17] and fescues [17] [19] [18], and therefore cultivars with apparently high or low elemental shoot concentrations may behave inconsistently over multiple sites and/or years. Environmental factors can have a large effect on the uptake of Cs and Sr, and identifying cultivars that display consistently low concentration ratios over time and between sites has previously been challenging [12].

As forage grasses exhibit limited self-pollination, individuals of the same cultivar are not genetically identical, and are usually referred to as 'populations' by grass breeders. Although 'cultivar' is usually used to describe commercial varieties, the term 'cultivar' is used in this paper to describe non-commercial varieties in addition to commercial varieties.

The aim of the study described was to investigate the variation in stable caesium (Cs) and strontium (Sr) uptake between 397 cultivars from four commonly used forage grass species using measurements from a programme of long term grass breeding experiments. Particular attention was paid to the identification of cultivars that have consistently low Cs and/or Sr concentration ratios at both sites, over two study years and from both spring and summer harvests.

Materials and methods

Overview

Caesium and strontium analyses were made on samples of 397 cultivars from four forage grass species. The grass species were: hybrid ryegrass (*Lolium perenne* L. x *Lolium multiflorum* Lam.), perennial ryegrass (*Lolium perenne* L.), Italian ryegrass (*Lolium multiflorum* Lam.) and tall fescue (*Festuca arundinacea* Shreb.). Samples were collected from established breeding experiments in Aberystwyth, UK and Edinburgh, UK, in the spring and summer of 2013 and 2014. Due to the ongoing structure of the ongoing breeding experiments not all cultivars were available for sampling at each site or time period. As one of our key objectives was to test whether cultivars behaved similarly in terms of Cs and Sr uptake over different seasons and in multiple years, we organised our work to maximise the number of samples from each site and time period.

Experimental design

The breeding programme is structured into individual experiments, each of which consisted of between 40 and 416 plots (Table 1), with between 10 and 104 cultivars grown in a randomised block design in a given field and sown between one and three years prior to our sampling.

| Experiment number | Site | Field number | Sowing date | Plot area (m ²) | No. of plots | No. of cultivars | No. of replicates of each cultivar | No. of cuts in 2013 | No. of cuts in 2014 | Spring cut 2013 | Summer cut 2013 | Spring cut 2014 | Summer cut 2014 |
|-------------------|-------------|--------------|-------------|-----------------------------|--------------|------------------|------------------------------------|---------------------|---------------------|-----------------|-----------------|-----------------|-----------------|
| 1 | Aberystwyth | 1 | 10/08/10 | 2 | 66 | 17 | 4 | 7 | 7 | 20/05/13 | - | - | - |
| 2 | Edinburgh | 4 | 10/08/10 | 2 | 66 | 17 | 4 | 7 | 7 | 20/05/13 | - | - | - |
| 3 | Aberystwyth | 1 | 10/08/10 | 1 | 64 | 16 | 4 | 7 | 7 | 20/05/13 | - | - | - |
| 4 | Aberystwyth | 1 | 10/08/10 | 1 | 40 | 10 | 4 | 7 | 7 | 01/06/13 | - | - | - |
| 5 | Aberystwyth | 1 | 10/08/10 | 1 | 72 | 18 | 4 | 7 | 7 | 20/05/13 | - | - | - |
| 6 | Edinburgh | 4 | 10/08/10 | 1 | 72 | 18 | 4 | 7 | 7 | 01/06/13 | - | - | - |
| 7 | Aberystwyth | 2 | 09/08/11 | 2 | 152 | 38 | 4 | 7 | 7 | 20/05/13 | 09/08/13 | 20/05/14 | 09/08/14 |
| 8 | Edinburgh | 4 | 22/08/11 | 2 | 100 | 25 | 4 | 7 | 7 | 20/05/13 | 27/08/13 | 19/05/14 | 20/08/14 |
| 9 | Aberystwyth | 2 | 09/08/11 | 4.68 | 108 | 27 | 4 | 9 | 5 | 01/06/13 | - | - | - |
| 10 | Aberystwyth | 2 | 09/08/11 | 4.68 | 78 | 19 | 4 | 9 | 5 | 01/06/13 | - | - | - |
| 11 | Aberystwyth | 2 | 09/08/11 | 4.68 | 42 | 10 | 4 | 6 | 8 | 01/06/13 | - | - | - |
| 12 | Aberystwyth | 3 | 14/09/12 | 1.2 | 84 | 21 | 4 | 7 | 7 | 04/06/13 | - | - | - |
| 13 | Aberystwyth | 3 | 14/09/12 | 2 | 248 | 62 | 4 | 7 | 7 | 22/05/13 | 29/08/13 | 19/05/14 | 12/08/14 |
| 14 | Aberystwyth | 3 | 13/09/12 | 4.68 | 108 | 27 | 4 | 5 | 9 | 22/05/13 | - | - | - |
| 15 | Edinburgh | 4 | 20/08/12 | 2 | 64 | 16 | 4 | 7 | 7 | 22/05/13 | 23/08/13 | 19/05/14 | 20/08/14 |
| 16 | Aberystwyth | 3 | 13/09/12 | 4.68 | 80 | 20 | 4 | 5 | 9 | 30/05/13 | - | - | - |
| 17 | Aberystwyth | 3 | 14/09/12 | 1.2 | 416 | 104 | 4 | 7 | 7 | 31/05/13 | - | - | - |
| 18 | Edinburgh | 4 | 20/08/12 | 2 | 126 | 42 | 3 | 7 | 7 | 12/06/13 | - | - | - |
| 19 | Edinburgh | 4 | 21/08/12 | 2 | 126 | 42 | 3 | 7 | 7 | 12/06/13 | - | - | - |

<https://doi.org/10.1371/journal.pone.0176040.t001>

Table 1. Sowing and harvest dates, number of cultivars, replicates, plots, cuts per year and the plot area in each of the 19 experiments sampled in this study.

<https://doi.org/10.1371/journal.pone.0176040.t001>

The Aberystwyth experiments were located in three fields at the Gogerddan Campus of the Institute of Biological and Environmental Sciences (IBERS). Field 1 (52°25'40.1"N 4°01'25.5"W) contained 2010-sown plots, Field 2 (52°26'01.1"N 4°00'42.1"W) the 2011-sown plots and Field 3 (52°26'03.6"N 4°00'27.5"W) the 2012-sown plots. The Edinburgh experiments at the Roddinglaw site of Science and Advice for Scottish Agriculture (SASA) (55°55'40.9"N, 3°20'25.3"W). The distribution of cultivars and species between the sampled experiments is given in Table 2.

| Site | Hybrid ryegrass | Perennial ryegrass | Italian ryegrass | Tall fescue |
|--------------|-----------------|--------------------|------------------|-------------|
| Aberystwyth | 100 | 100 | 8 | 10 |
| Edinburgh | 20 | 100 | 10 | 2 |
| Total number | 120 (20) | 200 (20) | 17 (7) | 12 (2) |

Number in parentheses are the number of cultivars shared between both sites.

<https://doi.org/10.1371/journal.pone.0176040.t002>

Table 2. Number of different cultivars grown in Aberystwyth, Edinburgh and in total.

<https://doi.org/10.1371/journal.pone.0176040.t002>

The cutting regime varied between experiments depending on the requirements of the breeding programme (Table 1). All cuts were carried out between April and November which is typical for UK grassland management (Table 1).

Not all of the samples from these replicates could be analysed, and our results are derived from between 1 and 4 replicates of each cultivar per experiment.

Preparation and maintenance of experiment plots

The plots were ploughed using a 5-furrow reversible plough prior to sowing. Experiments 1, 3, 7, 9, 10, 11, 13, 14 and 16 were drill sown. The rest of the experiments were broadcast sown. Fertilisers were applied before the first cut and after each cut (Table 3) in all experiments apart from 14 and 16, where the fertiliser was applied twice prior to the first cut. using a calibrated drop spreader (Gandy, Minnesota, USA).

| Experiments | Fertiliser before cut 1 (kg ha ⁻¹) | Fertiliser after cut 1 (kg ha ⁻¹) | Fertiliser after cut 2 (kg ha ⁻¹) | Fertiliser after cut 3 (kg ha ⁻¹) | Fertiliser after cut 4 (kg ha ⁻¹) | Fertiliser after cut 5 (kg ha ⁻¹) | Fertiliser after cut 6 (kg ha ⁻¹) | Fertiliser after cut 7 (kg ha ⁻¹) | Fertiliser after cut 8 (kg ha ⁻¹) | Fertiliser after cut 9 (kg ha ⁻¹) |
|--------------|--|---|---|---|---|---|---|---|---|---|
| 4,5,12,17 | 57 | 57 | 57 | 57 | 57 | 57 | 57 | 57* | 57* | 57* |
| 5,6,7,23 | 57 | 57 | 57 | 57 | 57 | 57 | 57 | 57* | 57* | 57* |
| 8,9,10,18,19 | 57 | 57 | 57 | 57 | 57 | 57 | 57 | 57* | 57* | 57* |
| 9,10 | 57 | 57 | 57 | 57 | 57 | 57 | 57 | 57* | 57* | 57* |
| 14,16 | 57 | 57 | 57 | 57 | 57 | 57 | 57* | 57* | 57* | 57* |
| 15 | 57 | 57 | 57 | 57 | 57 | 57 | 57* | 57* | 57* | 57* |

All fertilisers are shown in kg ha⁻¹. Superphosphate (17%P₂O₅, Contains 10% of 23 x 13 N P₂O₅ K₂O + 7% S), was used except for when the applications are specified as ** which the fertiliser used is 20 x 20 x 20 (N:P₂O₅:K₂O) + 7% S.

Table 3. Fertiliser treatments for the 19 experiments.

<https://doi.org/10.1371/journal.pone.0176040.t003>

Sample collection and preparation

The plots were cut to approximately 5 cm height using a custom Haldrup grass harvester (Haldrup GmbH, Ilshofen, Germany) as part of the normal operations of the breeding programme. For our work a subsample (200–320 g fresh mass) of sward was taken by hand from cuttings ejected from the harvester. Samples were collected and oven dried (80–90°C) for 2–5 days depending on the moisture content of the samples. Grass samples were then ground using a cutting mill, and c. 10 g dry mass (DM) of this ground sample was retained for further analyses.

A washing test was undertaken to determine whether the concentrations of Cs and Sr were affected by washing (see S1 Table and S1 Fig). On the basis of this test samples were not washed before drying thereby reducing the sample preparation workload.

Soil properties were measured using five soil cores of 10–15 cm depth from each experimental area, taken in a W-pattern using a soil auger between June 2013 and March 2014. The five cores were homogenised and bulked to form one sample per experiment. The pH_{water} was measured using a HI 9024 microcomputer pH meter (Hanna Instruments, Padova, Italy) using the method described by Emmett *et al.* [23]. To determine moisture content, soils were weighed (wet mass) and air-dried at ~30°C for 3–11 weeks then oven dried at 80°C for 3–4 days, after which they were reweighed (dry mass). Approximately 10 g was subsampled from the bulk, and ground in a ball mill using agate bowls and balls. Soil type was determined using the soil texture layer on the UKSO soils map viewer [24].

Acid digestion of grass samples

A subsample of ~0.3 g DM of ground grass was weighed into a digestion tube and 2 mL of concentrated nitric acid (80% trace element grade HNO₃), 1 mL of hydrogen peroxide (>30% H₂O₂) and 1 mL of ultra-pure 'Milli-Q' water (18.2 MΩ cm; Fisher Scientific UK Ltd, Loughborough, UK) added. These tubes were placed into a microwave digester (Multiwave 3000 with 48-vessel 48MF50 rotor; Anton Paar GmbH, Graz, Austria) with microwave settings as follows: power = 1400 W, temperature = 140°C, pressure = 2 MPa, time = 40 minutes. Once digested, each sample was diluted with 11 mL of ultra-pure water. An aliquot of 2 mL was then removed from this sample, and diluted with a further 8 mL of ultra-pure water prior to analysis using inductively-coupled plasma mass spectrometry (ICP-MS).

Soil ammonium nitrate extraction

Ammonium nitrate (NH₄NO₃) is commonly used to extract minerals from soil, and this 'extractable fraction' is considered to be a measure of what is available to the plant [25]. To determine extractable mineral concentrations approximately 2 g of dry soil was taken from the milled samples from each experiment, and weighed into centrifuge tubes. 20 mL of 1 M NH₄NO₃ was added to each tube. Tubes were then agitated using an end-over-end shaker for ~12 h overnight. The tubes were then centrifuged at 2500 rpm for 10 minutes and 10 mL of the supernatant was passed through a 22 μm syringe-driven filter into a universal tube. 1 mL of the filtered supernatant was then transferred via pipette into ICP tubes and diluted with 9 mL of 2% nitric acid ready for analysis using ICP-MS.

Mineral analysis

Multi-element analysis of diluted solutions was undertaken by ICP-MS. Samples from the spring cut 2013 of Experiments 1, 2, 4, 7, 11, 13, 14, 15, 16 and 18 were analysed using a Thermo-Fisher Scientific X-Series^{II} instrument (Thermo Fisher Scientific Inc., Waltham, MA, USA). All other samples from the spring cut 2013, summer cut 2013, spring cut 2014 and summer cut 2014 were analysed using a Thermo-Fisher Scientific iCAP-Q instrument (Thermo Fisher Scientific, Bremen, Germany), with appropriate cross-calibration between instruments. Detailed information about the ICP-MS methods used can be found in S1 Table.

Data analysis

All data analyses were conducted using R [26]. We interpreted Cs and Sr uptake by calculating concentration ratio (CR; Eq 1) for each of the grass samples using the concentration of the element in the grass divided by the corresponding ammonium nitrate extractable soil concentration. The CR is an internationally accepted approach for radionuclide uptake assessment (e.g. [27]) here we are using it to account for variation in soil concentration of Cs and Sr in across our experimental locations. Neither Cs or Sr are essential elements and there is no evidence to suggest that their *transfer* from soil to grass is dependent upon soil Cs or Sr concentrations [28]; we acknowledge that CR varies depending upon soil type [2] [27] but this does not negate the use of CRs for the purposes of this assessment.

Where a cultivar was grown in multiple experiments at the same site, mean Cs and Sr CRs for that cultivar were calculated using concentration ratios from all the Experiments at that site. All arithmetic means are denoted by \bar{x} .

$$\text{Concentration ratio} = \frac{\text{Concentration of element in the plant (mg kg}^{-1} \text{ DW)}}{\text{Concentration of element in the soil (mg kg}^{-1} \text{ DW)}}$$

(1)

Inter-cultivar variation, the difference between the concentration of an element in the highest accumulating cultivar and the lowest accumulating cultivar shows the full extent of variation for a group of cultivars. In this study, inter-cultivar variation was calculated using Eq (2).

$$\text{Inter-cultivar variation} = \frac{\text{Mean concentration in the highest accumulating cultivar}}{\text{Mean concentration in the lowest accumulating cultivar}}$$

(2)

Results

Between species variation in mean Cs concentration ratios

Tall fescue cultivars had a lower median Cs CR (0.0424) than median Cs CRs in hybrid ryegrasses, perennial ryegrass cultivars and Italian ryegrass cultivars grown in Aberystwyth, which were similar (0.180, 0.167, 0.149, respectively; Fig 1). Median Cs CRs were similar for hybrid ryegrass (0.578) and Italian ryegrass (0.531) and were higher in perennial ryegrass (0.923) cultivars grown in Edinburgh (Fig 1).




Fig 1. Mean caesium concentration ratios for hybrid ryegrass cultivars (Aberystwyth, number of cultivars = 100; Edinburgh, number of cultivars = 29), perennial ryegrass cultivars (Aberystwyth, number of cultivars = 189; Edinburgh, number of cultivars = 100), Italian ryegrass cultivars (Aberystwyth, number of cultivars = 8; Edinburgh, number of cultivars = 16) and tall fescue cultivars (Aberystwyth, number of cultivars = 10). The upper and lower 'hinges' refer to the 25th and 75th percentiles, the limits of the upper and lower whiskers refer to data that fall within 1.5* the inter-quartile range (IQR). The outliers (signified by ♦) show data outside this 1.5*IQR. <https://doi.org/10.1371/journal.pone.0176040.g001>

Inter-cultivar variation in mean Cs concentration ratios

Caesium CRs varied by 14-fold (range = 0.06–0.79) between the 100 hybrid ryegrass cultivars grown in Aberystwyth (Fig 1), and approximately 2-fold (range = 0.44–0.74) between the 29 hybrid ryegrass cultivars grown in Edinburgh. Caesium CRs for perennial ryegrass grown in Aberystwyth (189 cultivars) and Edinburgh (100 cultivars) varied by 7-fold (range = 0.06–0.44) and 13-fold (range = 0.12–1.64), respectively (Fig 1).

Less variation was observed between the Italian ryegrass and tall fescue cultivars sampled (Fig 1); Cs CRs varied by only ≈2-fold in the eight Italian ryegrass cultivars grown in Aberystwyth (range = 0.11–0.21), the 16 Italian ryegrass cultivars grown in Edinburgh (range = 0.38–0.81) and the 10 tall fescue varieties grown in Aberystwyth (range = 0.37–0.74).

Between species variation in mean Sr concentration ratios

Aberystwyth-grown hybrid ryegrass, perennial ryegrass and Italian ryegrass median Sr CRs were similar (1.64, 1.49 and 1.68, respectively) and were greater than the median Sr CR for tall fescue (1.00). Edinburgh-grown hybrid ryegrass had a lower median CR (0.71) when compared to Italian ryegrass (1.03), which had a lower Sr CR than the perennial ryegrass cultivars (1.20).

Inter-cultivar variation in mean Sr concentration ratios

The 29 hybrid ryegrass cultivars, 100 perennial ryegrass cultivars and 16 Italian ryegrass cultivars grown in Edinburgh displayed more variation in strontium CRs (4.4-fold (range = 0.325–1.00); 2.5-fold (range = 0.482–2.28) and 2.9-fold (range = 0.285–1.10) respectively; Fig 2) than those grown in Aberystwyth (hybrid ryegrass: 1.9-fold, (range = 1.35–2.61; number of cultivars = 100); perennial ryegrass: 2.1-fold, (range = 1.13–2.32; number of cultivars = 189) and Italian ryegrass cultivars 1.9-fold, (range = 1.37–2.57; number of cultivars = 8), Fig 2). The 10 tall fescue cultivars grown in Aberystwyth varied less (1.3-fold; range = 0.88–1.15).




Fig 2. Mean strontium concentration ratios for hybrid ryegrass cultivars (Aberystwyth, number of cultivars = 100; Edinburgh, number of cultivars = 29), perennial ryegrass cultivars (Aberystwyth, number of cultivars = 189; Edinburgh, number of cultivars = 100), Italian ryegrass cultivars (Aberystwyth, number of cultivars = 8; Edinburgh, number of cultivars = 16) and tall fescue cultivars (Aberystwyth, number of cultivars = 10).

The upper and lower 'hinges' refer to the 25th and 75th percentiles, the limits of the upper and lower whiskers refer to data that fall within 1.5* the inter-quartile range (IQR). The outliers (signified by ♦) show data outside this 1.5*IQR. <https://doi.org/10.1371/journal.pone.0176040.g002>

Variance component analysis

Variance components were calculated using the mixed effects model *lme4* package [29] using data from spring cut 2013. All the Cs and Sr CRs were natural log transformed, so the residuals were distributed normally. Concentration ratio (Cs/Sr), cultivar, species and experiment location were classed as random variables in the model. Cultivar was nested within species, and experiment number was nested within experiment location in the model. Environmental factors such as experiment location and experiment number explain a large amount of the variation in both Cs and Sr CRs (Table 4). Variety explains more of the variation in Cs (~3%) than Sr (~0.8%) CRs, whereas the opposite is true for species (Cs = >0.001%; Sr = ~0.4%).

| Caesium | Variance | Std Dev | % Variation |
|---------------------|-----------|-----------|-------------|
| Experiment number | 0.0268 | 0.2271 | 27.2 |
| Experiment location | 0.0367 | 0.05131 | 27.3 |
| Variety | 0.00417 | 0.00456 | 3.1 |
| Species | 1.174e-05 | 3.391e-05 | 8.374e-05 |
| Residuals | 0.0166 | 0.129 | 12.4 |
| Strontium | | | |
| Experiment number | 0.287 | 0.536 | 23.5 |
| Experiment location | 0.883 | 0.939 | 88.0 |
| Variety | 0.00049 | 0.00074 | 0.5 |
| Species | 0.00431 | 0.0066 | 0.4 |
| Residuals | 0.0069 | 0.0083 | 7.4 |

Table 4. Variance component analysis of caesium and strontium concentration ratios from spring cut 2013. <https://doi.org/10.1371/journal.pone.0176040.t004>

Site characteristics

The soil from Aberystwyth was characterised as sand to sandy loam, and the soil from Edinburgh as clay loam to sandy loam. The soil at both sites was classified as moderately acidic [30], as the pH ranged from 4.8–6.6 (\bar{x} = 5.8) in Aberystwyth and from 5.3–6.4 (\bar{x} = 5.9) in Edinburgh (Table 5). Soil moisture contents were similar for both sites, ranging from 13.8–18.3% (\bar{x} = 16.5%) in Aberystwyth and 16.8–19.3% (\bar{x} = 18.2%) in Edinburgh.

| Experiment number | Site | Sampling date | pH | Moisture (%) | Extractable Ca concentration (mg kg ⁻¹) | Extractable Sr concentration (mg kg ⁻¹) | Extractable K concentration (mg kg ⁻¹) | Extractable Cs concentration (mg kg ⁻¹) |
|-------------------|-------------|---------------|-----|--------------|---|---|--|---|
| 1 | Aberystwyth | 05/07/2013 | 4.8 | 15.5 | 0.0059 | 3.94 | 116 | 1910 |
| 2 | Edinburgh | 05/11/2013 | 5.3 | 16.9 | 0.0116 | 14.4 | 132 | 1520 |
| 3 | Aberystwyth | 05/07/2013 | 5.0 | 13.8 | 0.0049 | 4.92 | 154 | 1610 |
| 4 | Aberystwyth | 04/09/2014 | 5.1 | 18.3 | 0.0076 | 12.8 | 128 | 1540 |
| 5 | Aberystwyth | 05/07/2013 | 5.5 | 17.8 | 0.0062 | 4.77 | 98.7 | 1290 |
| 6 | Edinburgh | 01/10/2013 | 5.9 | 18.0 | 0.0119 | 14.4 | 122 | 1520 |
| 7 | Aberystwyth | 05/07/2013 | 5.3 | 17.2 | 0.0059 | 3.87 | 117 | 1650 |
| 8 | Edinburgh | 13/06/2014 | 6.4 | 19.3 | 0.0082 | 25.5 | 98.5 | 1670 |
| 9 | Aberystwyth | 05/07/2013 | 5.1 | 16.3 | 0.0052 | 3.87 | 158 | 1610 |
| 10 | Aberystwyth | 05/07/2013 | 5.7 | 18.2 | 0.0075 | 4.15 | 143 | 1610 |
| 11 | Aberystwyth | 05/07/2013 | 5.7 | 17.7 | 0.0055 | 4.16 | 131 | 1910 |
| 12 | Aberystwyth | 05/07/2013 | 5.1 | 16.8 | 0.0064 | 4.71 | 139 | 1540 |
| 13 | Aberystwyth | 05/07/2013 | 5.9 | 15.3 | 0.0065 | 5.29 | 137 | 1610 |
| 14 | Aberystwyth | 05/07/2013 | 5.4 | 15.7 | 0.0065 | 5.75 | 133 | 2010 |
| 15 | Edinburgh | 05/11/2013 | 5.5 | 16.8 | 0.0119 | 13.9 | 140 | 1530 |
| 16 | Aberystwyth | 05/07/2013 | 6.3 | 16.4 | 0.0090 | 5.54 | 127 | 1680 |
| 17 | Aberystwyth | 05/07/2013 | 6.2 | 16.9 | 0.105 | 9.93 | 138 | 1610 |
| 18 | Edinburgh | 05/11/2013 | 6.2 | 18.0 | 0.0128 | 14.6 | 88.2 | 1680 |
| 19 | Edinburgh | 05/11/2013 | 5.8 | 19.0 | 0.0102 | 11.2 | 138 | 1310 |

Table 5. Sampling data, pH, moisture content (%), extractable Cs, Sr, K and Ca concentrations (mg kg⁻¹) in the soils from the 19 experiments in Aberystwyth and Edinburgh. <https://doi.org/10.1371/journal.pone.0176040.t005>

Soil extractable Cs was higher in Aberystwyth (range = 0.0166–0.103; \bar{x} = 0.084 mg kg⁻¹) than in Edinburgh (range = 0.00824–0.0128; \bar{x} = 0.011 mg kg⁻¹), though the converse was true for extractable Sr; the extractable Sr concentration in Aberystwyth (range = 3.47–12.6; \bar{x} = 5.28 mg kg⁻¹) was almost three times lower than that in Edinburgh, (range = 11.2–25.5; \bar{x} = 15.7 mg kg⁻¹; Table 5). Concentrations of extractable Ca and K were relatively similar between the two sites (K: Aberystwyth, range = 90.7–191 mg kg⁻¹, \bar{x} = 136 mg kg⁻¹; Edinburgh, range = 86.2–139 mg kg⁻¹, \bar{x} = 116 mg kg⁻¹; Ca: Aberystwyth, range = 1010–2070 mg kg⁻¹, \bar{x} = 1680 mg kg⁻¹; Edinburgh, range = 1370–1790 mg kg⁻¹, \bar{x} = 1570; Table 5).

Consistency of inter-cultivar variation between cuts and locations

To compare the consistency of inter-cultivar variation in Cs and Sr uptake over time and between sites, the 17 hybrid ryegrass cultivars that were grown in both Aberystwyth and Edinburgh were selected from the larger population shown in Figs 1 and 2. These cultivars (Table 6) were sampled in spring and summer (spring = May-June, summer = August-September; Table 1) in both 2013 and 2014. The Cs and Sr CRs were estimated using the same method as above. Caesium and strontium CRs were higher in the summer cuts of these 17 hybrid ryegrass cultivars than in the spring cuts in every location and 2014 harvests tended to have higher Cs and Sr concentration ratios than 2013 harvests (Fig 3A and 3B).




Fig 3. Mean caesium concentration ratios of 17 hybrid ryegrass cultivars from Aberystwyth and Edinburgh. Individual bars from left-right represent the: spring cut 2013, summer cut 2013, spring cut 2014, summer cut 2014. The numbers above each panel refers to the cultivar number.
<https://doi.org/10.1371/journal.pone.0176040.g003>

| Cultivar number | Experiment numbers Aberystwyth | Experiment numbers Edinburgh |
|-----------------|--------------------------------|------------------------------|
| 11 | 1, 3, 7, 13 | 8, 15, 2 |
| 12 | 7, 13 | 8, 15 |
| 334 | 7, 13 | 8, 15 |
| 335 | 7, 13 | 8, 15 |
| 336 | 7, 13 | 8, 15 |
| 337 | 7, 13 | 8, 15 |
| 338 | 7, 13 | 8, 15 |
| 339 | 7, 13 | 8, 15 |
| 340 | 7, 13 | 8, 15 |
| 341 | 7, 13 | 8, 15 |
| 342 | 7, 13 | 8, 15 |
| 343 | 7, 13 | 8, 15 |
| 344 | 7, 13 | 8, 15 |
| 345 | 7, 13 | 8, 15 |
| 346 | 7, 13 | 8, 15 |
| 347 | 7, 13 | 8, 15 |
| 348 | 7, 13 | 8, 15 |

<https://doi.org/10.1371/journal.pone.0176040.t006>

Table 6. Experiment numbers of the 17 hybrid ryegrass cultivars grown in Aberystwyth and Edinburgh.
<https://doi.org/10.1371/journal.pone.0176040.t006>

The inter-varietal variation in Cs CRs for the 17 selected hybrid ryegrass cultivars ranged between 1.46 (spring cut 2013, Edinburgh) and 8.85 (summer cut 2014, Aberystwyth). For all cuts, the inter-varietal variation in Cs CRs was higher in Aberystwyth than in the equivalent cut in Edinburgh (Fig 3). The inter-varietal variation in Sr CRs (Fig 4) ranged from 1.43 (spring cut 2014, Edinburgh) to 1.79 (summer cut 2013, Edinburgh).




Fig 4. Mean strontium concentration ratios of 17 hybrid ryegrass cultivars from Aberystwyth and Edinburgh. Individual bars from left-right represent the: spring cut 2013, summer cut 2013, spring cut 2014, summer cut 2014. The numbers above each panel refers to the cultivar number.
<https://doi.org/10.1371/journal.pone.0176040.g004>

Consistency of lower concentration ratios between cuts and locations

The mean Cs CR and mean Sr CR of the 17 hybrid ryegrass cultivars in each of the eight site/harvest combinations were ranked. These ranks were then summed to create a total rank for each cultivar (S3 Table).

The total ranks were compared to a distribution of random ranks. This distribution was created by summing eight (one for each location-harvest combination) randomly-generated numbers between one and 17 to create a simulated 'total rank'. This process was iterated 1000 times, and the 5th and 95th percentile were calculated. Any cultivar with a total rank below the 5th percentile was considered significantly consistently low-ranking, and any cultivar with a total rank above the 95th percentile was deemed significantly consistently high-ranking.

Two cultivars (335 and 346), were identified as having consistently lower Sr CRs and three cultivars (334, 342 and 343) were identified as having consistently lower Cs CRs. One cultivar (339) had consistently lower Cs and Sr CRs (S3 Table in the Supplementary Information). Five cultivars were identified as having consistently higher Sr CRs (337, 340, 341, 342 and 344) and four cultivars were identified as having consistently higher Cs CRs (335, 338, 345 and 348).

Influence of study size on species-wide variation in Cs and Sr concentration ratios

It is likely that inter-cultivar variation will increase with increasing numbers of cultivars in an experiment, until the full genetic range has been represented. To test this, inter-varietal variation was calculated for two randomly selected cultivars. This was repeated in increments of two (4 cultivars, 6 cultivars, 8 cultivars etc....) to the maximum population size (max = 189 cultivars). This process was iterated 100 times and the means of these iterations were calculated and plotted (Cs: Fig 5; Sr: Fig 6). Simulations were not performed for Italian ryegrass and tall fescue, as there were too few cultivars for the simulations to be meaningful. All the curves derived from these simulations, apart from hybrid ryegrasses grown in Edinburgh (Fig 6), appear to be approaching a plateau at which the inter-cultivar variation would reach a maximum, which suggests that we have sampled nearly enough cultivars to be close to the full range. However, it is not possible to estimate this plateau accurately as the curves are constrained by the number of cultivars included in the experiments.




Fig 5. Inter-cultivar variation in caesium concentration ratios for 2 to maximum number of cultivars. Mean value of 100 simulations is shown. <https://doi.org/10.1371/journal.pone.0176040.g005>




Fig 6. Inter-cultivar variation in strontium concentration ratios for 2 to maximum number of cultivars. Mean value of 100 simulations is shown. <https://doi.org/10.1371/journal.pone.0176040.g006>

Discussion

The rationale of the study

Despite the importance of the soil-plant-milk pathway to human radioactive dose following nuclear accidents (e.g. [31]), and the agricultural importance of fodder crops (e.g. temporary grasslands cover nearly 1.4 million ha in the UK; [32]), research into exploiting variation in radiocaesium and radiostrontium uptake by different plant cultivars has focussed mainly on human food crop species [12]. Of these studies, most included fewer than 20 cultivars [12], exceptions being Penrose *et al.*, 2016 [33] (70 cultivars; *Brassica oleracea*) Ohmori *et al.*, [34] (85; *Oryza sativa*) and Moiseev *et al.*, [35] (99 cultivars; *Pisum sativum*). Therefore, to our knowledge, this study, where a total of 397 cultivars from four common forage grass species (hybrid ryegrass (*Lolium perenne* L. x *Lolium multiflorum* Lam.; 101 cultivars), perennial ryegrass (*Lolium perenne* L.; 269 cultivars), Italian ryegrass (*Lolium multiflorum* Lam.; 17 cultivars) and tall fescue (*Festuca arundinacea* Shreb.; 10 cultivars) is the largest study to investigate inter-cultivar variation in Cs and Sr uptake by plants.

Cultivars with consistently low Cs and Sr concentration ratios

We found three hybrid ryegrass cultivars that had consistently low Cs concentration ratios between four harvests over two years at two sites. We also found two cultivars that had consistently low Sr concentration ratios and one cultivar that had consistently low Cs and Sr concentration ratios, despite Cs and Sr having very different uptake mechanisms.

Concentration ratios were found to be lower in spring cuts than in summer cuts, which was consistent with findings of Salt *et al.* [36], who found almost 10-fold difference in stable Cs concentrations in bent grass (*Agrostis capillaris*) between both immature and mature leaves cut in mid-July and end of August. Seasonal effects on Cs uptake have been attributed to temperature, rainfall or soil moisture [37], which of these (or a combination) is responsible for the seasonal variation observed in this study is not clear. More research is required to elucidate the effect of these in UK conditions.

Inter-species variation in Cs and Sr

Our findings suggest that substitution of one forage grass species by another 'safer' species with a lower CR [10], could be a remediation strategy for contaminated areas. Tall fescue cultivars were found to have lower Cs CRs (0.04–0.07) and Sr CRs (0.88–1.15) than most hybrid ryegrass, perennial ryegrass and Italian ryegrass cultivars. We therefore, propose that tall fescue could be considered a good substitute for *Lolium* forage grasses following a contamination incident. However, tall fescue is known to have lower dry matter digestibility and voluntary intake in comparison to *Lolium* species, although percentage dry matter and crude protein have been found to be similar for tall fescue and perennial ryegrass and dry matter yield has been found to be higher than perennial ryegrass [38]. Therefore, the effect of using tall fescue instead of *Lolium* for forage on agronomic factors such as yield and quality parameters such as digestibility need to be evaluated prior to implementation of this as a remediation strategy.

Optimal number of cultivars

Our results suggest that increasing numbers of cultivars generally occurs concurrently with an increase in inter-cultivar variation, which implies that as the number of cultivars in an experiment increases, the likelihood of finding a larger variation between cultivars also increases. The inter-cultivar variations found in these experiments (Cs = up to 14-fold; Sr = up to 4.4-fold) were generally higher than those found in Alexakhin [39] (^{137}Cs inter-cultivar variation = 1.4–4.5 fold), where between 4–15 cultivars from 10 plant species were studied and Penrose *et al.* [12] (Cs inter-cultivar variation = 1.8; Sr inter-cultivar variation = 2.0) in which 115 experiments including between 2 and 28 cultivars (Cs) and 2 and 99 cultivars (Sr) from over 50 species were collated. Previous studies that have included a greater number of cultivars, for example Ohmori *et al.*, (2014) [34], which investigated variation in ^{137}Cs activity concentrations among 85 rice (*Oryza sativa*) cultivars and Penrose *et al.*, (2016) [33], which investigated variation in ^{137}Cs and ^{90}Sr concentration ratios in c. 70 *Brassica oleracea* cultivars have found larger inter-cultivar variation (Ohmori *et al.*, 2014 [34] = c.10-fold; Penrose *et al.*, 2016 [33]; ^{137}Cs = 35-fold, ^{90}Sr = 23-fold).

In this study, the relationship between the number of cultivars and the inter-cultivar variation in Cs and Sr concentration ratios was different for each of the species-location combinations, suggesting that the relationship is unique to each species in each location. More research on linkages between crop genetics, soil characteristics and environmental or management conditions is needed for this to be further elucidated. Whilst this study included a large number (29–189) of hybrid ryegrass and perennial ryegrass cultivars, it is likely that we would have observed larger inter-cultivar variation in Cs and Sr concentration if more cultivars had been studied. It is therefore recommended that studies in the future regarding inter-cultivar variation in Cs and Sr include as many cultivars as possible.

Inter-cultivar variation in Cs and Sr

Our results suggest that cultivar substitution could be an effective remediation strategy for contaminated grasslands, and could reduce transfer of Cs to forage by up to 14-fold, and Sr by up to approximately 4-fold. To our knowledge, only two other studies (Øhlenschläger & Gissel-Nielsen, [14]; [15]) have considered variation in Cs uptake between forage grass cultivars. Both papers, in which six studies were carried out, included only two perennial ryegrass cultivars (Darbor and Partoro). The inter-cultivar variation in ^{137}Cs uptake ranged from 1.0–1.3 in their studies (Øhlenschläger & Gissel-Nielsen, [14]; [15]), which was lower than the inter-cultivar variation in Cs found in our study (up to 14-fold). It is likely that the greater variation between cultivars in our study is due to the greater number of cultivars included.

The maximum reduction in Cs and Sr (Cs = up to 14-fold; Sr = up to 4.4-fold) by planting 'safer' cultivars with lower CRs are similar to—if not greater than—the existing mechanical and chemical soil amendments currently used for remediation of contaminated agricultural land. Mechanical soil amendment techniques such as ploughing can reduce radiocaesium and radiostrontium transfer to plants by up to 4-fold, whilst soil amendments such as liming or the addition of organic matter can reduce transfer by 1.5–4.0 fold and 1.3–3.0 fold, respectively [7]. Application of mineral fertilisers can reduce transfer of radiocaesium by 1.3–16-fold [40] and radiostrontium by up to 2.0-fold [41] depending on the soil type. As noted above, for Cs there are additionally effective binders which can be administered to animals to reduce the uptake of radiocaesium from the gut and hence transfer to milk and/or meat [8], though there are no feasible binders for Sr. Whilst dietary calcium levels can be manipulated to reduce radiostrontium transfer to milk it is likely that a maximum reduction of radiostrontium of at least c. two-fold could be achieved in most farming systems without exceeding recommended dietary calcium intakes [40].

Cultivar-selection could be used in conjunction with other existing strategies to add additional reductions in Cs and Sr transfer to food products. Farmers are advised to reseed 10–15% of their grassland per year, with priority given to weed-affected fields and fields with reduced yield. Therefore, reseeding early following a nuclear incident would not be very different from normal practice. Furthermore, it is likely that following an incident low-Cs and low-Sr seeds would be subsidised, reducing the cost to the farmer of reseeding. Reseeding involves both ploughing and fertilisation, and therefore these three remediation strategies used together may well reduce Cs and Sr transfer to livestock further. It is also likely that cultivar, or species substitution of forage grasses will be more acceptable to stakeholders than, for instance, the use of feed additives.

Conclusions

Using existing ongoing grass breeding experiments allowed us to undertake the largest and most comprehensive examination of inter-cultivar variation in Cs and Sr uptake in plants to date over two sites, sampled at multiple time points over two years.

All tall fescue cultivars had lower Cs and Sr CRs than the majority of cultivars from the other three species, and therefore could be a candidate species for species substitution following a contamination incident.

Three (from 17) hybrid ryegrass cultivars were found to have consistently low Cs concentration ratios, two cultivars were found to have consistently low Sr concentration ratios, one cultivar was found to have lower Cs and Sr concentration ratios over time and between sites. This is the first time, to the authors' knowledge, that forage grass cultivars with consistently lower Cs and Sr concentration ratios have been identified. The identification of cultivars with consistently lower concentration ratios suggests that substitution of cultivars with high concentration ratios for cultivars with lower concentration ratios could be an effective remediation strategy in contaminated grasslands. However, the sites were relatively similar in terms of soil characteristics and climate, and therefore cultivars with low concentration ratios should be tested for consistency over sites with more diverse environmental conditions.

We have found that inter-cultivar variation increases with increasing numbers of cultivars included in the experiment. We suggest that future research calculating inter-cultivar variation in Cs and Sr endeavours to include the largest feasible number of cultivars in order to increase the chances of reaching the maximum inter-cultivar variation.

Supporting information

S1 Fig. Ratio of the mean concentration of 28 elements in unwashed samples to the mean concentration of 28 elements in washed samples.

The black line indicates a 1:1 ratio, where there is no difference in concentration between the washed and unwashed samples.
<https://doi.org/10.1371/journal.pone.0176040.s001>
 (TIFF)

S1 Table. Washing test and ICP-MS methods.

<https://doi.org/10.1371/journal.pone.0176040.s002>
 (XLSX)

S2 Table. Experiment number, plot area, species and variety number, sowing year, sowing and harvest date and K, Ca, Rb, Sr and Cs concentration (mg kg⁻¹).

<https://doi.org/10.1371/journal.pone.0176040.s003>
 (XLS)

S3 Table. Minimum and maximum concentration ratios (CRs) and inter-varietal variation (IVV; maximum CR/minimum CR) for 17 hybrid ryegrass cultivars grown in Aberystwyth and Edinburgh.

<https://doi.org/10.1371/journal.pone.0176040.s004>
 (XLSX)

S4 Table. Rankings (R) of 17 hybrid cultivars grown in both Aberystwyth and Edinburgh.

The lowest ranking (1) denotes the cultivar with the lowest concentration ratio, the highest ranking (17) refers to the cultivar with the highest concentration ratio.

<https://doi.org/10.1371/journal.pone.0176040.s005>
 (XLSX)

S5 Table. Mean concentration of Cs, Sr, Ca, Mg and K (mg kg⁻¹) in perennial ryegrass, Italian ryegrass, hybrid ryegrass and tall fescue.

Results are shown for Aberystwyth and Edinburgh.
<https://doi.org/10.1371/journal.pone.0176040.s006>
 (XLSX)

S6 Table. Mean concentration of Cs, Sr, Ca, Mg and K (mg kg⁻¹) for perennial ryegrass, Italian ryegrass, hybrid ryegrass and tall fescue in each field.

<https://doi.org/10.1371/journal.pone.0176040.s007>
 (XLSX)

Acknowledgments

The authors would like to thank Scott Young (University of Nottingham) for ICP-MS analyses, Raymond Davies and Dylan Jones (IBERS) for their help harvesting the samples, and Peter Henrys (CEH Lancaster) for his advice on statistical methods. Beth Penrose was supported by a studentship from the UK Natural Environment Research Council (NE/K500951/1).

Author Contributions

Conceptualization: BP NAB NMJC JAL MRB.

Data curation: BP.

Formal analysis: BP.

Funding acquisition: NB NMJC MB.

Investigation: BP JAL RT.

Methodology: BP NAB NMJC JAL RT MRB.

Software: BP.

Visualization: BP.

Writing – original draft: BP NAB NMJC JAL MRB.

Writing – review & editing: BP NAB NMJC JAL RT MRB.

References

1. Shaw G. Radioactivity in the terrestrial environment. Oxford: Elsevier; 2007.
2. Wright SM, Smith JT, Beresford NA, Scott WA. Monte-Carlo prediction of changes in areas of west Cumbria requiring restrictions on sheep following the Chernobyl accident. *Radiat Environ Biophys.* 2003;42: 41–47. pmid:12684828
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
3. IAEA. Environmental Consequences of the Chernobyl Accident and Their Remediation: Twenty Years of Experience. International Atomic Agency; 2006.

4. Nakanishi TM, Tanoi K. Agricultural Implications of the Fukushima Nuclear Accident [Internet]. Japan: Springer; 2013. Available: [10.1007/978-4-431-54328-2](https://doi.org/10.1007/978-4-431-54328-2)
5. Molchanova I, Mikhailovskaya L, Antonov K, Pozolotina V, Antonova E. Current assessment of integrated content of long-lived radionuclides in soils of the head part of the East Ural Radioactive Trace. *J Environ Radioact.* 2014;138: 238–248. pmid:25262151
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
6. Kolosha V. Atlas Ukraine Radioactive Contamination. Ministry of Ukraine of emergencies and affairs of population protection from the consequences of Chernobyl catastrophe. Kiev; 2008.
7. IAEA. Guidelines for Remediation Strategies to Reduce the Radiological Consequences of Environmental Contamination. International Atomic Agency; 2012.
8. Beresford NA, Fesenko S, Konoplev A, Skuterud L, Smith JT, Voigt G. Thirty years after the Chernobyl accident: What lessons have we learnt? *J Environ Radioact.* 2016;157: 77–89. pmid:27018344
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
9. Beresford N., Mayes R., Colgrove P., Barnett C., Bryce L, Dodd B., et al. A comparative assessment of the potential use of alginates and dietary calcium manipulation as countermeasures to reduce the transfer of radiostrontium to the milk of dairy animals. *J Environ Radioact.* 2000;51: 321–334.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
10. White PJ, Swarup K, Escobar-Gutiérrez AJ, Bowen HC, Willey NJ, Broadley MR. Selecting plants to minimise radiocaesium in the food chain. *Plant Soil.* 2003;249: 177–186.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
11. Beresford NA, Barnett C., Howard BJ, Rantavaara A, Rissanen K, Reales N, et al. EURANOS Compendium of Countermeasures for the Management of Food Production Systems [Internet]. Available: <http://www.euranos.fzk.de>.
12. Penrose B, Beresford NA, Broadley MR, Crout NMJ. Inter-variety variation in caesium and strontium uptake by plants: a meta-analysis. *J Environ Radioact.* 2015;139: 103–117. pmid:25464046
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
13. Howard BJ, Beresford NA, Nisbet A, Cox G, Oughton DH, Hunt J, et al. The STRATEGY project: decision tools to aid sustainable restoration and long-term management of contaminated agricultural ecosystems. *J Environ Radioact.* 2005;83: 275–295. pmid:15905001
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
14. Oehlenschläger M, Gissel-Nielsen G. Transfer of radiocaesium to barley, rye grass and pea [Internet]. Risoe National Lab.; 1989. Report No.: RISO-M—2831. Available: http://inis.iaea.org/Search/search.aspx?orig_q=RN:21042934
15. Ølenschläger M, Gissel-Nielsen G. Differences in the Ability for Barley and Rye Grass Varieties to Absorb Caesium through the Roots. *Acta Agric Scand.* 1991;41: 321–328.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
16. Minson DJ. Preface. Forage in Ruminant Nutrition. Academic Press; 1990. pp. xi–xii. Available: <http://www.sciencedirect.com/science/article/pii/B9780124983106500055>
17. Slepser DA, Vogel KP, Asay KH, Mayland HF. Using plant breeding and genetics to overcome the incidence of grass tetany. *J Anim Sci.* 1989;67: 3456–3462. pmid:2693420
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
18. Nguyen HT, Slepser DA. Genetic variability of mineral concentrations in *Festuca arundinacea* Schreb. *TAG Theor Appl Genet Theor Angew Genet.* 1981;59: 57–63.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
19. Reeder LR, Nguyen HT, Slepser DA, Brown JR. Genetic Variability of Mineral Concentrations in Tall Fescue Grown under Controlled Conditions. *Crop Sci.* 1986;26: 514.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
20. Broadley MR, White PJ. Some elements are more equal than others: soil-to-plant transfer of radiocaesium and radiostrontium, revisited. *Plant Soil.* 2012;355: 23–27.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
21. Hides D, Thomas T. Variation in the magnesium content of grasses and its improvement by selection. *J Sci Food Agric.* 1981;32: 990–991.
[View Article](#) • [PubMed/NCBI](#) • [Google Scholar](#)
22. Mayland HF, Slepser DA. Developing a tall fescue for reduced grass tetany risk. XVII International Grassland Congress: summaries of invited (spoken) papers and poster papers. Palmerston North: The Congress; 1993. pp. 1095–1096.
23. Emmett BA, Frogbrook ZL, Chamberlain P, Griffiths R, Pickup R, Poskitt J, et al. CS Technical Report No.3/07 Soils Manual. Natural Environment Research Council; 2008.
24. British Geological Survey. UKSO Soils map viewer [Internet]. 2015 [cited 13 Sep 2015]. Available: <http://mapapps2.bgs.ac.uk/ukso/home.html?>

25. Stuanes AO, Ogner G, Opem M. Ammonium nitrate as extractant for soil exchangeable cations, exchangeable acidity and aluminum. *Commun Soil Sci Plant Anal.* 1984;15: 773–778.
View Article • PubMed/NCBI • Google Scholar
26. R Core Team. R: A language and environment for statistical computing [Internet]. Vienna: R Foundation for Statistical Computing; 2013. Available: <http://www.R-project.org/>
27. International Atomic Energy Agency (IAEA). Quantification of Radionuclide Transfer in Terrestrial and Freshwater Environments for Radiological Assessments [Internet]. Vienna: INTERNATIONAL ATOMIC ENERGY AGENCY; 2009. Available: <http://www-pub.iaea.org/books/IAEABooks/8103/Quantification-of-Radionuclide-Transfer-in-Terrestrial-and-Freshwater-Environments-for-Radiological-Assessments>
28. Beresford N. A., Wright S. M. Non-linearity in radiocaesium soil to plant transfer: Fact or fiction? *Radioprotection.* 2005;40: S67–S72.
View Article • PubMed/NCBI • Google Scholar
29. Bates D, Mächler M, Bolker B, Walker S. Fitting Linear Mixed-Effects Models Using lme4. *J Stat Softw.* 2015;67.
View Article • PubMed/NCBI • Google Scholar
30. Peverill K, Sparrow L, Reuter D. Soil analysis and interpretation manual. CSIRO Publishing; 2001.
31. Beresford NA, Voigt G, Wright SM, Howard BJ, Barnett CL, Prister B, et al. Self-help countermeasure strategies for populations living within contaminated areas of Belarus, Russia and Ukraine. *J Environ Radioact.* 2001;56: 215–239. pmid:11446120
View Article • PubMed/NCBI • Google Scholar
32. Department for Environment, Food and Rural Affairs. 2015 [cited 29 Jan 2016]. Available: <https://www.gov.uk/government/statistics/agriculture-in-the-united-kingdom-2014>
33. Penrose B, Johnson née Payne KA, Arkhipov A, Maksimenko A, Gaschak S, Meacham MC, et al. Inter-cultivar variation in soil-to-plant transfer of radiocaesium and radiostrontium in Brassica oleracea. *J Environ Radioact.* 2016;155–156: 112–121. pmid:26945429
View Article • PubMed/NCBI • Google Scholar
34. Ohmori Y, Inui Y, Kajikawa M, Nakata A, Sotta N, Kasai K, et al. Difference in cesium accumulation among rice cultivars grown in the paddy field in Fukushima Prefecture in 2011 and 2012. *J Plant Res.* 2014;127: 57–66. pmid:24338062
View Article • PubMed/NCBI • Google Scholar
35. Moiseev I, Tikhomirov F, Rerikh L. Effect of Cultivar Characteristics on Cesium-137 and Potassium Accumulation in Wheat and Pea Yields. *Vestn Mosk Univ.* 1977;3: 105–109.
View Article • PubMed/NCBI • Google Scholar
36. Salt CA, Kay JW, Jarvis KE. The influence of season and leaf age on concentrations of radiocaesium (¹³⁷Cs), stable caesium (¹³³Cs) and potassium in *Agrostis capillaris*. *Environ Pollut.* 2004;130: 359–369. pmid:15182969
View Article • PubMed/NCBI • Google Scholar
37. Ehlken S, Kirchner G. (1996) Seasonal variations in soil-to-grass transfer of fallout strontium and cesium and of potassium in North German soils. *J Environ Radioact.* 1996;33: 147–181.
View Article • PubMed/NCBI • Google Scholar
38. Cougnon M, Baert J, Van Waes C, Reheul D. Performance and quality of tall fescue (*Festuca arundinacea* Schreb.) and perennial ryegrass (*Lolium perenne* L.) and mixtures of both species grown with or without white clover (*Trifolium repens* L.) under cutting management. *Grass Forage Sci.* 2014;69: 666–677.
View Article • PubMed/NCBI • Google Scholar
39. Alexakhin RM. Countermeasures in agricultural production as an effective means of mitigating the radiological consequences of the Chernobyl accident. *Sci Total Environ.* 1993;137: 9–20.
View Article • PubMed/NCBI • Google Scholar
40. Prister B, Loshchilov N, Perepelyatnikova L, Perepelyatnikov G, Bondar P. Efficiency of measures aimed at decreasing the contamination of agricultural products in areas contaminated by the Chernobyl NPP accident. *Sci Total Environ.* 1992;112: 79–87. pmid:1574707
View Article • PubMed/NCBI • Google Scholar
41. Fesenko SV, Alexakhin RM, Balonov MI, Bogdevitch IM, Howard BJ, Kashparov VA, et al. An extended critical review of twenty years of countermeasures used in agriculture after the Chernobyl accident. *Sci Total Environ.* 2007;383: 1–24. pmid:17573097
View Article • PubMed/NCBI • Google Scholar