

Effects of Maternal Basking and Food Quantity during Gestation Provide Evidence for the Selective Advantage of Matrotrophy in a Viviparous Lizard

Keisuke Itonaga, Susan M. Jones, Erik Wapstra*

School of Zoology, University of Tasmania, Hobart, Tasmania, Australia

Abstract

The evolution of matrotrophy (i.e., direct supply of nutrients by the mother during gestation) may be associated with high maternal energy availability during gestation. However, we lack knowledge about the selective advantages of matrotrophic viviparity (live-bearing) in reptiles. In reptiles, the interaction between body temperature and food intake affect maternal net energy gain. In the present study, we examined the effects of basking and food availability (2 by 2 factorial design) during gestation on offspring phenotype in a matrotrophic viviparous lizard (*Pseudemoia entrecasteauxii*). Subsequently, we investigated if the maternal effects were context-dependent using offspring growth rate as an indicator of the adaptive significance of matrotrophy. Offspring were exposed either to the same thermal conditions as their mothers experienced or to thermal conditions different from those experienced by their mothers. We provide the first evidence that an interaction between maternal thermal and maternal food conditions during gestation strongly affects offspring phenotype, including date of birth, body size and performance ability, which affect offspring fitness. Offspring growth rate was dependent on offspring thermal conditions, but was not influenced by maternal effects or offspring sex. Matrotrophic viviparity provided gravid females with the means to enhance offspring fitness through greater energetic input to offspring when conditions allowed it (i.e., extended basking opportunity with high food availability). Therefore, we suggest that selective advantages of matrotrophic viviparity in *P. entrecasteauxii* may be associated with high maternal energy availability during gestation.

Citation: Itonaga K, Jones SM, Wapstra E (2012) Effects of Maternal Basking and Food Quantity during Gestation Provide Evidence for the Selective Advantage of Matrotrophy in a Viviparous Lizard. PLoS ONE 7(7): e41835. doi:10.1371/journal.pone.0041835

Editor: Jeff Demuth, University of Texas Arlington, United States of America

Received: September 6, 2011; **Accepted:** June 29, 2012; **Published:** July 26, 2012

Copyright: © 2012 Itonaga et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Funding was provided by Ecological Society of Australia Student Research Awards (<http://www.ecolsoc.org.au/What%20we%20do/Prizes/StudentResearchPrize.html>). EW thanks the Australian Research Council for support. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: Erik.Wapstra@utas.edu.au

Introduction

Maternal effects can include non-genetic effects that influence offspring phenotype. They are recognised as one of the most important components of evolutionary ecology [1]. Since the early 20th century, a large number of studies have documented maternal effects in both plants and animals (e.g., insects: [2]; herbs: [3]; fish: [4]; reptiles: [5]). However, the adaptive significance of maternal effects is still an unresolved issue in evolutionary biology [6–8]. This is because maternal effects can occur without any important ecological implications [9,10] or conversely may have positive or negative impacts on offspring fitness [6].

Despite the uncertain benefits of maternal effects, the evolution of viviparity (live-bearing) in squamate reptiles has been linked to maternal effects. Shine [11] suggested that the prolonged maternal control over embryonic developmental conditions in the uterus may provide selective advantages for offspring of viviparous reptiles (i.e., the Maternal Manipulation Hypothesis). Reptiles are ectothermic, so environmental conditions (especially temperature) strongly affect physiological processes. For example, embryonic temperature during embryogenesis has been shown to affect offspring phenotype in many species [12–14]. In viviparous reptiles, the temperature of embryonic development is controlled by maternal thermoregulation (reviewed in [15]), and this

maternal control during gestation can enhance offspring fitness (e.g., [5,16,17]).

In viviparous animals, offspring fitness may involve a number of other maternal factors. For example, the magnitude of placental nutrient support during embryogenesis strongly affects offspring fitness in matrotrophic viviparous animals which utilise a source supplied by the mother to nourish embryos during pregnancy (reviewed in [18]). Indeed, Trexler and DeAngelis [19] predict that matrotrophic viviparity is more likely to evolve when food availability during gestation is high and constant. This model developed for fish has largely been ignored in studies of the evolution of viviparity and subsequent evolution of matrotrophy in reptiles. Past studies generally concentrated on the key selective force of maternal effects mediated through temperature [11], however maternal food availability may be important [20–22]. This is because in reptiles net energy gain is strongly associated with the interaction between body temperature and food intake [23,24]. However, no study has investigated such interaction-induced selective advantages in matrotrophic viviparous reptiles.

Context-dependent or anticipatory maternal effects can enhance offspring fitness [6,7]. They occur when there is a correlation between maternal and offspring environments, such that the reproducing females predict the offspring environment in response to their current environment and adjust offspring

phenotype accordingly [6,7,25]. In reptiles, growth rate is an important component of offspring fitness [26–28], and benefits of growth rates can be dependent on the postnatal environment [29]. Therefore, offspring growth rate may be a reliable indicator as to whether mothers adjust offspring growth rate to postnatal environments. There is some evidence that maternal thermal conditions during gestation can influence offspring growth rate in viviparous reptiles [13,30]. So, if context-dependent maternal effects have evolved in viviparous reptiles, it is possible that gravid females adjust offspring growth rate in response to maternal thermal (or other environmental) conditions, and that this enhances offspring, and their own, fitness.

In this study, we focused on how the interaction between maternal thermal and maternal food conditions (i.e., maternal net energy availability) during gestation affects offspring phenotype in a matrotrophic viviparous reptile. Matrotrophy may benefit offspring development (and fitness) when maternal net energy is high during gestation [19,31], by enhancing fitness-related offspring phenotypic traits such as date of birth, body size, performance ability and amount of fat reserves [5,20,32–34]. Conversely, matrotrophy may be costly to offspring development (and fitness) when maternal net energy is low during gestation [35,36]. In such situations, however, matrotrophic females may display strategies such as bet-hedging (i.e., variation in energy allocation among embryos) to prevent the costs of matrotrophy from outweighing its benefits in terms of maternal fitness [19,37,38].

We addressed these questions using southern grass skinks, *Pseudemoia entrecasteauxii*. They are small matrotrophic viviparous skinks [snout-vent length (SVL) 40–60 mm] and their nutrient provisioning for embryonic development is roughly half from the yolk and half via the placenta [21,39]. This is an ideal model species to investigate the evolution of matrotrophic reproduction in reptiles; we have previously demonstrated the importance of transfer of nutrients, organic substances and hormones across the placenta [22,40,41] during embryo development, and here explore the importance of variation in two key environmental factors that affect female energy availability during gestation (temperature and food availability). This species is restricted to cold-temperate regions where food and thermal availability vary annually and seasonally [42–45]. In addition, females increase basking behaviour and maintain high feeding rates during gestation [22,46] suggesting both maternal body temperature and energy gain are important to offspring development. We manipulated maternal basking (i.e., female body temperature) and maternal food availability (2 by 2 factorial design) during gestation and then measured date of birth, offspring body size, offspring performance ability, offspring fat reserves and variation in within-clutch offspring size (i.e., indicator of a bet-hedging strategy) to examine the effects of the interaction between maternal thermal and maternal food conditions during gestation on offspring. Subsequently, we measured offspring growth rate using reciprocal transplant experiments (manipulating thermal conditions only) to examine the potential for context-dependent maternal effects (e.g., [22,47]). A similar design was used previously by Swain and Jones [32] to explore facultative versus obligate placentotrophy in a largely non-matrotrophic unrelated species (*Niveoscincus metallicus*); here we explore a larger question about the selective advantages of matrotrophic viviparity in reptiles.

Results

Maternal and offspring characteristics

There were no differences in initial maternal SVL among treatments ($F_{3, 159} = 1.20$, $P = 0.3126$) (Table 1). Although all

females grew during gestation, maternal growth rate (mm/day) was significantly affected by the interaction between maternal basking and maternal food availability during gestation (Table 1 and 2). Post-hoc tests [Ryan-Einot-Gabriel-Welsch multiple range tests (REGWQ tests)] showed that females given extended basking with high food availability during gestation had a higher growth rate than females in other gestation conditions (Table 1). Although maternal growth rates varied between maternal treatments, there was no variation in postpartum maternal SVL between treatments (Table 1 and 2), presumably reflecting differences in gestation length between treatments (see below).

There was no effect of maternal basking availability during gestation and no interaction between maternal basking and maternal food availability during gestation on postpartum maternal body mass. However, the main effect of maternal food availability during gestation was significant (Table 2). Females with high food availability during gestation were heavier postpartum than females with low food availability (Table 1). Maternal treatments did not affect clutch size (Table 2). There was no interaction between maternal basking and maternal food availability on relative clutch mass (RCM), but the main effects of maternal basking and maternal food availability during gestation were significant (Table 2). Females given 12 hours basking during gestation showed significantly higher RCM than females given four hours basking during gestation. Similarly, females with low food availability during gestation showed significantly higher RCM than females with high food availability during gestation, presumably reflecting differences in maternal postpartum body mass (Table 1).

Maternal food availability during gestation significantly affected within-clutch variation in offspring size (i.e., CV), but maternal basking opportunity during gestation and the interaction between maternal basking and maternal food availability during gestation had no effect (Table 2). Within-clutch variation in offspring size was lower in females that had high food availability (6.22 ± 0.60) than in females that had low food availability during gestation (8.14 ± 0.68) (Table 1). Some females produced both fully-developed offspring and premature offspring in the same clutch. Six cases of this were observed in females given 12 hours basking with low food availability during gestation, four cases in females given four hours basking with high food availability during gestation, and seven cases in females given four hours basking with low food availability during gestation. Overall, we observed more premature offspring in females with low food availability during gestation (Table 1).

Date of birth was affected by an interaction between maternal basking and maternal food availability during gestation (Table 2): REGWQ tests showed that females given 12 hours basking with high food availability during gestation gave birth about five days earlier than females given 12 hours basking with low food availability during gestation. Furthermore, mean birth date of all females in 12 hours basking treatments was about 55 days earlier than for females in four hours basking treatments (Fig. 1a and 1b).

Most offspring phenotypic traits including SVL, mass and fat reserves were significantly affected by the interaction between maternal basking and maternal food availability during gestation (Table 2). REGWQ tests showed that females given 12 hours basking with high food availability during gestation produced offspring with larger SVL, heavier mass and larger fat reserves relative to body mass than females in other gestation conditions (Table 1 and Fig. 2a, 2b, 2d). Females given 4 hours basking with high food availability during gestation also produced offspring with larger SVL and heavier mass than females given 4 hours basking with low food availability during gestation or females given 12 hours basking with low food availability during gestation

Table 1. Characteristics of maternal and offspring *Pseudemoia entrecasteauxii* under four gestational regimes.

		12 hours basking	12 hours basking	4 hours basking	4 hours basking
		+high food	+low food	+high food	+low food
Maternal characteristics	Total sample size	40	40	40	40
	Number of females giving birth	37	36	36	36
	Initial snout-vent length (mm)	44.93±0.76	46.26±0.90	46.04±0.72	44.92±0.43
	Postpartum snout-vent length (mm)	47.32±0.64	47.19±0.90	47.68±0.64	46.12±0.47
	Growth rate during gestation (mm/day)	0.054±0.005	0.019±0.003	0.016±0.002	0.011±0.001
	Postpartum body mass (g)	2.07±0.06	1.53±0.08	2.10±0.08	1.65±0.05
	Number of offspring	106	107	102	105
	Number of premature offspring	0	29	7	30
	Number of stillborn offspring	0	2	8	9
	Clutch size	2.84±0.17	2.94±0.20	2.83±0.16	2.92±0.18
	Relative clutch mass (unit-less: g/g)	0.26±0.01	0.30±0.02	0.22±0.01	0.26±0.01
	Within-clutch variation in offspring size (coefficient of variation)	5.59±0.02	7.28±0.79	6.93±1.14	9.00±1.10
	Offspring characteristics	Snout-vent length (mm)	20.98±0.50	19.87±0.16	20.50±0.12
Body mass (mg)		195.31±3.86	159.89±4.42	168.68±3.95	156.50±4.92
Sprint speed (m s ⁻¹)		0.46±0.01	0.38±0.01	0.40±0.01	0.37±0.01
Dry fat reserves relative to body mass (Student residual)		0.53±0.20 (35)	-0.40±0.08 (30)	0.01±0.21 (30)	-0.19±0.16 (30)

Characteristics of maternal *Pseudemoia entrecasteauxii* including snout-vent length, growth rate during gestation, body mass, clutch size, relative clutch mass and within-clutch variation in offspring size, and characteristics of offspring *Pseudemoia entrecasteauxii* from females given combinations of extended and restricted basking opportunities and food supply during gestation including snout-vent length, body mass, sprint speed and fat reserves relative to body mass at birth. Values are means ± S.E. (n).

doi:10.1371/journal.pone.0041835.t001

(Table 1 and Fig. 2a, 2b, 2d). Offspring sprint speed was influenced by both maternal basking and maternal food availability during gestation, but there was no interaction (Table 2). Offspring from females given 12 hours basking opportunities during gestation sprinted faster than offspring from females given four hour basking opportunities during gestation. Offspring from females with high food availability during gestation sprinted faster

than offspring from females with low food availability during gestation (Table 1 and Fig. 2c).

Assessing context-dependent maternal effects based on offspring growth rates

Offspring growth rate was affected by offspring basking availability, but not by maternal gestation conditions, offspring

Table 2. The results of statistical analyses for maternal and offspring characteristics.

		Basking condition	Food supply	Basking*Food
Maternal characteristics	Postpartum snout-vent length	$F_{1, 142} = 0.03, P = 0.5868$	$F_{1, 142} = 1.67, P = 0.1989$	$F_{1, 142} = 1.60, P = 0.2076$
	Postpartum body mass	$F_{1, 142} = 2.83, P = 0.0948$	$F_{1, 142} = 54.31, P < 0.0001$	$F_{1, 142} = 1.38, P = 0.2413$
	Growth rate during gestation	$F_{1, 142} = 52.80, P < 0.0001$	$F_{1, 142} = 43.14, P < 0.0001$	$F_{1, 142} = 20.74, P < 0.0001$
	Clutch size	$F_{1, 142} = 0.10, P = 0.7567$	$F_{1, 142} = 0.00, P = 0.9711$	$F_{1, 142} = 0.04, P = 0.8463$
	Relative clutch mass	$F_{1, 128} = 5.05, P = 0.0264$	$F_{1, 128} = 7.73, P = 0.0063$	$F_{1, 128} = 0.12, P = 0.7309$
	Within-clutch variation in offspring size	$F_{1, 120} = 2.30, P = 0.1317$	$F_{1, 120} = 7.09, P = 0.0088$	$F_{1, 120} = 0.02, P = 0.8887$
	Date of birth	$F_{1, 134} = 18.41, P < 0.0001$	$F_{1, 134} = 21.63, P < 0.0002$	$F_{1, 134} = 11.44, P = 0.0009$
Offspring characteristics	Snout-vent length	$F_{1, 140} = 2.30, P = 0.1313$	$F_{1, 140} = 27.24, P < 0.0001$	$F_{1, 140} = 4.25, P = 0.0411$
	Body mass	$F_{1, 137} = 15.80, P = 0.0001$	$F_{1, 137} = 34.05, P < 0.0001$	$F_{1, 137} = 7.98, P = 0.0054$
	Sprint speed	$F_{1, 145} = 8.82, P = 0.0035$	$F_{1, 145} = 17.29, P < 0.0001$	$F_{1, 145} = 1.49, P = 0.2236$
	Fat reserves relative to body mass	$F_{1, 121} = 0.73, P = 0.3946$	$F_{1, 121} = 10.26, P = 0.0017$	$F_{1, 121} = 4.29, P = 0.0405$

Summary of the results of statistical analyses of the effects of maternal basking and food availability (combinations of extended and restricted basking opportunities and food supply) and their interaction during gestation on female and offspring in *Pseudemoia entrecasteauxii*. Significant results are indicated in bold.

doi:10.1371/journal.pone.0041835.t002




Figure 1. Frequency of births (3 day increments) for *Pseudemoia entrecasteauxii*. White bars indicate high food availability during gestation, while black bars indicate low food availability during gestation. Graph A shows date of birth in offspring from females given extended basking availability during gestation. Graph B shows date of birth in offspring from females given restricted basking availability during gestation. Three day increments begin with the first recorded births on Dec 7 2007.
doi:10.1371/journal.pone.0041835.g001

sex or any interactions between maternal basking opportunity, maternal food availability, offspring basking opportunity and offspring sex (Table 3). Random choice of a single offspring for each clutch resulted in relatively even distribution of sexes among the treatments with no clear bias resulting from maternal treatments (27 males and 33 females from 12 hours of basking and 36 males and 24 females from 4 hours of basking). Offspring given 12 hours basking availability grew faster (0.056 ± 0.043 mm day⁻¹) than offspring given four hours basking availability (0.043 ± 0.002 mm day⁻¹) (Fig. 3).

Discussion

Our experiment provides the first evidence that maternal thermal conditions and maternal food conditions during gestation and importantly the interaction between them, strongly affect fitness related offspring phenotypic traits in a matrotrophic viviparous reptile. We showed that females given extended basking with high food availability during gestation produced offspring with earlier birth date, larger size, larger fat reserves relative to body mass, and faster sprint speed compared with females given other treatments. These effects may be explained by an acceleration of embryonic development due to increased maternal body temperature through extended basking opportunity [13,48,49], combined with increased net energy gain when food is abundant [23,24], providing for enhanced nutrient transport across the placenta to developing embryos. High nutrient support for embryonic development via the placenta may increase offspring body size and body condition involving an increase in




Figure 2. Effects of maternal gestation conditions on offspring *Pseudemoia entrecasteauxii*. A: snout-vent length (SVL), B: body mass, C: sprint speed and D: fat reserves relative to body mass at birth from females given combinations of extended and restricted basking opportunities and food supply during gestation. Values are means \pm S.E. ●: Offspring from females given high food availability during gestation. ○: Offspring from females given low food availability during gestation.
doi:10.1371/journal.pone.0041835.g002

both muscle [50] and fat reserves [51]; see Thompson and Stewart [52] for related arguments.

Date of hatching or birth, body size and locomotor performance ability are known to be important determinants of offspring fitness in a variety of taxa including reptiles (e.g., [5,33,53–57]). The potential fitness benefits related to offspring phenotype, such as early birth, large body size and high locomotor performance in reptiles include a higher survival over winter [56,58], a higher early-age survivorship [59,60], a higher predator avoidance [61,62] and a larger size at maturation [12,34]. Specifically, these effects are important to *P. entrecasteauxii* because their predator avoidance strategy is primarily sprinting [63,64] and because adult body size in both sexes is positively associated with reproductive success (i.e., courtship and fecundity) ([65]; in the present study: maternal initial SVL and clutch size showed a positive relationship, $r = 0.29$, $P < 0.0001$). Therefore, if offspring




Figure 3. Offspring growth rate over five weeks in *Pseudemoia entrecasteauxii* under the two thermal regimes. ●: Offspring from females given 12 hours basking opportunity with high food availability during gestation. ○: Offspring from females given 12 hours basking opportunity with low food availability during gestation. ■: Offspring from females given 4 hours basking opportunity with high food availability during gestation. □: Offspring from females given 4 hours basking opportunity with low food availability during gestation. Sample sizes among offspring given 12 hours basking opportunity per day are ● = 17, ○ = 13, ■ = 16 and □ = 14. Sample sizes among offspring given 4 hours basking opportunity per day are ● = 18, ○ = 17, ■ = 13 and □ = 12. Values are means ± S.E. Some overlapping points have been displaced laterally to aid clarity.
doi:10.1371/journal.pone.0041835.g003

size at birth is positively correlated with size at a later date and importantly at maturation as it is in other small Australian lizards [5,14,34,66], large offspring size at birth has an advantage in terms of future reproduction. However, large offspring size does not always have positive fitness effects; it can reduce survival rate when postnatal food is limited [29]. We found that large offspring from females given extended basking with high food availability during

gestation had large fat reserves relative to body mass. Even when feeding opportunities are limited in postnatal environments, large fat reserves may help these offspring to meet immediate crises. Thus, we suggest that high maternal net energy availability, (i.e., extended basking and abundant food during gestation) results in offspring with phenotypic traits such as early birth date, large body size, large fat reserves and fast sprint speed which may confer significant fitness benefits in *P. entrecasteauxii* (see also [40] for the importance of the quality of the female diet).

By contrast, when gestation conditions were unfavourable (i.e., restricted basking opportunity and especially limited food), we found that *P. entrecasteauxii* produced premature offspring and offspring with delayed birth, small size, low fat reserves relative to body mass and/or slow sprint speed. Such gestation conditions are described as low net energy gain conditions [24]. Furthermore, these offspring phenotypes have been shown to reduce survival and subsequent reproductive success in other reptiles (e.g., [33,67]), and could be particularly important in our species if accompanied by reduced transport of crucial nutrients and organic substances such as leucine [21] and carotene [40] across the placenta to embryos. Therefore, we suggest that low net energy gain in females during gestation may increase the costs of matrotrophy in part because females may favour allocation of limited resources to their own needs including future reproduction rather than offspring [22].

We thus suggest that the fitness costs and benefits of matrotrophy in *P. entrecasteauxii* may depend on maternal net energy availability during gestation (*sensu* [19]), along with nutrient availability. The advantage of matrotrophy is that it can enable an adjustment of inputs to offspring when conditions allow. Therefore, we suggest that predictably high energy availability during gestation (especially in comparison to energy availability and predictability pre-gestation) may lead to consistent advantages of placental nutrient supply. In time, this may have been an important selective force for the evolution of matrotrophy and its associated mechanisms, in a similar way to which current

Table 3. The results of statistical analyses for offspring growth rate.

Source	F value	P value
Maternal basking	$F_{1, 104} = 0.00$	$P = 0.9804$
Maternal food	$F_{1, 104} = 0.01$	$P = 0.9059$
Offspring basking	$F_{1, 104} = 18.42$	$P < 0.0001$
Offspring sex	$F_{1, 104} = 0.17$	$P = 0.6813$
Maternal basking*maternal food	$F_{1, 104} = 0.44$	$P = 0.5090$
Maternal basking*offspring basking	$F_{1, 104} = 0.00$	$P = 0.9936$
Maternal basking*offspring sex	$F_{1, 104} = 0.01$	$P = 0.9309$
Maternal food*offspring basking	$F_{1, 104} = 0.47$	$P = 0.4945$
Maternal food*offspring sex	$F_{1, 104} = 0.42$	$P = 0.5187$
Offspring basking*offspring sex	$F_{1, 104} = 0.03$	$P = 0.8554$
Maternal basking*maternal food*offspring basking	$F_{1, 104} = 0.00$	$P = 0.9770$
Maternal basking*maternal food*offspring sex	$F_{1, 104} = 1.70$	$P = 0.1951$
Maternal basking*offspring food*offspring sex	$F_{1, 104} = 0.01$	$P = 0.9044$
Maternal food*offspring basking*offspring sex	$F_{1, 104} = 0.89$	$P = 0.3474$
Maternal basking*maternal food*offspring basking*offspring sex	$F_{1, 104} = 0.94$	$P = 0.3347$

Summary of the results of statistical analyses of the effects of maternal basking and food availability (combinations of extended and restricted basking opportunities and food supply) during gestation, offspring basking availability (extended and restricted basking opportunities), offspring sex and their interactions on offspring growth rates in *Pseudemoia entrecasteauxii*. The significant result is indicated in bold.

doi:10.1371/journal.pone.0041835.t003

thermal effects on offspring have been used to argue for viviparity [5,16,17]. In *P. entrecasteauxii*, as in other cold-temperate-zone viviparous reptiles, vitellogenesis occurs in autumn and early spring (e.g., [44,68,69]) when food availability is relatively low and unpredictable compared with the gestational periods of *P. entrecasteauxii* (i.e., late-spring to summer). One way to counter the costs on offspring fitness of matrotrophy when conditions are unfavourable is to alter energy allocation [19,38]. We found that *P. entrecasteauxii* displayed a selective energy allocation among embryos and produced some better offspring within a single clutch at the sacrifice of fitness of other siblings when energy availability was limited during gestation.

So while we have demonstrated the benefits of matrotrophy in some situations (e.g., high food condition during gestation), why it has only evolved in a limited number of viviparous reptiles worldwide (i.e., 4 to 6 origins: [70]) is a key question. A possible explanation for the restricted number of evolutionary transitions to matrotrophy is that the costs of matrotrophy outweigh benefits in many viviparous species. For example, *P. entrecasteauxii* is one of a limited number of species which demonstrate an increase in basking and foraging to enhance maternal energy (and potentially nutrient and organic substance) gain during gestation [23,24,40,46] presumably to support embryonic nutritional requirements. However, these behaviours can increase predation risks, especially for gravid females [46,71]. Therefore, it is possible that the costs of such behaviours can outweigh the benefits in some viviparous reptiles. Furthermore, unpredictable food levels during gestation usually favour lecithotrophic reproduction (i.e., supply of nutrients by the yolk) [72,73]. Another possibility is that placental structure limits nutrient transfer. In reptiles, viviparity has about 100 independent origins [74] with a variety of placental support and complexity [74–76]. Therefore, some evolutionary origins of placental structure may not have the potential for further evolution of matrotrophic reproduction.

Despite the effects of gestation conditions on offspring phenotype and the putative links to fitness in the wild, we found that postnatal conditions significantly affected offspring growth while maternal gestation conditions did not (i.e., we found no evidence for context-dependent maternal effects). In addition, offspring sex did not affect their growth which suggests that context-dependent sex allocation may not occur in this species in response to thermal or food conditions [25,77]. There was no variation in growth rate among offspring in the same thermal regimes, even though their maternal gestation conditions were different. Thus, the maternal effects we demonstrated do not necessarily translate to effects on offspring growth (see also [47,78]). Similarly, incubation temperature in Western fence lizard (*Sceloporus occidentalis*) strongly affected offspring morphology, but did not influence offspring growth rates [27,79]. Mainwaring et al. [80] also found that in a passerine bird (*Cyanistes caeruleus*) postnatal environmental effects were more important than maternal effects in determining offspring growth rate. In addition, context-dependent maternal effects are only predicted under a specific set of circumstances (notably that females or offspring have a priori knowledge during development of the condition they will experience postnatally [6,7]; for our species it is possible that the thermal conditions experienced by females while gravid are not strongly reflected in the conditions their offspring experience).

In conclusion, we have provided the first demonstration that two key environmental factors (temperature and food availability) interact to influence offspring phenotype in a matrotrophic viviparous reptile. Matrotrophy allows maternal effects on offspring phenotype to respond to more than one environmental factor [20,22], in this case basking opportunity and/or food intake

during gestation. Trade-offs between fitness costs and benefits related to maternal net energy availability during gestation may have been an important determinant for the evolution of matrotrophy in *P. entrecasteauxii*. Thus, we suggest that matrotrophy may have evolved as a response to high net energy availability during gestation, which may enhance offspring fitness throughout maternal effects.

Materials and Methods

Maternal conditions during gestation

One hundred and sixty female *P. entrecasteauxii* were collected from one population at the Peter Murrell Reserve in Kingston, southern Tasmania, Australia (41°50'S, 146°36'E) between 15 and 31 October 2007, shortly after ovulation took place. We do not know the history of these females and recognise that their nutritional and thermal history may influence the expression of maternal effects, including for example transgenerational maternal effects. However, given that all females were collected from the same population over a restricted temporal window and because our design split females randomly into their treatments, we do believe this to be an important source of error. The lizards were taken to the Herpetology Laboratory at the School of Zoology at The University of Tasmania where they were measured SVL (± 0.01 mm) and weighed (± 1 mg). The presence of ovulated follicles was confirmed by palpation of the female's abdomen. Females were randomly assigned to one of the following four treatment groups (2 by 2 factorial design):

1. 12 hours basking opportunity per day with high food availability (three mealworms per lizard, three times per week);
2. 12 hours basking opportunity per day with low food availability (two mealworms per lizard, once a week);
3. four hours basking opportunity per day with high food availability; and
4. four hours basking opportunity per day with low food availability.

The thermal conditions reflect the annual variation in hours of sunlight typically encountered by Tasmanian skinks in the natural population and has been used by us for other reptile species to realistically vary thermal opportunity (e.g., [5,25,45,81,82]). The amount of food (high and low) per lizard during gestation was based on our standard husbandry conditions for this and similar sized skinks (i.e., six mealworms per lizard per week). Over long experience (e.g., [5,25,32]), we find that females maintained on such a diet remain healthy, do not lose weight nor gain excessive fat stores. We therefore both increased and decreased this amount of food to provide a comparatively high and low level of food. We recognize that variation in the frequency of feeding may affect female metabolism [83], but given the speed of digestion and that prey are not always consumed immediately, we do not believe that this introduces confounding effects in our analyses. They were maintained in an air-conditioned room under bright fluorescent tube lighting (≈ 20000 lux) and UV lighting (14L: 10D). Each lizard was housed individually in a plastic terrarium (length: 300 mm, width: 200 mm, depth: 100 mm) which contained paper pellets as a substrate, one terracotta saucer and one wooden block as basking sites, and one plastic plate as a shelter; water, supplemented with multi-vitamins and calcium, was available *ad libitum*. Basking heat was supplied by a 25-w spotlight positioned ≈ 80 mm above a basking surface and the thermal gradient in each plastic terrarium was 12–40°C, while the basking light was on allowing free thermoregulation during

this time. All lizards were maintained in these conditions until parturition. Each plastic terrarium was positioned randomly within the experimental group, and was repositioned weekly to minimize position effects.

Maternal and offspring characteristics

For each female, the following data were recorded: date of birth (birthing synchrony), postpartum SVL, maternal growth rate during gestation, postpartum mass, clutch size, RCM and variation in offspring size within-clutch. Gestation length was estimated by assuming that all females ovulated on Nov. 1. Growth rate during gestation was calculated as $(\text{postpartum SVL} - \text{initial SVL}) / (\text{gestation length})^{-1}$. Relative clutch mass was calculated as $(\text{total offspring mass}) / (\text{female postpartum body mass immediately after parturition})^{-1}$ [46], therefore, RCM is unit-less (g per g). Variation in offspring size within-clutch was measured using the coefficient of variation (CV) [84]. The range of clutch sizes was one to six; however, only 10 females produced a single offspring: we removed those offspring from the data set for the CV. The CV expresses variability in within-clutch offspring size relative to the magnitude of the mean within-clutch offspring size. The estimates of within-clutch variation in offspring size are not affected by differences in clutch size among females [85].

For all offspring, we recorded whether they were born alive or dead, SVL, body mass, body pigmentation and sprint speed as a measure of whole body performance [13] on the day of birth. Offspring were not sexed at birth because hemipenes eversion as a predictor of gonadal sex has not been verified in this species and moreover offspring are very small at birth (less than 200 mg) and we did not want to risk tail loss (or stress) prior to growth experiments (see below). Offspring with a weakly pigmented body (transparent body colour and/or red chest colour) were defined as “premature” offspring. Prior to the sprint trials, each lizard was held in a petri dish in a water bath ($28 \pm 1^\circ\text{C}$) for 30 min to reach the optimal temperature for sprinting for this species [86]. Sprint time along the track (120 cm long and 8 cm wide) was recorded by five equally spaced (20 cm) infrared light beams. The fastest sprint speed over a 20 cm distance was taken as the maximum sprint speed (e.g., [63]). Lizards were encouraged to run by occasional gentle taps on the tail using a soft paint brush.

In clutches of two or more offspring, one offspring was selected randomly and killed on the day of birth to determine dry fat reserves. *Pseudemoia entrecasteauxii* stores fat as both abdominal fat bodies and in caudal fat stores. To estimate the mass of abdominal fat bodies for each offspring, the abdominal fat bodies were dissected out and transferred to a pre-weighed (± 0.1 mg) eppendorf tube. The abdominal fat bodies were dried in a 60°C oven at least for two days, and re-weighed (± 0.1 mg). To estimate the mass of caudal fat storage for each sample, we used the method of Chapple and Swain [87], modifying the volume of diethyl ether for the small size of the offspring. The tail was removed at the highest fracture plane, cut into small pieces and transferred to a pre-weighed (± 0.1 mg) eppendorf tube. The tail was then dried in a 60°C oven for at least for two days and the tube was reweighed (± 0.1 mg). The dried tail mass was calculated as $(\text{dried tail plus tube mass} - \text{tube mass})$. The dried tail was then immersed in diethyl ether (1 ml) for at least two days to dissolve the lipid stores, and then transferred to fresh pre-weighed (± 0.1 mg) eppendorf tubes, and placed in a fume cabinet overnight to allow the diethyl ether to evaporate completely. The tube was then re-weighed (± 0.1 mg), to determine the mass of tail minus fat.

Assessing context-dependent maternal effects based on offspring growth rates

We measured offspring growth rate using reciprocal transplant experiments (manipulating thermal condition only). We randomly selected one offspring per litter from each maternal treatment. The rest of the offspring was released in the field. Each offspring was given a unique toe clip for permanent identification. These offspring were allocated into one of two conditions, either 12 hours or four hours basking opportunity per day. This means that the thermal conditions for half of the offspring were identical to those their mothers had experienced during gestation, while the other half of the offspring received a different thermal treatment. Up to five individuals were housed in a plastic terrarium (length: 300 mm, width: 200 mm, depth: 100 mm). Each cage contained paper pellets as a substrate, two terracotta saucers and one wooden block as basking sites, and one plastic plate as a shelter. Water, supplemented with multi-vitamins, was available *ad libitum*. Lizards were fed human baby food (HEINZ® pear flavour), supplemented with multi-vitamins, calcium powder and protein powder three times per week. Other conditions were as described for the females. We measured offspring SVL weekly for five weeks (e.g., [13,22,47]). The growth rate of each individual was calculated using the slope of the least squares regression line of offspring SVL against time (in weeks). In addition, at the end of growth experiment, we determined offspring sex by eversion of hemipenes to investigate if sex had an influence on growth rate and to confirm there were no strong sex biases resulting from maternal treatments.

Statistical analyses

All statistical analyses were performed with SAS® 9.1 for Windows. Differences in maternal initial SVL between maternal treatments were examined using one-way ANOVA. Influences of maternal gestation conditions (i.e., basking and food availability) and their interaction on maternal characteristics, including the date of birth, postpartum SVL, growth rate during gestation, postpartum body mass, clutch size, RCM and within-clutch variation in offspring size (CVs) were examined using full model two-way ANOVA. Basking (12 hours, four hours) and food availability (high, low) were considered as fixed factors. Assumptions of normality were checked by examining plots of standardized residuals against estimated values and the normal probability curve of the residuals; consequently the data for postpartum mass, clutch size and CV were log transformed, and the data for growth rate during gestation and date of birth were square root transformed to meet the assumptions of ANOVA.

Influences of maternal gestation conditions (i.e., basking and food availability) and their interaction on offspring phenotype, including SVL, body mass and sprint speed, were examined using full mixed-model ANOVA. In the mixed-model ANOVA, maternal identity was treated as a random factor to account for litter/clutch effects. Fat reserves were examined using full model two-way ANOVA. To examine differences in offspring fat reserves, we used residuals (i.e., fat reserves relative to body mass), which were generated from a regression analysis of all values for offspring fat reserves (abdominal fat bodies plus caudal fat storage) against offspring body mass. This is because, in general, body mass and fat reserves are positively correlated [88]. In the present study, this assumption was supported overall ($r^2 = 0.4371$, $P < 0.0001$) and also in each maternal treatment. Maternal basking and maternal food availability during gestation were considered as fixed factors. Data for offspring sprint were log transformed to meet the assumptions of ANOVA.

To investigate context-dependent maternal effects, we used a full model four-way ANOVA to analyse the effect of maternal basking and maternal food availability during gestation, offspring basking availability and offspring sex and their interactions on offspring growth rate. Maternal basking and maternal food availability during gestation and offspring basking availability and offspring sex were considered as fixed factors.

All research was carried out with approval of the Animal Ethics Committee (A0009687) of the University of Tasmania, and a permit from the Department of Primary Industries, Water and Environment (FA07177).

References

- Mousseau TA, Fox CW (1998) Maternal Effects as Adaptations. New York: Oxford University Press.
- Dobzhansky T (1935) Maternal effect as a cause of the difference between the reciprocal crosses in *Drosophila pseudoobscura*. P Natl Acad Sci USA 21: 443–446.
- Galloway LF (2005) Maternal effects provide phenotypic adaptation to local environmental conditions. New Phytol 166: 93–99. doi: 10.1111/j.1469-8137.2004.01314.x
- Lindholm AK, Hunt J, Brooks R (2006) Where do all the maternal effects go? Variation in offspring body size through ontogeny in the live-bearing fish *Poecilia parae*. Biol Letters 2: 586–589. doi: 10.1098/rsbl.2006.0546
- Wapstra E, Uller T, While GM, Olsson M, Shine R (2010) Giving offspring a head start in life: field and experimental evidence for selection on maternal basking behaviour in lizards. J Evol Biol 23: 651–657. doi: 10.1111/j.1420-9101.2009.01924.x
- Marshall DJ, Uller T (2007) When is a maternal effect adaptive? Oikos 116: 1957–1963. doi: 10.1111/j.2007.0030-1299.16203.x
- Uller T (2008) Developmental plasticity and the evolution of parental effects. Trends Ecol Evol 23: 432–438. doi: 10.1016/j.tree.2008.04.005
- Mousseau TA, Uller T, Wapstra E, Badyaev AV (2009) Maternal effects as adaptations: past and present. Phil Trans R Soc B 364: 1035–1038. doi: 10.1098/rstb.2008.0303
- Fox CW, Czesak ME (2000) Evolutionary ecology of progeny size in arthropods. Annu Rev Entomol 45: 341–369.
- Einum S, Fleming IA (2004) Environmental unpredictability and offspring size: conservative versus diversified bet-hedging. Evol Ecol Res 6: 443–455.
- Shine R (1995) A new hypothesis for the evolution of viviparity in reptiles. Am Nat 145: 809–823.
- Harlow PS, Taylor JE (2000) Reproductive ecology of the jacky dragon (*Amphibolurus muricatus*): an agamid lizard with temperature-dependent sex determination. Austral Ecol 25: 640–652.
- Wapstra E (2000) Maternal basking opportunity affects juvenile phenotype in a viviparous lizard. Funct Ecol 14: 345–352.
- Warner DA, Shine R (2008) The adaptive significance of temperature-dependent sex determination in a reptile. Nature 451: 566–U5. doi: 10.1038/nature06519
- Robert KA, Thompson MB (2010) Viviparity and temperature-dependent sex determination. Sex Dev 1–2: 119–128. doi: 10.1159/000260373.
- Webb JK, Shine R, Christian KA (2006) The adaptive significance of reptilian viviparity in the tropics: testing the maternal manipulation hypothesis. Evolution 60: 115–122.
- Li H, Qu YF, Hu LJ, Ji X (2009) Evolution of viviparity in cold-climate lizards: testing the maternal manipulation hypothesis. Evol Ecol 23: 777–790. doi: 10.1007/s10682-008-9272-2
- Fowden AL, Giussani DA, Forhead AJ (2006) Intrauterine programming of physiological systems: causes and consequences. Physiology 21: 29–37. doi:10.1152/physiol.00050.2005.
- Trexler JC, DeAngelis DL (2003) Resource allocation in offspring provisioning: an evaluation of the conditions favoring the evolution of matrotrophy. Am Nat 162: 574–585.
- Cadby CD, Jones SM, Wapstra E (2011) Potentially adaptive effects of maternal nutrition during gestation on offspring phenotype of a viviparous reptile. J Exp Biol 214: 4234–4239. doi: 10.1242/jeb.057349
- Itonaga K, Wapstra E, Jones SM (2012) A novel pattern of placental leucine transfer during mid to late gestation in a highly placentotrophic viviparous lizard. J Exp Zool B: In press.
- Itonaga K, Jones SM, Wapstra E (2012) Do gravid females become selfish? Female allocation of energy during gestation. Physiol Biochem Zool 85: 231–242. doi: 10.1086/665367
- Spencer RJ, Thompson MB, Hume ID (1998) The diet and digestive energetics of an Australian short-necked turtle, *Emydura macquarii*. Comp Biochem Phys A 121: 341–349.
- Tsai TS, Lee HJ, Tu MC (2009) Bioenergetic modeling reveals that Chinese green tree vipers select postprandial temperatures in laboratory thermal gradients that maximize net energy intake. Comp Biochem Phys A 154: 394–400. doi: 10.1016/j.cbpa.2009.07.011
- Pen I, Uller T, Feldmeyer B, Harts A, While GM, et al. (2010) Climate driven population divergence in sex determining systems. Nature 468: 436–438. doi: 10.1038/nature09512
- Huey RB, Stevenson RD (1979) Integrating thermal physiology and ecology of ectotherms - discussion of approaches. Am Zool 19: 357–366.
- Sinervo B, Adolph SC (1994) Growth plasticity and thermal opportunity in *Sceloporus* lizards. Ecology 75: 776–790.
- Warner DA, Andrews RM (2002) Laboratory and field experiments identify sources of variation in phenotypes and survival of hatchling lizards. Biol J Linn Soc 76: 105–124.
- Olsson M, Shine R (2002) Growth to death in lizards. Evolution 56: 1867–1870.
- Caley MJ, Schwarzkopf L (2004) Complex growth rate evolution in a latitudinally widespread species. Evolution 58: 862–869.
- Ostrovsky AN, Gordon DP, Lidgard S (2009) Independent evolution of matrotrophy in the major classes of Bryozoa: transitions among reproductive patterns and their ecological background. Mar Ecol Prog Ser 378: 113–124. doi: 10.3354/meps07850
- Swain R, Jones SM (2000) Maternal effects associated with gestation conditions in a viviparous lizard. Herpetol Monogr 14: 432–440.
- Miles DB (2004) The race goes to the swift: fitness consequences of variation in sprint performance in juvenile lizards. Evol Ecol Res 6: 63–75.
- Uller T, Olsson M (2010) Offspring size and timing of hatching determine survival and reproductive output in a lizard. Oecologia 162: 663–671. doi: 10.1007/s00442-009-1503-x
- Jones GP, McCormick MI (2002) Numerical and energetic processes in the ecology of coral reef fishes. In: Sale PF, editor. Coral Reef Fishes - Dynamics and Diversity in a Complex Ecosystem. London: Academic Press. pp. 221–238.
- Liang H, Xiong WH, Zhang ZB (2007) Effect of maternal food restriction during gestation on early development of F1 and F2 offspring in the rat-like hamster (*Cricetus triton*). Zoology 110: 118–126. doi: 10.1016/j.zool.2006.11.004
- Kozlowski J, Stearns SC (1989) Hypotheses for the production of excess zygotes: models of bet-hedging and selective abortion. Evolution 43: 1369–1377.
- Gaillard JM, Yoccoz NG (2003) Temporal variation in survival of mammals: a case of environmental canalization? Ecology 84: 3294–3306.
- Stewart JR, Thompson MB (1993) A novel pattern of embryonic nutrition in a viviparous reptile. J Exp Biol 174: 97–108.
- Itonaga K, Jones SM, Wapstra E (2011) Effects of variation in maternal carotenoid intake during gestation on offspring innate immune response in a matrotrophic viviparous reptile. Funct Ecol 25: 1318–1326. doi: 10.1111/j.1365-2435.2011.01882.x
- Itonaga K, Wapstra E, Jones SM (2011) Evidence for placental transfer of maternal corticosterone in a viviparous lizard. Comp Biochem Phys A. 160: 184–189.
- Brown G (1988) The diet of *Leiophasma entrecasteauxii* (Lacertilia: Scincidae) from south-western Victoria, with notes on its relationship with the reproductive cycle. Aust Wildl Res 15: 605–614
- Wapstra E, Swain R (1996) Feeding ecology of the Tasmanian spotted skink, *Niveoscincus ocellatus* (Squamata: Scincidae). Aust J Zool 44: 205–213.
- Wapstra E, Swain R, Jones SM, O'Reilly J (1999) Geographic and annual variation in the timing of reproductive cycles in a temperate skink, *Niveoscincus ocellatus*. Aust J Zool 47: 539–550.
- Wapstra E, Uller T, Sinn DL, Olsson M, Mazurek K, et al. (2009) Climate effects on offspring sex ratio on a viviparous lizard. J Anim Ecol 78: 84–90. doi: 10.1111/j.1365-2656.2008.01470.x
- Shine R (1980) "Costs" of reproduction in reptiles. Oecologia 46: 92–100.
- Cadby CD, Jones SM, Wapstra E (2010) Are increased concentrations of maternal corticosterone adaptive to offspring? A test using a placentotrophic lizard. Funct Ecol 24: 409–416. doi: 10.1111/j.1365-2435.2009.01637.x
- Burger J (1991) Effects of incubation temperature on behavior of hatchling pine snakes: implications for reptilian distribution. Behav Ecol Sociobiol 28: 297–303.
- VanDamme R, Bauwens D, Braña F, Verheyen RF (1992) Incubation temperature differentially affects hatchling time, egg survival, and hatchling performance in the lizard *Podarcis muralis*. Herpetologica 48: 220–228.

Acknowledgments

We thank Chloé Cadby, Geoffrey While, Tubasa Kato and Jo McEvoy for field and laboratory assistance and thank Louise Oxley and Debbie Ploughman for English assistance. All research was carried out with approval of the Animal Ethics Committee of the University of Tasmania, and a permit under Department of Primary Industries, Water and Environment.

Author Contributions

Conceived and designed the experiments: KI SJ EW. Performed the experiments: KI. Analyzed the data: KI EW. Contributed reagents/materials/analysis tools: KI EW SJ. Wrote the paper: KI EW SJ.

50. Swain R, Jones SM (1997) Maternal transfer of ^3H -labelled leucine in the viviparous lizard *Niveoscincus metallicus* (Scincidae: Lygosominae). *J Exp Zool* 277: 139–145.
51. Jones SM, Swain R (2006) Placental transfer of ^3H -oleic acid in three species of viviparous lizards: a route for supplementation of embryonic fat bodies? *Herpetol Monogr* 20: 186–193.
52. Thompson MB, Stewart JR (1997) Embryonic metabolism and growth in lizards of the genus *Eumeces*. *Comp Biochem Phys A* 118: 647–654.
53. Perrins CM (1970) Timing of birds breeding seasons. *IBIS* 112: 242–255.
54. Watkins TB (1996) Predator-mediated selection on burst swimming performance in tadpoles of the Pacific tree frog, *Pseudacris regilla*. *Physiol Zool* 69: 154–167.
55. Einum S, Fleming IA (2000) Selection against late emergence and small offspring in Atlantic salmon (*Salmo salar*). *Evolution* 54: 628–639.
56. Warner DA, Shine R (2007) Fitness of juvenile lizards depends on seasonal timing of hatching, not offspring body size. *Oecologia* 154: 65–73. doi: 10.1007/s00442-007-0809-9
57. Feder C, Martin JGA, Festa-Bianchet M, Berube C, Jorgenson J (2008) Never too late? Consequences of late birthdate for mass and survival of bighorn lambs. *Oecologia* 156: 773–781. doi: 10.1007/s00442-008-1035-9
58. Atkins N, Swain R, Wapstra E, Jones SM (2007) Late stage deferral of parturition in the viviparous lizard *Niveoscincus ocellatus* (Gray 1845): implications for offspring quality and survival. *Biol J Lin Soc* 60: 735–746.
59. Brown GP, Shine R (2005) Female phenotype, life history, and reproductive success in free-ranging snakes (*Tropidomorphus mairii*). *Ecology* 86: 2763–2770.
60. Kissner KJ, Weatherhead PJ (2005) Phenotypic effects on survival of neonatal northern watersnakes *Nerodia sipedon*. *J Anim Ecol* 74: 259–265. doi: 10.1111/j.1365-2656.2005.00919.x
61. Janzen FJ, Tucker JK, Paukstis GL (2000) Experimental analysis of an early life-history stage: selection on size of hatchling turtles. *Ecology* 81: 2290–2304.
62. Burgess EA, Booth DT, Lanyon JM (2006) Swimming performance of hatchling green turtles is affected by incubation temperature. *Coral Reefs* 25: 341–349. doi: 10.1007/s00338-006-0116-7
63. Melville J, Swain R (2003) Evolutionary correlations between escape behaviour and performance ability in eight species of snow skinks (*Niveoscincus*; Lygosominae) from Tasmania. *J Zool London* 261: 79–89. doi: 10.1017/S0952836903003984
64. Itonaga K, Edwards A, Jones SM, Wapstra E (2012) Interpopulational variation in costs of reproduction related to pregnancy in a viviparous lizard. *Ethol Ecol Evol*: In press.
65. Stapley J (2006) Individual variation in preferred body temperature covaries with social behaviours and colour in male lizards. *J Therm Biol* 31: 362–369. doi: 10.1016/j.jtherbio.2006.01.008
66. While GM, Uller T, McEvoy J, Wapstra E (2009) Long-lasting effects of among-but not within-litter timing of birth in a viviparous lizard. *Evol Ecol Res* 11: 1259–1270.
67. Baron JP, Le Galliard JF, Tully T, Ferrière R (2010) Cohort variation in offspring growth and survival: prenatal and postnatal factors in a late-maturing viviparous snake. *J Anim Ecol* 79: 640–649. doi: 10.1111/j.1365-2656.2010.01661.x
68. Jones SM, Swain R (1996) Annual reproductive cycle and annual cycles of reproductive hormones in plasma of female *Niveoscincus metallicus* from Tasmania. *J Herpetol* 30: 140–146.
69. Murphy K, Hudson S, Shea G (2006) Reproductive seasonality of three cold-temperate viviparous skinks from southeastern Australia. *J Herpetol* 40: 454–464.
70. Blackburn DG, Flemming AF (2010) Reproductive specialization in a viviparous african skink and its implications for evolution and conservation. *Herp Conserv Biol* 5: 263–270.
71. Huey RB, Pianka ER (1981) Ecological consequences of foraging mode. *Ecology* 62: 991–999.
72. Thibault RE, Schultz RJ (1978) Reproductive adaptations among viviparous fishes (*Cyprinodontiformes poeciliidae*). *Evolution* 32: 320–333.
73. Wourms JP, Lombardi J (1992) Reflections on the evolution of piscine viviparity. *Am Zool* 32: 276–293.
74. Blackburn DG (1992) Convergent evolution of viviparity, matrotrophy, and specializations for fetal nutrition in reptiles and other vertebrates. *Am Zool* 32: 313–321.
75. Blackburn DG, Vitt LJ (2002) Specializations of the chorioallantoic placenta in the Brazilian scincid lizard, *Mabuya heathi*: a new placental morphotype for reptiles. *J Morphol* 254: 121–131. doi: 10.1002/jmor.10005
76. Stewart JR, Thompson MB (2003) Evolutionary transformations of the fetal membranes of viviparous reptiles: a case study of two lineages. *J Exp Zool* 299A: 13–32. doi: 10.1002/jez.a.10288
77. Wapstra E, Olsson M, Shine R, Edwards A, Swain R, Joss JMP (2004) Maternal basking behaviour determines offspring sex in a viviparous reptile. *Proc R Soc Lond, Ser B: Biol Sci* 271: S230–S232. doi: 10.1098/rsbl.2003.0152
78. Janczak AM, Heikkilä M, Valros A, Torjesen P, Andersen IL, et al. (2007) Effects of embryonic corticosterone exposure and post-hatch handling on tonic immobility and willingness to compete in chicks. *Appl Anim Behav Sci* 107: 275–286. doi: 10.1016/j.applanim.2006.10.002
79. Buckley CR, Jackson M, Youssef M, Irschick DJ, Adolph SC (2007). Testing the persistence of phenotypic plasticity after incubation in the western fence lizard, *Sceloporus occidentalis*. *Evol Ecol Res* 9: 169–183.
80. Mainwaring MC, Dickens M, Hartley IR (2010) Environmental and not maternal effects determine variation in offspring phenotypes in a passerine bird. *J Evolution Biol* 23: 1302–1311. doi: 10.1111/j.1420-9101.2010.01997.x
81. While GM, Wapstra E (2009) Effects of basking opportunity on birthing asynchrony in a viviparous lizard. *Anim Behav* 77: 1465–1470. doi: 10.1016/j.anbehav.2009.03.001
82. Uller T, While GM, Cadby CD, Harts A, O'Connor K, et al. (2011) Thermal opportunity, maternal effects, and offspring survival at different climatic extremes in a viviparous lizard. *Evolution* 65: 2313–2324.
83. Lourdaís O, Bonnet X, Doughty P (2002) Costs of anorexia during pregnancy in a viviparous snake (*Vipera aspis*). *J Exp Zool* 292: 487–493. doi: 10.1002/jez.10065
84. Crean AJ, Marshall DJ (2009) Coping with environmental uncertainty: dynamic bet hedging as a maternal effect. *Phil Trans R Soc B* 364: 1087–1096. doi: 10.1098/rstb.2008.0237
85. Sokal RR, Rohlf FJ (1995) *Biometry: The Principles and Practice of Statistics in Biological Research*, 3rd edition. New York: W.H. Freeman and Company.
86. Melville J (1998) *The evolution of locomotory mode in the lizard genus Niveoscincus*. PhD dissertation. Hobart: University of Tasmania.
87. Chapple DG, Swain R (2002) Distribution of energy reserves in a viviparous skink: does tail autotomy involve the loss of lipid stores? *Austral Ecol* 27: 565–572.
88. Sinervo B, Hedges R, Adolph SC (1991). Decreased sprint speed as a cost of reproduction in the lizard *Sceloporus occidentalis*: variation among populations. *J Exp Biol* 155: 323–336.