



Vitamin D₃ Supplementation Reduces Subsequent Brain Injury and Inflammation Associated with Ischemic Stroke

Megan A. Evans^{1,2} · Hyun Ah Kim^{1,2} · Yeong Hann Ling^{1,2} · Sandy Uong² · Antony Vinh^{1,2} · T. Michael De Silva^{1,2} · Thiruma V. Arumugam^{3,4} · Andrew N. Clarkson^{5,6} · Graeme R. Zosky⁷ · Grant R. Drummond^{1,2} · Brad R. S. Broughton² · Christopher G. Sobey^{1,2}

Received: 11 January 2018 / Accepted: 16 February 2018 / Published online: 23 February 2018
© The Author(s) 2018. This article is an open access publication

Abstract

Acute inflammation can exacerbate brain injury after ischemic stroke. Beyond its well-characterized role in calcium metabolism, it is becoming increasingly appreciated that the active form of vitamin D, 1,25-dihydroxyvitamin D₃ (1,25-VitD₃), has potent immunomodulatory properties. Here, we aimed to determine whether 1,25-VitD₃ supplementation could reduce subsequent brain injury and associated inflammation after ischemic stroke. Male C57Bl6 mice were randomly assigned to be administered either 1,25-VitD₃ (100 ng/kg/day) or vehicle i.p. for 5 day prior to stroke. Stroke was induced via middle cerebral artery occlusion for 1 h followed by 23 h reperfusion. At 24 h post-stroke, we assessed infarct volume, functional deficit, expression of inflammatory mediators and numbers of infiltrating immune cells. Supplementation with 1,25-VitD₃ reduced infarct volume by 50% compared to vehicle. Expression of pro-inflammatory mediators IL-6, IL-1 β , IL-23a, TGF- β and NADPH oxidase-2 was reduced in brains of mice that received 1,25-VitD₃ versus vehicle. Brain expression of the T regulatory cell marker, Foxp3, was higher in mice supplemented with 1,25-VitD₃ versus vehicle, while expression of the transcription factor, ROR- γ , was decreased, suggestive of a reduced Th17/ $\gamma\delta$ T cell response. Immunohistochemistry indicated that similar numbers of neutrophils and T cells were present in the ischemic hemispheres of 1,25-VitD₃- and vehicle-supplemented mice. At this early time point, there were also no differences in the impairment of motor function. These data indicate that prior administration of exogenous vitamin D, even to vitamin D-replete mice, can attenuate infarct development and exert acute anti-inflammatory actions in the ischemic and reperfused brain.

Keywords Vitamin D · Inflammation · Stroke · Middle cerebral artery occlusion · Mouse

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s12017-018-8484-z>) contains supplementary material, which is available to authorized users.

✉ Christopher G. Sobey
c.sobey@latrobe.edu.au

¹ Vascular Biology Immunopharmacology Group, Department of Physiology, Anatomy and Microbiology, La Trobe University, Bundoora, VIC 3083, Australia

² Cardiovascular Disease Program, Department of Pharmacology, Biomedicine Discovery Institute, Monash University, Clayton, VIC 3800, Australia

³ Department of Physiology, Yong Loo Lin School of Medicine, National University of Singapore, Singapore 119228, Singapore

Introduction

Stroke is the world's second leading cause of death, contributing to 6.7 million deaths annually (Mozaffarian et al. 2016). It is also the most frequent cause of permanent

⁴ School of Pharmacy, Sungkyunkwan University, Suwon, South Korea

⁵ Department of Anatomy, Brain Health Research Centre and Brain Research New Zealand, University of Otago, Dunedin 9054, New Zealand

⁶ School of Life Sciences, La Trobe University, Bundoora, VIC 3083, Australia

⁷ School of Medicine, Faculty of Health Science, University of Tasmania, Hobart, TAS 7005, Australia

disability in adults, with half of all survivors discharged into care (Mozaffarian et al. 2016). Currently, there is only one approved pharmacological agent available to treat stroke, recombinant tissue plasminogen activator (rtPA), which must be administered within a 4.5-h window of stroke onset and only after a CT scan has diagnosed a thrombotic cause (Del Zoppo et al. 2009). Due to these strict limitations, < 10% of stroke patients are eligible to receive rtPA (Reeves et al. 2005; Kleindorfer et al. 2008). Consequently, there is a desperate need to identify modifiable mechanisms capable of limiting the impact of acute stroke.

Secondary brain injury following stroke is driven by local inflammation, production of reactive oxygen species and the infiltration of circulating immune cells (Anrather and Iadecola 2016). Thus, targeting these inflammatory processes has been of intense interest to stroke researchers. However, one immunomodulatory molecule that has received very little attention as a potential stroke therapy is vitamin D, a fat-soluble vitamin that functions as a steroid hormone. Vitamin D is synthesized predominantly from 7-dehydrocholesterol in response to skin exposure to ultraviolet light, but can also be obtained through dietary supplementation (Holick 2007). To become biologically active, vitamin D must first be converted to 1,25-dihydroxyvitamin D₃ (1,25-VitD₃) via two hydroxylation steps. This occurs firstly in the liver by 25-hydroxylase and then typically in the kidney by 1- α -hydroxylase (CYP27B) (Holick 2007). The latter hydroxylation step can also occur in macrophages, T cells and neurons, which also express 1- α -hydroxylase (Lugg et al. 2015). Once in this active form, vitamin D can engage with the vitamin D receptor (VDR) which is located on a number of cell types including leukocytes, endothelial cells, astrocytes and neurons (Provvedini et al. 1983; Merke et al. 1989; Langub et al. 2001; Lee et al. 2008). Vitamin D is best characterized to promote calcium absorption from the small intestine, but recent findings indicate that it may also control expression of a large number of genes, particularly those involved in inflammatory processes (Lugg et al. 2015).

1,25-VitD₃ exerts such immunomodulatory actions through a variety of cellular and molecular mechanisms. Firstly, 1,25-VitD₃ can prevent the development of pathogenic T helper (Th) 1, Th17 and $\gamma\delta$ T cells, and can promote the formation of anti-inflammatory Th2 and T regulatory cells (Zeitelhofer et al. 2017; Chang et al. 2010a; Gregori et al. 2002; Joshi et al. 2011; Nashold et al. 2013; Sloka et al. 2011; Cantorna et al. 2004; Hart et al. 2011; Chen et al. 2005). Studies have also shown that 1,25-VitD₃ promotes the generation of tolerogenic dendritic cells (Takeda et al. 2010; Gorman et al. 2010) and can prevent the release of pro-inflammatory cytokines from monocytes and microglia (Korf et al. 2012; Zhang et al. 2012; Boontanrart et al. 2016; Verma and Kim 2016). Further, 1,25-VitD₃ may inhibit the production of reactive oxygen species by decreasing

expression of NADPH oxidase (NOX) enzymes (Dong et al. 2012) and enhancing expression of antioxidants such as superoxide dismutase and glutathione (Jain and Micinski 2013; Dong et al. 2012).

Observational studies have documented that patients with lower serum levels of vitamin D experience larger infarct volumes and worse functional outcomes following stroke (Tu et al. 2014; Wang et al. 2014; Turetsky et al. 2015; Daubail et al. 2013; Park et al. 2015), suggesting that vitamin D may play a protective role during cerebral ischemia. We recently reported that low baseline levels of vitamin D, resulting from a vitamin D-deficient diet, had no discernible impact on selected outcome measures within 24 h of large vessel occlusion stroke (Evans et al. 2017). Here, we have instead examined the effect of elevated baseline levels of vitamin D achieved by supraphysiological doses of vitamin D given to vitamin D-replete animals during the 5 days prior to stroke, in an analogous manner to high dose supplementation regimes in humans (Wong et al. 2014; Sotirchos et al. 2016). For this, we adopted a similar supplementation regime that was found to reduce vascular injury in mice following hindlimb ischemia (Wong et al. 2014). Indeed, we report that 1,25-VitD₃ supplementation can reduce post-stroke brain injury, reduce expression of pro-inflammatory cytokines, modulate the phenotype of T cells and increase the number of M2-polarized (anti-inflammatory) macrophages/microglia in the brain.

Materials and Methods

Animals

A total of 92 male C57Bl6 mice (7–10 week old; 21–30 g) were used for this study. Mice were housed under a 12-h light/dark cycle and had free access to water and food pellets. Mice were excluded from the study if during the surgical procedure to induce middle cerebral artery occlusion: [1] > 0.2 ml of blood was lost ($n = 1$); [2] subarachnoid hemorrhage occurred ($n = 2$); [3] death occurred during ischemia ($n = 2$); [4] cerebral blood flow failed to reach $\geq 80\%$ pre-ischemic levels upon reperfusion ($n = 1$) and [5] death occurred after reperfusion and prior to the designated time for euthanasia ($n = 7$). All animals were randomly assigned to groups, and the investigator performing the surgical procedure or data analysis was, wherever possible, blinded to the treatment group.

Administration of 1-25,Dihydroxyvitamin D₃

Vitamin D₃ was administered as its active form, 1 α , 25-dihydroxyvitamin D₃ (1,25-VitD₃; Sigma; 100 ng/kg/day) which was dissolved in a solvent mixture of sterile water, propylene

glycol and ethanol in a 5:4:1 ratio. Animals were injected i.p. for 5 consecutive days prior to experimental stroke and again on the day of the procedure, as previously described (Wong et al. 2014).

Middle Cerebral Artery Occlusion

Mice underwent either sham surgery or focal cerebral ischemia as previously described (Evans et al. 2017). Cerebral ischemia was produced in anesthetized mice (ketamine: 80 mg/kg plus xylazine: 10 mg/kg i.p.) by occlusion of the middle cerebral artery (MCA) using a 6.0 silicone-coated monofilament (Doccol Corporation). Rectal temperature was monitored and maintained at 37.0 ± 0.5 °C. MCA occlusion (MCAO) was sustained for 60 min, and the filament then retracted to allow reperfusion. Both successful occlusion (> 70% reduction in cerebral blood flow; CBF) and reperfusion ($\geq 80\%$ return of CBF to the pre-ischemic level) were confirmed by transcranial laser-Doppler flowmetry (PeriMed). Sham-operated mice were anesthetized and the right carotid bifurcation exposed, but no filament was inserted. Neck wounds were then closed with sutures and covered with Betadine® (Sanofi) and spray dressing. Head wounds were closed with superglue, and mice were returned to their cages after regaining consciousness.

Functional Assessment

Mice were assessed for functional deficits at approximately 30 min prior to euthanasia. This comprised a 6-point scoring system for neurological deficits: 0 = normal motor function, 1 = flexion of torso and contralateral forelimb when lifted by the tail, 2 = circling to the contralateral side when held by the tail on a flat surface but normal posture at rest, 3 = leaning to the contralateral side at rest, 4 = no spontaneous motor activity, 5 = death. A hanging grip test was performed as a measure of grasping ability and forelimb strength in which mice were suspended by their forelimbs on a wire between 2 posts 60 cm above a soft pillow for up to 60 s. The time until the animal fell was recorded (a score of 0 s was assigned to animals that fell immediately and a score of 60 s was assigned to animals that did not fall), and the average time of 3 trials with 5 min rests in between was calculated. Spontaneous locomotor activity was assessed using a parallel rod floor apparatus using ANY-maze software coupled to an automated video-tracking system as previously described (Lee et al. 2015).

Assessment of Infarct Volume

Cerebral infarct volumes were determined as previously described (Evans et al. 2017). At 24 h post-stroke, mice were killed by inhalation of isoflurane followed by

decapitation. Brains were immediately removed, snap-frozen in liquid nitrogen and stored at -80 °C. Evenly spread (separated by ~ 420 μm) coronal sections (30 μm) spanning the infarct were cut, thaw-mounted onto poly-L-lysine coated glass slides and stained with 0.1% thionin (Sigma) to delineate the infarct area. Infarct volume was quantified using image analysis software (ImageJ, NIH), correcting for brain edema, according to the following formula: $CIV = (LHA - (RHA - RIA)) \times (\text{thickness of section} + \text{distance between sections})$; where *CIV* is corrected infarct volume, *LHA* is left hemisphere area, *RHA* is right hemisphere area and *RIA* is right hemisphere infarct area. Edema-corrected infarct volumes of individual brain sections were then added, giving an approximation of the total infarct volume.

Real-Time Polymerase Chain Reaction (rt-PCR)

At 24 h following stroke or sham surgery, mice were euthanized by isoflurane overdose and perfused with RNase-free phosphate-buffered saline (PBS). After removing the cerebellum and olfactory bulbs, the brain was separated into left and right hemispheres and snap-frozen in liquid nitrogen for RNA extraction. Spleens were also removed, cut in half and snap-frozen in liquid nitrogen. Tissues were stored at -80 °C until required. Total RNA was extracted using Qiazol® reagent (Qiagen) and the RNeasy Mini Kit with on-column DNase step (Qiagen) followed by cDNA conversion using the Quantitect Reverse Transcription kit (for Taqman® gene expression assays; Qiagen). The cDNA was then used as a template in real-time PCR to measure mRNA expression of *Vdr*, *Cyp27b*, *Cyp24a*, *Cxcl12*, *Tbx21*, *Stat4*, *Rorc*, *Gata3*, *Stat6*, *Foxp3*, *Tnfa*, *Il1 β* , *Il6*, *Il21*, *Il23a*, *Tgfb1*, *Ccl2*, *Ccl5*, *Gp91phox*, *Mrc1*, *Il10* and *Icam1*. *Gapdh* and *β -actin* were assessed as housekeeping genes for brain and spleen tissue, respectively. Assays were performed according to the manufacturer's instructions using the Bio-Rad CFX96™ real-time PCR machine (Bio-Rad). Data were normalized to the housekeeping gene and calculated as change in fold expression relative to sham using the formula: $\text{fold-change} = 2^{-\Delta\Delta C_t}$.

Immunofluorescence

Six serial coronal sections (10 μm thick) per animal were collected at six regions: bregma + 0.06, -0.78 , -1.2 , -1.62 , -2.04 , -2.46 mm. Frozen brain sections (10 μm) were fixed in 4% paraformaldehyde for 15 min and washed in 0.01 M PBS (3×10 min). Sections were then blocked with 10% goat serum (Sigma) for 60 min to block non-specific binding of the secondary antibody. Sections were then incubated overnight at 4 °C with either rabbit anti-CD3 (1:200; Abcam) or rabbit anti-CD206 (1:500; Abcam). On

the following day, they were washed (PBS; 3×10 min) and incubated for a maximum of 2 h with either goat anti-rabbit Alexa Fluor 594 (1:500; Thermofisher Scientific) or goat anti-rabbit Alexa Fluor 488 (1:500; Thermofisher Scientific). Finally, sections were again washed and then mounted with Vectashield medium containing 4,6-diamidino-2-phenylindole (DAPI) (Vector Laboratories), and a coverslip was applied. All tissue-mounted slides were viewed, analyzed and photographed with an Olympus fluorescence microscope. Numbers of immunoreactive cells were counted manually per whole ischemic hemisphere and then averaged across the six regions, as indicated above.

3,3'-Diaminobenzidine (DAB) Immunohistochemistry

Frozen brain sections (at the regions indicated for immunofluorescence) were fixed in 4% paraformaldehyde for 15 min, washed in PBS and then incubated in peroxidase blocking solution (Dako) for 10 min to block endogenous peroxidases followed by 10% goat serum for 60 min. They were then incubated overnight at room temperature in rabbit anti-myeloperoxidase (1:100; Abcam). The following day, sections were washed and incubated for 2 h in anti-rabbit IgG horse-radish peroxidase conjugate (1:200; Dako), washed again, and DAB (Dako) was then applied for 5–10 min. Sections were then rinsed in dH₂O, dehydrated in increasing concentrations of ethanol (70 and 100% vol/vol), cleared in xylene and mounted in DPX. Tissue-mounted slides were viewed, analyzed and photographed using an Olympus light microscope. Numbers of immunoreactive cells were counted manually per whole ischemic hemisphere and then averaged across the six regions, as indicated above.

Statistical Analysis

Data are presented as mean \pm standard error of the mean (SEM), with the exception of neurological deficit scores, which are presented as median. Statistical analyses were performed using GraphPad Prism version 6.0 (GraphPad Software Inc. San Diego, CA, USA). Between-group comparisons were compared using one-way ANOVA, or Student's unpaired *t* test, as appropriate. If differences were detected by ANOVA, individual groups were compared with Tukey's multiple comparisons test, where indicated. Neurological deficit scores were compared using a Kruskal–Wallis test followed by Dunn's multiple comparisons test. If there were two independent variables, data were compared using a two-way ANOVA. Statistical significance was accepted if $P < 0.05$.

Results

Effect of Cerebral Ischemia on Vitamin D-Associated Genes in Brain and Spleen

The effect of stroke was first assessed on the expression of VDR and metabolizing enzymes in the brain and spleen at 24 h in otherwise untreated animals. Stroke increased expression of the VDR by \sim twofold in both organs (Fig. 1a, b). The vitamin D-activating enzyme, *Cyp27b*, was reduced by \sim 30% in the brain, but was unchanged in the spleen following stroke (Fig. 1a, b). Expression of the inactivating enzyme, *Cyp24a*, was increased by \sim 2.5 fold in the brain, but was undetectable in spleen (Fig. 1a, b).

Effects of Vitamin D₃ Supplementation on Infarct Volume and Functional Deficits Following Stroke

To determine the effect of elevated baseline vitamin D prior to stroke on the extent of subsequent infarct development, mice were treated with 1,25-VitD₃ (100 ng/kg/day)




Fig. 1 Post-stroke expression of vitamin D-associated genes. mRNA expression of the vitamin D receptor (*Vdr*), 1- α -hydroxylase (*Cyp27b*) and 24-hydroxylase (*Cyp24a*) was examined in **a** the brain and **b** the spleen at 24 h after either stroke or sham surgery. Sham: $n = 6$ –9 per group and stroke: $n = 8$ –12 per group. N.D. denotes not detected. * $P < 0.05$, Student's unpaired *t* test. Data are presented as mean \pm SEM

for 5 days and then subjected to focal cerebral ischemia. At 24 h post-stroke, we found that mice which received 1,25-VitD₃ supplementation had ~ 50% smaller infarct volume than those which received vehicle (Fig. 2a, b). This finding was not associated with any differences in the level of cerebral blood flow during, or immediately after, cerebral ischemia (Fig. S1). Examining the distribution of the infarcts, 1,25-VitD₃-supplemented animals tended to have a reduced infarct area in most coronal sections (Fig. 2c).

However, mice in both groups displayed similar functional deficits at this early time point (Fig. 3a–e).

Effect of Vitamin D₃ Supplementation on T Cell Phenotype in the Brain and Spleen After Stroke

Previous reports suggest that Th1 and $\gamma\delta$ T cells exacerbate brain injury following stroke while Th2 and T regulatory cells play a protective role by dampening excessive inflammation (Gu et al. 2012; Gelderblom et al. 2012; Benakis




Fig. 2 1,25-Dihydroxyvitamin D₃ (VitD₃) treatment reduces infarct development. **a** Representative coronal brain sections from vehicle- and VitD₃-treated animals (infarct area is outline in white). **b** Infarct volume and **c** distribution of infarct area from vehicle- and

VitD₃-treated animals. Vehicle: *n* = 14 per group and VitD₃: *n* = 12 per group. **P* < 0.05, Student’s unpaired *t* test, where appropriate. Data are presented as mean ± SEM

Fig. 3 1,25-Dihydroxyvitamin D₃ (VitD₃) treatment does not influence functional outcome at 24 h following stroke. **a** Neurological deficit scores and **b** latency to fall on hanging grip test of sham, vehicle- and VitD₃-treated animals. Parallel rod floor test examining **c** total distance travelled, **d** percentage time mobile and **e** average speed of sham, vehicle- and VitD₃-treated animals. Sham: *n* = 6–8 per group, vehicle: *n* = 14 per group and VitD₃: *n* = 11–14 per group. ***P* < 0.01, ****P* < 0.001, Kruskal–Wallis test followed by Dunn’s multiple comparisons test (**a**) or one-way ANOVA followed by Tukey’s multiple comparison test (**b–e**). Data are presented as mean ± SEM, with the exception of neurological deficit scores which are presented as median





Fig. 4 1,25-Dihydroxyvitamin D₃ (VitD₃) supplementation alters mRNA expression of T cell transcription factors in the brain and spleen following stroke. **a** Expression of Th1 transcription factors, *Tbx21* and *Stat4*, and Th17 transcription factor, *Rorc*, within the brain of sham, vehicle- (Veh) and VitD₃-treated mice at 24 h post-surgery. **b** Expression of Th2 transcription factors, *Gata3* and *Stat6*, and T regulatory cell transcription factor, *Foxp3*, within the brain of sham, vehicle- and VitD₃-treated mice at 24 h post-surgery. **c** Expression of *Tbx21*, *Stat4*, and *Rorc* within the spleen of sham, vehicle- and VitD₃-treated mice at 24 h post-surgery. **d** Expression of *Gata3*, *Stat6* and *Foxp3* within the spleen of sham, vehicle- and VitD₃-treated mice 24 h post-surgery. Sham: *n* = 6–9 per group, vehicle: *n* = 11–13 per group, VitD₃: *n* = 10–12 per group. **P* < 0.05, ***P* < 0.01, ****P* < 0.001, one-way ANOVA followed by Tukey's multiple comparisons test. Data are presented as mean ± SEM



et al. 2016; Liesz et al. 2009). It has been shown that vitamin D can modulate the immune response to injury by polarizing T cells toward an anti-inflammatory phenotype (Hart et al.


2011). We found no effect of stroke or 1,25-VitD₃ on mRNA expression of Th1 transcription factors, *Tbx21* or *Stat4*, or Th2 transcription factors, *Gata3* or *Stat6* in the brain

(Fig. 4a, b). However, stroke resulted in an elevation of the Th17/γδ T cell transcription factor, *Rorc*, and this effect was mitigated by 1,25-VitD₃ treatment (Fig. 4a). Moreover, we noted an increase in the T regulatory cell transcription factor, *Foxp3*, after stroke, and this was augmented in 1,25-VitD₃-supplemented animals (Fig. 4b). In the spleen, 1,25-VitD₃ had no effect on expression of *Tbx21*, *Stat4*, *Gata3*, *Stat6* or *Rorc* (Fig. 4c, d). However, *Foxp3* expression was slightly higher after stroke in 1,25-VitD₃-treated mice than in sham mice or in those treated with vehicle (Fig. 4d).

Effect of Vitamin D₃ Supplementation on Expression of Pro-inflammatory Mediators in the Brain Following Stroke

As vitamin D₃ has immunomodulatory actions, we also examined mRNA expression of various inflammatory mediators known to be involved in ischemic brain injury. Indeed, 1,25-VitD₃-treated animals had lower expression of *Il1β*, *Il6*, *Il23a*, *Tgfb1* and *Gp91phox* (NOX-2) than vehicle-treated animals (Fig. 5b, c, e, f, i). However, there was no effect of

Fig. 5 1,25-Dihydroxyvitamin D₃ (VitD₃) supplementation alters mRNA expression of inflammatory mediators in the brain following stroke. mRNA expression of **a** *Tnfa*, **b** *Il1β*, **c** *Il6*, **d** *Il21*, **e** *Il23a*, **f** *Tgfb1*, **g** *Ccl2*, **h** *Ccl5*, **i** *Gp91phox*, **j** *Mrc1*, **k** *Icam1* and **l** *Il10* within the brains of sham, vehicle (Veh)- and VitD₃-treated mice at 24 h post-surgery. Sham: *n* = 6–7 per group, vehicle: *n* = 12 per group and VitD₃: *n* = 9–11 per group. **P* < 0.05, ***P* < 0.01, ****P* < 0.001, one-way ANOVA followed by Tukey’s multiple comparisons test. Data are presented as mean ± SEM



1,25-VitD₃ on *Tnfa*, *Il21*, *Ccl2*, *Ccl5*, *Mrc1*, *Icam1* or *Il10* (Fig. 5a, d, g, h, j, k, l).


Effect of Vitamin D₃ Supplementation on Numbers of Infiltrating Leukocytes and M2-Polarized Macrophages/Microglia in the Brain Following Stroke

We tested for any effect of vitamin D on migration of immune cells toward the site of injury, by quantifying leukocyte infiltration into the ischemic hemisphere using immunohistochemistry. We noted a tendency for 1,25-VitD₃-treated animals to have fewer MPO⁺ neutrophils in the brain at 24 h post-stroke, whereas there was no effect on CD3⁺ T cells (Fig. 6a, b). Additionally, there was a trend for greater numbers of “M2” polarized microglia/macrophages (defined as CD206⁺) after stroke in 1,25-VitD₃-treated animals (Fig. 6c).

Discussion

Inflammation is a major contributor to secondary brain injury after ischemic stroke and thus represents a potential target for therapy (Anrather and Iadecola 2016). Beyond its well-characterized role in calcium metabolism, vitamin D has potent immunomodulatory properties and can alter the immune response to injury in various disease settings (Nashold et al. 2013; Takeda et al. 2010; Martorell et al. 2016; Schedel et al. 2016). If vitamin D was found to exert such effects in post-stroke brain injury, it could represent a novel direction for acute therapy. Indeed, here we report data supporting this concept. This neuroprotective effect appears to occur in association with reduced expression of pro-inflammatory mediators in the brain. Moreover, our data suggest that 1,25-VitD₃ supplementation alters the phenotype of T cells and increases numbers

Fig. 6 Quantification of leukocytes in the brain post-stroke. Immunohistochemistry was used to determine the numbers of **a** myeloperoxidase (MPO)⁺ cells, **b** CD3⁺ cells and **c** CD206⁺ cells per right (ischemic) hemisphere in vehicle- and 1,25-dihydroxyvitamin D₃ (VitD₃)-treated animals at 24-h post-stroke. Vehicle: *n* = 9–11 per group and VitD₃: *n* = 9–10 per group. Student's unpaired *t* test. Data are presented as mean ± SEM. Arrows on representative images indicate examples of positive cells, and scale bars represent either 20 μm (**b**) or 50 μm (**a**, **c**)



of M2 macrophages/microglia in the ischemic brain, both of which may contribute to the neuroprotection by 1,25-VitD₃ treatment.

Previous studies have demonstrated that the VDR and the vitamin D regulatory enzymes, 1- α -hydroxylase and 24-hydroxylase, to be expressed in non-classical tissues such as the brain and activated immune cells, suggesting that vitamin D may exert paracrine functions (Penna et al. 2007; Overbergh et al. 2000; Provvedini et al. 1983; Eyles et al. 2005). Additionally, studies have documented that the expression of the VDR and these enzymes can be altered during inflammation and disease (Luo et al. 2013; Yao et al. 2015; von Essen et al. 2010; Yang et al. 2011; Liu et al. 2006; Spanier et al. 2012). In the current study, we examined expression of the VDR (*Vdr*), 1- α -hydroxylase (*Cyp27b*) and 24-hydroxylase (*Cyp24a*), in both the brain and spleen at 24 h after stroke or sham surgery. We found that expression of the VDR was elevated in both organs after stroke. Interestingly, we observed that expression of the vitamin D-activating enzyme, 1- α -hydroxylase, was reduced in the brain after stroke, while expression of the vitamin D inactivating enzyme, 24-hydroxylase, was increased. However, in the spleen, we observed that expression of 1- α -hydroxylase and 24-hydroxylase was unchanged and undetected, respectively. These findings may imply that local levels of the active form of endogenous vitamin D may be reduced in the brain after stroke, raising the possibility that supplementation with exogenous 1,25-VitD₃ may be of benefit.

Indeed, we found that 1,25-VitD₃-supplemented animals developed a smaller infarct volume than vehicle-treated controls by 24 h. However, at this time point, there were no apparent differences in functional outcome. While 24 h is a relatively early time point for examining outcomes after stroke, we know from our previous work that infarct size is fully developed within 24 h in this model of stroke (Evans et al. 2018). Therefore, in seeking to test whether vitamin D might exert a neuroprotective effect to limit infarct development potentially by inhibiting inflammation, we chose to examine outcomes at 24 h. However, we do acknowledge the importance of evaluating the effect of 1,25-VitD₃ at later time points after stroke, particularly on functional recovery. Our findings are analogous to those reported by two previous studies using rat models of stroke, whereby 1,25-VitD₃ pretreatment reduced infarct volume (Fu et al. 2013; Oermann et al. 2004). However, neither of these studies examined functional outcome. Moreover, the precise mechanisms by which 1,25-VitD₃ reduced brain injury in those studies were unclear.

To this end, we tested for evidence that 1,25-VitD₃ may modulate the immune response to ischemic stroke. Vitamin D can modulate the phenotype of T cells (Hart et al. 2011; Cantorna et al. 1996). For instance, in mouse models of multiple sclerosis vitamin D can down-regulate signaling

pathways essential for development of Th1 and Th17 cells (Zeitelhofer et al. 2017; Mattner et al. 2000; Muthian et al. 2006; Chang et al. 2010b; Joshi et al. 2011). Moreover, vitamin D can promote the formation of Th2 and T regulatory cells (Hart et al. 2011) and limit the development of $\gamma\delta$ T cells (Chen et al. 2005). Several studies have revealed that Th1 and $\gamma\delta$ T cells can aggravate brain injury after stroke, and that blocking their invasion may be neuroprotective (Gu et al. 2012; Yilmaz et al. 2006; Gelderblom et al. 2012; Shichita et al. 2009). Th2 and T regulatory cells are thought to be injury-limiting in the setting of stroke (Gu et al. 2012; Liesz et al. 2009). We thus examined whether the neuroprotection by 1,25-VitD₃ may be associated with modulation of T cell phenotypes. In the brain, we found that neither stroke nor 1,25-VitD₃ had any effect on expression of Th1 or Th2 transcription factors. However, 1,25-VitD₃ blunted expression of the Th17/ $\gamma\delta$ T cell transcription factor, *Rorc* (ROR- γ t), and enhanced expression of the T regulatory cell transcription factor, *Foxp3*. In the spleen 1,25-VitD₃ increased expression of *Foxp3*, but had no effect on Th1, Th2 or Th17/ $\gamma\delta$ transcription factors. Collectively, these data may indicate that 1,25-VitD₃ promotes the formation of T regulatory cells while inhibiting development of Th17/ $\gamma\delta$ T cells, consistent with a neuroprotective profile.

1,25-VitD₃ supplementation reduced mRNA expression of pro-inflammatory cytokines, *Il1 β* (IL-1 β), *Il6* (IL-6), *Tgfb1* (TGF- β) and *Il23a* (IL-23a). Interestingly, these cytokines are thought to play key roles in the function of both Th17 and $\gamma\delta$ T cells (Vantourout and Hayday 2013). Treatment with 1,25-VitD₃ had no effect on expression of *Il10* (IL-10), an immunosuppressive cytokine often involved in T regulatory cell function (Taylor et al. 2006); however, T regulatory cells may limit injury and excessive inflammation via other mechanisms (Sakaguchi et al. 2009). We also observed a reduction in *gp91phox* (NOX2) expression, a key producer of superoxide and mediator cellular damage following ischemic stroke (De Silva et al. 2011).

As 1,25-VitD₃ can reduce leukocyte recruitment to injured tissues (Pedersen et al. 2007; Korf et al. 2012; Grishkan et al. 2013), we examined its effect on leukocyte infiltration into the brain following stroke. We observed a trend for 1,25-VitD₃-treated animals to have fewer infiltrating neutrophils but no apparent effect on T cells. It is also possible that 1,25-VitD₃ reduces recruitment of other types of immune cell subsets or that it modulates their functional phenotype rather than migration to the site of post-stroke injury. It is important to note that 24 h represents a relatively early pathological time point after stroke with significant immune cell infiltration continuing after this time point (Gelderblom et al. 2009; Benakis et al. 2016). We also examined the effect of 1,25-VitD₃ on the numbers of M2-polarized macrophages/microglia in the brain after stroke. Studies have reported that M2 macrophages/microglia are likely to be protective in the

setting of stroke by reducing inflammation and coordinating repair processes (Benakis et al. 2014; Chu et al. 2015; Hu et al. 2012). We observed a strong trend for 1,25-VitD₃ to augment numbers of CD206⁺ M2 macrophages/microglia in the brain after stroke.

As mentioned above, there is a strong rationale to gain a deeper understanding of how altered levels of vitamin D—prior and/or subsequent to stroke—might impact on the degree of ensuing brain injury. Following on from our previous finding that low baseline vitamin D levels did not impact on outcome measures at 24 h (Evans et al. 2017), here we have instead assessed the effect of increasing baseline levels achieved by five daily supraphysiological doses of vitamin D prior to stroke. Indeed, the present data suggest that supplementing mice with the active form of vitamin D prior to stroke can reduce the extent of brain injury. With regard to therapeutic relevance for acute clinical stroke, our data are important in terms of proof-of-concept. However, a limitation is that 1,25-VitD₃ was administered only prior to stroke induction, and clearly, studies are now required in which post-stroke treatment of 1,25-VitD₃ is evaluated. While our data suggest that 1,25-VitD₃ can also modulate the immune response to brain injury following stroke, at least part of this protection may occur via non-immune mechanisms, such as inhibiting excitotoxicity (Taniura et al. 2006; Brewer et al. 2001), stimulating production of neurotrophic factors (Neveu et al. 1994; Naveilhan et al. 1996; Landel et al. 2016) or improving blood brain barrier integrity (Won et al. 2015). It is also noteworthy that we administered 1,25-VitD₃ to mice that were vitamin D replete. Whether a similar or greater level of neuroprotection might be achieved in vitamin D-deficient animals by 1,25-VitD₃ therapy will be important to clarify.

In conclusion, these findings indicate that administration of vitamin D can attenuate infarct development following stroke possibly by modulating the inflammatory response to cerebral ischemia. Therefore, vitamin D supplementation may represent a novel direction for limiting the impact of acute stroke.

Funding We wish to acknowledge support from Monash Faculty Postgraduate Scholarship (MAE), Australian Postgraduate Award (YHL) and NHMRC Senior Research Fellowships (GRD and CGS).

Compliance with Ethical Standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical Approval All procedures performed in these studies involving animals were in accordance with the ethical standards of the institution or practice at which the studies were conducted. Specifically, the studies were approved by Monash University Animal Ethics Committees and performed in accordance with the National Health and Medical Research Council of Australia guidelines for the care and use of ani-

mals in research. This article does not contain any studies with human participants performed by any of the authors.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

- Anrather, J., & Iadecola, C. (2016). Inflammation and stroke: An overview. *Neurotherapeutics*, 13(4), 661–670. <https://doi.org/10.1007/s13311-016-0483-x>.
- Benakis, C., Brea, D., Caballero, S., Faraco, G., Moore, J., Murphy, M., et al. (2016). Commensal microbiota affects ischemic stroke outcome by regulating intestinal $\gamma\delta$ T cells. *Nature Medicine*, 22(5), 516–523. <https://doi.org/10.1038/nm.4068>.
- Benakis, C., Garcia-Bonilla, L., Iadecola, C., & Anrather, J. (2014). The role of microglia and myeloid immune cells in acute cerebral ischemia. *Frontiers in Cellular Neuroscience*, 8, 461. <https://doi.org/10.3389/fncel.2014.00461>.
- Boontanart, M., Hall, S. D., Spanier, J. A., Hayes, C. E., & Olson, J. K. (2016). Vitamin D3 alters microglia immune activation by an IL-10 dependent SOCS3 mechanism. *Journal of Neuroimmunology*, 292, 126–136. <https://doi.org/10.1016/j.jneuroim.2016.01.015>.
- Brewer, L. D., Thibault, V., Chen, K.-C., Langub, M. C., Landfield, P. W., & Porter, N. M. (2001). Vitamin D hormone confers neuroprotection in parallel with downregulation of L-Type calcium channel expression in hippocampal neurons. *Journal of Neuroscience*, 21(1), 98–108.
- Cantorna, M. T., Hayes, C. E., & DeLuca, H. F. (1996). 1,25-Dihydroxyvitamin D3 reversibly blocks the progression of relapsing encephalomyelitis, a model of multiple sclerosis. *Proceedings of the National Academy of Sciences, USA*. <https://doi.org/10.1073/pnas.93.15.7861>.
- Cantorna, M. T., Zhu, Y., Froicu, M., & Wittke, A. (2004). Vitamin D status, 1,25-dihydroxyvitamin D3, and the immune system. *American Journal of Clinical Nutrition*, 80(6 Suppl), 1717s–1720s.
- Chang, J. H., Cha, H. R., Lee, D. S., Seo, K. Y., & Kweon, M. N. (2010a). 1,25-Dihydroxyvitamin D3 inhibits the differentiation and migration of T(H)17 cells to protect against experimental autoimmune encephalomyelitis. *PLoS ONE*, 5, e12925. <https://doi.org/10.1371/journal.pone.0012925>.
- Chang, S. H., Chung, Y., & Dong, C. (2010b). Vitamin D suppresses Th17 cytokine production by inducing C/EBP homologous protein (CHOP) expression. *Journal of Biological Chemistry*, 285(50), 38751–38755. <https://doi.org/10.1074/jbc.C110.185777>.
- Chen, L., Cencioni, M. T., Angelini, D. F., Borsellino, G., Battistini, L., & Brosnan, C. F. (2005). Transcriptional profiling of $\gamma\delta$ T cells identifies a role for vitamin D in the immunoregulation of the V γ 9 V δ 2 response to phosphate-containing ligands. *The Journal of Immunology*, 174(10), 6144–6152. <https://doi.org/10.4049/jimmunol.174.10.6144>.
- Chu, H. X., Broughton, B. R., Kim, H. A., Lee, S., Drummond, G. R., & Sobey, C. G. (2015). Evidence that Ly6C(hi) monocytes are protective in acute ischemic stroke by promoting M2 macrophage polarization. [Research Support, Non-U.S. Gov't]. *Stroke*, 46(7), 1929–1937. <https://doi.org/10.1161/strokeaha.115.009426>.

- Daubail, B., Jacquin, A., Guillard, J. C., Hervieu, M., Ossey, G. V., Rouaud, O., et al. (2013). Serum 25-hydroxyvitamin D predicts severity and prognosis in stroke patients. *European Journal of Neurology*, *20*(1), 57–61. <https://doi.org/10.1111/j.1468-1331.2012.03758.x>.
- De Silva, T. M., Brait, V. H., Drummond, G. R., Sobey, C. G., & Miller, A. A. (2011). Nox₂ oxidase activity accounts for the oxidative stress and vasomotor dysfunction in mouse cerebral arteries following ischemic stroke. *PLoS ONE*, *6*(12), e28393. <https://doi.org/10.1371/journal.pone.0028393>.
- Del Zoppo, G. J., Saver, J. L., Jauch, E. C., & Adams, H. P., Jr. (2009). Expansion of the time window for treatment of acute ischemic stroke with intravenous tissue plasminogen activator: a science advisory from the American Heart Association/American Stroke Association. [Practice guideline]. *Stroke*, *40*(8), 2945–2948. <https://doi.org/10.1161/strokeaha.109.192535>.
- Dong, J., Wong, S. L., Lau, C. W., Lee, H. K., Ng, C. F., Zhang, L., et al. (2012). Calcitriol protects renovascular function in hypertension by down-regulating angiotensin II type I receptors and reducing oxidative stress. *European Heart Journal*, *33*(23), 2980–2990. <https://doi.org/10.1093/eurheartj/ehr459>.
- Evans, M. A., Kim, H. A., De Silva, T. M., Arumugam, T. V., Clarkson, A. N., Drummond, G. R., et al. (2017). Diet-induced vitamin D deficiency has no effect on acute post-stroke outcomes in young male mice. *Journal of Cerebral Blood Flow and Metabolism*. <https://doi.org/10.1177/0271678x17719208>.
- Evans, M. A., Lim, R., Kim, H. A., Chu, H. X., Gardiner-Mann, C. V., Taylor, K. W. E., et al. (2018). Acute or delayed systemic administration of human amnion epithelial cells improves outcomes in experimental stroke. *Stroke*. <https://doi.org/10.1161/strokeaha.117.019136>.
- Eyles, D. W., Smith, S., Kinobe, R., Hewison, M., & McGrath, J. J. (2005). Distribution of the vitamin D receptor and 1 alpha-hydroxylase in human brain. *Journal of Chemical Neuroanatomy*, *29*(1), 21–30. <https://doi.org/10.1016/j.jchemneu.2004.08.006>.
- Fu, J., Xue, R., Gu, J., Xiao, Y., Zhong, H., Pan, X., et al. (2013). Neuroprotective effect of calcitriol on ischemic/reperfusion injury through the NR3A/CREB pathways in the rat hippocampus. *Mol Med Rep*, *8*(6), 1708–1714. <https://doi.org/10.3892/mmr.2013.1734>.
- Gelderblom, M., Leyboldt, F., Steinbach, K., Behrens, D., Choe, C. U., Siler, D. A., et al. (2009). Temporal and spatial dynamics of cerebral immune cell accumulation in stroke. *Stroke*, *40*(5), 1849–1857. <https://doi.org/10.1161/strokeaha.108.534503>.
- Gelderblom, M., Weymar, A., Bernreuther, C., Velden, J., Arunachalam, P., Steinbach, K., et al. (2012). Neutralization of the IL-17 axis diminishes neutrophil invasion and protects from ischemic stroke. *Blood*, *120*(18), 3793–3802.
- Gorman, S., Judge, M. A., & Hart, P. H. (2010). Topical 1,25-dihydroxyvitamin D3 subverts the priming ability of draining lymph node dendritic cells. *Immunology*, *131*(3), 415–425. <https://doi.org/10.1111/j.1365-2567.2010.03315.x>.
- Gregori, S., Giarratana, N., Smiroldo, S., Uskokovic, M., & Adorini, L. (2002). A 1 α ,25-dihydroxyvitamin D analog enhances regulatory T-cells and arrests autoimmune diabetes in NOD mice. *Diabetes*, *51*(5), 1367–1374. <https://doi.org/10.2337/diabetes.51.5.1367>.
- Grishkan, I. V., Fairchild, A. N., Calabresi, P. A., & Gocke, A. R. (2013). 1,25-dihydroxyvitamin D3 selectively and reversibly impairs T helper-cell CNS localization. *Proceedings of the National Academy of Sciences of the United States of America*, *110*(52), 21101–21106. <https://doi.org/10.1073/pnas.1306072110>.
- Gu, L., Xiong, X., Zhang, H., Xu, B., Steinberg, G. K., & Zhao, H. (2012). Distinctive effects of T cell subsets in neuronal injury induced by cocultured splenocytes in vitro and by in vivo stroke in mice. *Stroke*, *43*(7), 1941–1946.
- Hart, P. H., Gorman, S., & Finlay-Jones, J. J. (2011). Modulation of the immune system by UV radiation: more than just the effects of vitamin D? *Nature Reviews Immunology*, *11*(9), 584–596. <https://doi.org/10.1038/nri3045>.
- Holick, M. F. (2007). Vitamin D deficiency. *New England Journal of Medicine*, *357*(3), 266–281. <https://doi.org/10.1056/NEJMr070553>.
- Hu, X., Li, P., Guo, Y., Wang, H., Leak, R. K., Chen, S., et al. (2012). Microglia/macrophage polarization dynamics reveal novel mechanism of injury expansion after focal cerebral ischemia. *Stroke*, *43*(11), 3063–3070. <https://doi.org/10.1161/STROKEAHA.112.659656>.
- Jain, S. K., & Micinski, D. (2013). Vitamin D upregulates glutamate cysteine ligase and glutathione reductase, and GSH formation, and decreases ROS and MCP-1 and IL-8 secretion in high-glucose exposed U937 monocytes. *Biochemical and Biophysical Research Communications*, *437*(1), 7–11. <https://doi.org/10.1016/j.bbrc.2013.06.004>.
- Joshi, S., Pantalena, L. C., Liu, X. K., Gaffen, S. L., Liu, H., Rohowsky-Kochan, C., et al. (2011). 1,25-Dihydroxyvitamin D(3) ameliorates Th17 autoimmunity via transcriptional modulation of interleukin-17A. *Molecular and Cellular Biology*, *31*(17), 3653–3669. <https://doi.org/10.1128/MCB.05020-11>.
- Kleindorfer, D., Lindsell, C. J., Brass, L., Koroshetz, W., & Broderick, J. P. (2008). National US estimates of recombinant tissue plasminogen activator use: ICD-9 codes substantially underestimate. *Stroke*, *39*(3), 924–928. <https://doi.org/10.1161/STROKEAHA.107.490375>.
- Korf, H., Wenes, M., Stijlemans, B., Takiishi, T., Robert, S., Miani, M., et al. (2012). 1,25-Dihydroxyvitamin D3 curtails the inflammatory and T cell stimulatory capacity of macrophages through an IL-10-dependent mechanism. *Immunobiology*, *217*(12), 1292–1300. <https://doi.org/10.1016/j.imbio.2012.07.018>.
- Landel, V., Millet, P., Baranger, K., Loriod, B., & Feron, F. (2016). Vitamin D interacts with Esr1 and Igf1 to regulate molecular pathways relevant to Alzheimer's disease. *Molecular Neurodegeneration*, *11*, 22. <https://doi.org/10.1186/s13024-016-0087-2>.
- Langub, M. C., Herman, J. P., Malluche, H. H., & Koszewski, N. J. (2001). Evidence of functional vitamin D receptors in rat hippocampus. *Neuroscience*, *104*(1), 49–56.
- Lee, S., Evans, M. A., Chu, H. X., Kim, H. A., Widdop, R. E., Drummond, G. R., et al. (2015). Effect of a selective Mas receptor agonist in cerebral ischemia in vitro and in vivo. *PLoS ONE*, *10*(11), e0142087. <https://doi.org/10.1371/journal.pone.0142087>.
- Lee, J. H., O'Keefe, J. H., Bell, D., Hensrud, D. D., & Holick, M. F. (2008). Vitamin D deficiency an important, common, and easily treatable cardiovascular risk factor? *Journal of the American College of Cardiology*, *52*(24), 1949–1956. <https://doi.org/10.1016/j.jacc.2008.08.050>.
- Liesz, A., Suri-Payer, E., Veltkamp, C., Doerr, H., Sommer, C., Rivest, S., et al. (2009). Regulatory T cells are key cerebroprotective immunomodulators in acute experimental stroke. *Nature Medicine*, *15*(2), 192–199. <https://doi.org/10.1038/nm.1927>.
- Liu, P. T., Stenger, S., Li, H., Wenzel, L., Tan, B. H., Krutzik, S. R., et al. (2006). Toll-like receptor triggering of a vitamin D-mediated human antimicrobial response. *Science*, *311*(5768), 1770–1773. <https://doi.org/10.1126/science.1123933>.
- Lugg, S. T., Howells, P. A., & Thickett, D. R. (2015). Optimal vitamin D supplementation levels for cardiovascular disease protection. *Disease Markers*, *2015*, 1–10. <https://doi.org/10.1155/2015/864370>.
- Luo, W., Hershberger, P. A., Trump, D. L., & Johnson, C. S. (2013). 24-Hydroxylase in cancer: impact on vitamin D-based anticancer therapeutics. *Journal of Steroid Biochemistry and Molecular Biology*, *136*, 252–257. <https://doi.org/10.1016/j.jsbmb.2012.09.031>.

- Martorell, S., Hueso, L., Gonzalez-Navarro, H., Collado, A., Sanz, M. J., & Piqueras, L. (2016). Vitamin D receptor activation reduces angiotensin-II-induced dissecting abdominal aortic aneurysm in apolipoprotein E-knockout mice. *Arteriosclerosis, Thrombosis, and Vascular Biology*, *36*(8), 1587–1597. <https://doi.org/10.1161/ATVBAHA.116.307530>.
- Mattner, F., Smirolto, S., Galbiati, F., Muller, M., Di Lucia, P., Poliani, P. L., et al. (2000). Inhibition of Th1 development and treatment of chronic-relapsing experimental allergic encephalomyelitis by a non-hypercalcemic analogue of 1,25-dihydroxyvitamin D3. *European Journal of Immunology*, *30*(2), 498–508. [https://doi.org/10.1002/1521-4141\(200002\)30:2<498::AID-IMMU498>3.0.CO;2-Q](https://doi.org/10.1002/1521-4141(200002)30:2<498::AID-IMMU498>3.0.CO;2-Q).
- Merke, J., Milde, P., Lewicka, S., Hugel, U., Klaus, G., Mangelsdorf, D. J., et al. (1989). Identification and regulation of 1,25-dihydroxyvitamin D3 receptor activity and biosynthesis of 1,25-dihydroxyvitamin D3. Studies in cultured bovine aortic endothelial cells and human dermal capillaries. *Journal of Clinical Investigation*, *83*(6), 1903–1915. <https://doi.org/10.1172/jci114097>.
- Mozaffarian, D., Benjamin, E. J., Go, A. S., Arnett, D. K., Blaha, M. J., Cushman, M., et al. (2016). Heart disease and stroke statistics-2016 update: A report from the American Heart Association. *Circulation*, *133*(4), e38–e360. <https://doi.org/10.1161/CIR.0000000000000350>.
- Muthian, G., Raikwar, H. P., Rajasingh, J., & Bright, J. J. (2006). 1,25 Dihydroxyvitamin-D3 modulates JAK-STAT pathway in IL-12/IFN γ axis leading to Th1 response in experimental allergic encephalomyelitis. *Journal of Neuroscience Research*, *83*(7), 1299–1309. <https://doi.org/10.1002/jnr.20826>.
- Nashold, F. E., Nelson, C. D., Brown, L. M., & Hayes, C. E. (2013). One calcitriol dose transiently increases Helios + FoxP3 + T cells and ameliorates autoimmune demyelinating disease. *Journal of Neuroimmunology*, *263*(1–2), 64–74. <https://doi.org/10.1016/j.jneuroim.2013.07.016>.
- Naveilhan, P., Neveu, I., Wion, D., & Brachet, P. (1996). 1,25-Dihydroxyvitamin D3, an inducer of glial cell line-derived neurotrophic factor. *NeuroReport*, *7*(13), 2171–2175.
- Neveu, I., Naveilhan, P., Jehan, F., Baudet, C., Wion, D., De Luca, H. F., et al. (1994). 1,25-Dihydroxyvitamin D3 regulates the synthesis of nerve growth factor in primary cultures of glial cells. *Molecular Brain Research*, *24*(1–4), 70–76. [https://doi.org/10.1016/0169-328X\(94\)90119-8](https://doi.org/10.1016/0169-328X(94)90119-8).
- Oermann, E., Bidmon, H. J., Witte, O. W., & Zilles, K. (2004). Effects of 1 α ,25 dihydroxyvitamin D3 on the expression of HO-1 and GFAP in glial cells of the photothrombotically lesioned cerebral cortex. *Journal of Chemical Neuroanatomy*, *28*(4), 225–238. <https://doi.org/10.1016/j.jchemneu.2004.07.003>.
- Overbergh, L., Decallonne, B., Valckx, D., Verstuyf, A., Depovere, J., Laureys, J., et al. (2000). Identification and immune regulation of 25-hydroxyvitamin D-1- α -hydroxylase in murine macrophages. *Clinical and Experimental Immunology*, *120*(1), 139–146. <https://doi.org/10.1046/j.1365-2249.2000.01204.x>.
- Park, K.-Y., Chung, P.-W., Kim, Y. B., Moon, H.-S., Suh, B.-C., Won, Y. S., et al. (2015). Serum vitamin D status as a predictor of prognosis in patients with acute ischemic stroke. *Cerebrovascular Diseases*, *40*(1–2), 73–80. <https://doi.org/10.1159/000434691>.
- Pedersen, L. B., Nashold, F. E., Spach, K. M., & Hayes, C. E. (2007). 1,25-dihydroxyvitamin D3 reverses experimental autoimmune encephalomyelitis by inhibiting chemokine synthesis and monocyte trafficking. *Journal of Neuroscience Research*, *85*(11), 2480–2490. <https://doi.org/10.1002/jnr.21382>.
- Penna, G., Amuchastegui, S., Giarratana, N., Daniel, K. C., Vulcano, M., Sozzani, S., et al. (2007). 1,25-Dihydroxyvitamin D3 selectively modulates tolerogenic properties in myeloid but not plasmacytoid dendritic cells. *Journal of Immunology*, *178*(1), 145–153.
- Provvedini, D. M., Tsoukas, C. D., Deftos, L. J., & Manolagas, S. C. (1983). 1,25-Dihydroxyvitamin D3 receptors in human leukocytes. *Science*, *221*(4616), 1181–1183.
- Reeves, M. J., Arora, S., Broderick, J. P., Frankel, M., Heinrich, J. P., Hickenbottom, S., et al. (2005). Acute stroke care in the US: results from 4 pilot prototypes of the Paul Coverdell National Acute Stroke Registry. *Stroke*, *36*(6), 1232–1240. <https://doi.org/10.1161/01.STR.0000165902.18021.5b>.
- Sakaguchi, S., Wing, K., Onishi, Y., Prieto-Martin, P., & Yamaguchi, T. (2009). Regulatory T cells: how do they suppress immune responses? *International Immunology*, *21*(10), 1105–1111. <https://doi.org/10.1093/intimm/dxp095>.
- Schedel, M., Jia, Y., Michel, S., Takeda, K., Domenico, J., Joetham, A., et al. (2016). 1,25D3 prevents CD8(+)/Tc2 skewing and asthma development through VDR binding changes to the Cyp11a1 promoter. *Nature Communication*, *7*, 10213. <https://doi.org/10.1038/ncomms10213>.
- Shichita, T., Sugiyama, Y., Ooboshi, H., Sugimori, H., Nakagawa, R., Takada, I., et al. (2009). Pivotal role of cerebral interleukin-17-producing gammadeltaT cells in the delayed phase of ischemic brain injury. *Nature Medicine*, *15*(8), 946–950. <https://doi.org/10.1038/nm.1999>.
- Sloka, S., Silva, C., Wang, J., & Yong, V. W. (2011). Predominance of Th2 polarization by vitamin D through a STAT6-dependent mechanism. *Journal of Neuroinflammation*, *8*(1), 56. <https://doi.org/10.1186/1742-2094-8-56>.
- Sotirchos, E. S., Bhargava, P., Eckstein, C., Van Haren, K., Baynes, M., Ntranos, A., et al. (2016). Safety and immunologic effects of high- vs low-dose cholecalciferol in multiple sclerosis. *Neurology*, *86*(4), 382–390. <https://doi.org/10.1212/wnl.0000000000002316>.
- Spanier, J. A., Nashold, F. E., Olson, J. K., & Hayes, C. E. (2012). The Ifng gene is essential for Vdr gene expression and vitamin D(3)-mediated reduction of the pathogenic T cell burden in the central nervous system in experimental autoimmune encephalomyelitis, a multiple sclerosis model. *Journal of Immunology*, *189*(6), 3188–3197. <https://doi.org/10.4049/jimmunol.1102925>.
- Takeda, M., Yamashita, T., Sasaki, N., Nakajima, K., Kita, T., Shinohara, M., et al. (2010). Oral administration of an active form of vitamin D3 (calcitriol) decreases atherosclerosis in mice by inducing regulatory T cells and immature dendritic cells with tolerogenic functions. *Arteriosclerosis, Thrombosis, and Vascular Biology*, *30*(12), 2495–2503. <https://doi.org/10.1161/atvba.110.215459>.
- Taniura, H., Ito, M., Sanada, N., Kuramoto, N., Ohno, Y., Nakamichi, N., et al. (2006). Chronic vitamin D3 treatment protects against neurotoxicity by glutamate in association with upregulation of vitamin D receptor mRNA expression in cultured rat cortical neurons. *Journal of Neuroscience Research*, *83*(7), 1179–1189. <https://doi.org/10.1002/jnr.20824>.
- Taylor, A., Verhagen, J., Blaser, K., Akdis, M., & Akdis, C. A. (2006). Mechanisms of immune suppression by interleukin-10 and transforming growth factor- β : the role of T regulatory cells. *Immunology*, *117*(4), 433–442. <https://doi.org/10.1111/j.1365-2567.2006.02321.x>.
- Tu, W. J., Zhao, S. J., Xu, D. J., & Chen, H. (2014). Serum 25-hydroxyvitamin D predicts the short-term outcomes of Chinese patients with acute ischaemic stroke. *Clinical Science (London, England: 1979)*, *126*(5), 339–346. <https://doi.org/10.1042/cs20130284>.
- Turetsky, A., Goddeau, R. P., Jr., & Henninger, N. (2015). Low serum vitamin D is independently associated with larger lesion volumes after ischemic stroke. *Journal of Stroke and Cerebrovascular Diseases*, *24*(7), 1555–1563. <https://doi.org/10.1016/j.jstrokecerebrovasdis.2015.03.051>.
- Vantourout, P., & Hayday, A. (2013). Six-of-the-best: unique contributions of [gamma][delta] T cells to immunology. *Nature Reviews: Immunology*, *13*(2), 88–100. <https://doi.org/10.1038/nri3384>.

- Verma, R., & Kim, J. Y. (2016). 1,25-Dihydroxyvitamin D3 facilitates M2 polarization and upregulates TLR10 expression on human microglial cells. *Neuroimmunomodulation*, 23(2), 75–80. <https://doi.org/10.1159/000444300>.
- von Essen, M. R., Kongsbak, M., Schjerling, P., Olgaard, K., Odum, N., & Geisler, C. (2010). Vitamin D controls T cell antigen receptor signaling and activation of human T cells. *Nature Immunology*, 11(4), 344–349. <https://doi.org/10.1038/ni.1851>.
- Wang, Y., Ji, H., Tong, Y., & Zhang, Z. B. (2014). Prognostic value of serum 25-hydroxyvitamin D in patients with stroke. *Neurochemical Research*, 39(7), 1332–1337. <https://doi.org/10.1007/s11064-014-1316-0>.
- Won, S., Sayeed, I., Peterson, B. L., Wali, B., Kahn, J. S., & Stein, D. G. (2015). Vitamin D prevents hypoxia/reoxygenation-induced blood-brain barrier disruption via vitamin D receptor-mediated NF- κ B signaling pathways. *PLoS ONE*, 10(3), e0122821. <https://doi.org/10.1371/journal.pone.0122821>.
- Wong, M. S., Leisegang, M. S., Kruse, C., Vogel, J., Schurmann, C., Dehne, N., et al. (2014). Vitamin D promotes vascular regeneration. *Circulation*, 130(12), 976–986. <https://doi.org/10.1161/CIRCULATIONAHA.114.010650>.
- Yang, L., Brozovic, S., Xu, J., Long, Y., Kralik, P. M., Waigel, S., et al. (2011). Inflammatory gene expression in OVE26 diabetic kidney during the development of nephropathy. *Nephron Experimental Nephrology*, 119(1), e8–e20.
- Yao, T., Ying, X., Zhao, Y., Yuan, A., He, Q., Tong, H., et al. (2015). Vitamin D receptor activation protects against myocardial reperfusion injury through inhibition of apoptosis and modulation of autophagy. *Antioxidants & Redox Signaling*, 22(8), 633–650. <https://doi.org/10.1089/ars.2014.5887>.
- Yilmaz, G., Arumugam, T. V., Stokes, K. Y., & Granger, D. N. (2006). Role of T lymphocytes and interferon-gamma in ischemic stroke. *Circulation*, 113(17), 2105–2112.
- Zeitelhofer, M., Adzemovic, M. Z., Gomez-Cabrero, D., Bergman, P., Hochmeister, S., N'Diaye, M., et al. (2017). Functional genomics analysis of vitamin D effects on CD4+ T cells in vivo in experimental autoimmune encephalomyelitis. *Proceedings of the National Academy of Sciences of the United States of America*. <https://doi.org/10.1073/pnas.1615783114>.
- Zhang, Y., Leung, D. Y., Richers, B. N., Liu, Y., Remigio, L. K., Riches, D. W., et al. (2012). Vitamin D inhibits monocyte/macrophage proinflammatory cytokine production by targeting MAPK phosphatase-1. *Journal of Immunology*, 188(5), 2127–2135. <https://doi.org/10.4049/jimmunol.1102412>.