

RESEARCH ARTICLE

Negligible differences in metabolism and thermal tolerance between diploid and triploid Atlantic salmon (*Salmo salar*)

Alyssa J. Bowden^{1,2,*}, Sarah J. Andrewartha², Nick G. Elliott^{1,2}, Peter B. Frappell¹ and Timothy D. Clark^{1,2,3}

ABSTRACT

The mechanisms that underlie thermal tolerance in aquatic ectotherms remain unresolved. Triploid fish have been reported to exhibit lower thermal tolerance than diploids, offering a potential model organism to better understand the physiological drivers of thermal tolerance. Here, we compared triploid and diploid juvenile Atlantic salmon (*Salmo salar*) in freshwater to investigate the proposed link between aerobic capacity and thermal tolerance. We measured specific growth rates (SGR) and resting (aerobic) metabolic rates (RMR) in freshwater at 3, 7 and 9 weeks of acclimation to 10, 14 and 18°C. Additionally, maximum metabolic rates (MMR) were measured at 3 and 7 weeks of acclimation, and critical thermal maxima (CT_{max}) were measured at 9 weeks. Mass, SGR and RMR differed between ploidies across all temperatures at the beginning of the acclimation period, but all three metrics were similar across ploidies by week 7. Aerobic scope (MMR–RMR) remained consistent across ploidies, acclimation temperatures and time. At 9 weeks, CT_{max} was independent of ploidy, but correlated positively with acclimation temperature despite the similar aerobic scope between acclimation groups. Our findings suggest that acute thermal tolerance is not modulated by aerobic scope, and the altered genome of triploid Atlantic salmon does not translate to reduced thermal tolerance of juvenile fish in freshwater.

KEY WORDS: Critical thermal maxima, Elevated temperature, Oxygen consumption, Salmonids, Ploidy

INTRODUCTION

Triploid fish have been proposed as useful experimental models to understand the mechanisms underlying environmental tolerance because their altered genome may influence tolerance levels under challenging conditions (Maxime, 2008). Indeed, triploid brown trout (*Salmo trutta*) exhibited higher mortality rates than those of diploids when exposed to a high temperature challenge (18°C) for 3 weeks, and triploid mortalities reached 50% after 12 weeks at this temperature (Altimiras et al., 2002). Moreover, triploid rainbow trout (*Oncorhynchus mykiss*) suffered immediate mortalities when exposed to 21°C, and a total mortality of 69% was recorded after 3 weeks compared with a mortality of 39% in diploid conspecifics (Ojolick et al., 1995).

Such observations raise the possibility of using triploids to elucidate the physiological mechanisms underlying temperature

tolerance in fish. Triploid fish are typically produced by subjecting eggs to one or more pressure shocks within the first hour or two of fertilisation, resulting in the retention rather than extrusion of a polar body that gives rise to a third chromosome (Teskeredžić et al., 1993). Triploids compensate for this extra genetic material by having larger but fewer cells than their diploid counterparts, resulting in the two ploidies achieving similar sizes as adults (Swarup, 1959; Small and Benfey, 1987). Growth rates of triploids, however, have been inconsistent across studies and have been the same as, greater than or less than the growth rates of their diploid counterparts (Galbreath et al., 1994; McGeachy et al., 1995; McCarthy et al., 1996).

Furthermore, triploids are thought to have similar haematocrit (packed erythrocyte volume) as that of diploids but lower haemoglobin concentration, subsequently reducing blood oxygen carrying capacity (Benfey and Sutterlin, 1984a; Benfey et al., 1984; Graham et al., 1985). Blood oxygen transport of triploids may be further reduced owing to the smaller surface area to volume ratio of the enlarged erythrocytes, which could inhibit oxygen diffusion dynamics (Sadler et al., 2000). In light of the proposed link between oxygen transport capacity and thermal performance (Pörtner and Knust, 2007), a logical extension is that inferior oxygen transport capacity in triploids drives their reportedly lower thermal tolerance. Having said that, the role of oxygen transport capacity in governing thermal tolerance has been debated and requires additional investigation using different model organisms (Clark et al., 2013; Brijs et al., 2015; Ern et al., 2016).

Despite the apparent reduction in oxygen carrying capacity of triploid blood, contrasting results exist regarding their aerobic metabolic rates at both high and low acclimation temperatures. Higher routine metabolism has been reported in triploid Atlantic salmon (*Salmo salar* Linnaeus 1758) at an acclimation temperature of 12°C, but lower routine metabolism at a higher temperature (18°C) was considered to be indicative of lower thermal tolerance in triploids (Atkins and Benfey, 2008). In contrast, no differences in routine metabolic rate were found between diploid and triploid Atlantic salmon at 15°C, or between diploid and triploid rainbow trout at 19°C (Benfey and Sutterlin, 1984b; Oliva-Teles and Kaushik, 1990). Although maximum metabolic rate (MMR) has received less attention, a study on Atlantic salmon reported no differences in MMR between ploidies at 15°C (Lijalad and Powell, 2009). Scant data and disparate findings point to a requirement for more comprehensive investigations to determine whether aerobic capacity may be causally linked with acute and/or chronic thermal tolerance across ploidies (Benfey et al., 1997; Galbreath et al., 2006).

Here, we investigated acute and chronic thermal tolerance of diploid and triploid juvenile Atlantic salmon with an aim to clarify some of the conflicting findings reported previously. Specifically, our objectives were to compare the growth performance, resting metabolic rate (RMR), MMR and aerobic scope at three acclimation temperatures (10, 14 and 18°C), and then to quantify routine

¹Institute of Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania 7000, Australia. ²CSIRO Agriculture and Food, Hobart, Tasmania 7000, Australia. ³Deakin University, School of Life and Environmental Sciences, Geelong, Victoria 3216, Australia.

*Author for correspondence (alyssa_bowden@outlook.com)

 A.J.B., 0000-0001-6024-1891

metabolic rate during assessments of critical thermal maxima (CT_{max}) in the different acclimation groups. We tested the hypothesis that triploids exhibit lower chronic and acute thermal tolerance than those of diploids, and that these differences are at least partly explained by lower aerobic capacity of triploids.

MATERIALS AND METHODS

Animal husbandry

Diploid and triploid Atlantic salmon ($n=35$ and $n=40$, respectively) were sourced from the Salmon Enterprises of Tasmania Pty Ltd (SALTAS) freshwater hatchery in Wayatinah, Tasmania (water temperature $\sim 9^{\circ}\text{C}$) and transported to Hobart, Tasmania. All experiments were conducted at the Aquaculture Physiology laboratory at CSIRO under the animal ethics permit A0013794. Both groups were from the 2015 commercial spawning and had been incubated at $\sim 8^{\circ}\text{C}$. The fish were separated by ploidy into two 200 l tanks that were supplied by a freshwater recirculating system held at 10°C . After 7 days of recovery from transport, the fish were individually weighed (mean \pm s.e.m.; diploids: 46.63 ± 2.31 g, triploids: 68.75 ± 2.08 g) and their fork lengths were measured (diploids: 163.41 ± 2.95 mm, triploids: 188.63 ± 2.13 mm). Additionally, the fish were tagged intraperitoneally with a passive integrated transponder (PIT) to track individual performance and with coloured elastomer in the adipose fin to visually discern ploidy (yellow for diploids and green for triploids), before being returned to their respective tanks for 3 days to recover from tagging. Thereafter, mixed groups of diploids and triploids ($n=5-8$ per ploidy) were assigned to six 68 l tanks ($n=11-13$ per tank) with recirculating freshwater and maintained at 10°C for 3 weeks.

Water temperature was subsequently raised to 14°C (to represent current-day summer sea temperatures of $14-15^{\circ}\text{C}$ in southeast Tasmania) in two tanks and to 18°C (forecast of regional sea temperatures under a business-as-usual emissions scenario) in two other tanks ($2^{\circ}\text{C day}^{-1}$) using 600 W titanium heaters with a digital Nema thermostat (Aquasonic, Wauchope, NSW, Australia). The remaining two tanks were maintained at 10°C to represent current-day winter temperatures. For each temperature, the heaters were placed in a common sump supplying the two tanks and the water was maintained at $\pm 0.5^{\circ}\text{C}$ of the desired temperature. Water was changed periodically to maintain water quality (ammonia < 0.7 mg l^{-1} ; nitrite < 0.2 mg l^{-1} ; nitrate < 20 mg l^{-1}). The fish were fed to satiation daily, and were allowed to acclimate for 3 weeks to their respective temperatures prior to the start of experiments. Dissolved oxygen was maintained above 85% air saturation and the lighting regime was kept at 10 h:14 h light:dark. During the ~ 4 weeks following tagging, there were two mortalities in the 10°C acclimation group (two diploids), two in the 14°C acclimation group (one diploid, one triploid) and one in the 18°C acclimation group (one triploid). We attribute these mortalities to transport and/or tagging effects rather than to temperature.

Respirometry

Oxygen consumption rates of individual fish were measured in six intermittently closed respirometers (2.8 l), following previously described methods (Clark et al., 2013). The respirometers were submerged in a 160 l temperature-controlled water bath that was maintained at $> 90\%$ air saturation with compressed air. The temperature of the water bath was adjusted as necessary to match the acclimation temperature of the fish involved in the respirometry trial. Each respirometer was flushed every 15 min by a pump at 0.5 l min^{-1} to replenish the oxygen levels. Water was continuously mixed by a submersible pump (0.1 l min^{-1}) within a closed

recirculation loop. Dissolved oxygen concentration was measured in each respirometer by fibre optic oxygen probes (FireSting O_2 , Pyroscience, Aachen, Germany) sealed in the recirculation loop, and recorded at 5 s intervals using an eight channel PowerLab/8sp and LabChart 7 Pro software (ADInstruments Pty Ltd, Bella Vista, NSW, Australia).

Resting and maximum oxygen consumption rates were measured at 3 and 7 weeks, after the acclimation temperatures had been established. Fish from an acclimation temperature treatment were fasted for 24 h, and then dip-netted out of their holding tanks in the morning and lightly anaesthetised with 0.02 ml l^{-1} Aqu-i-S (50% active isoeugenol); their mass, length and PIT tag number were recorded. The fish were then placed in individual respirometers at their respective acclimation temperatures, and RMR was determined from oxygen consumption measurements (see below) during the 17–20 h overnight recovery. The following morning, the fish were removed from the chambers and individually exercised in a 40 l round swim tank. Water temperature in the swim tank was maintained using an Aquasonic heater and Nema thermostat, and dissolved oxygen levels were maintained above 95% air saturation through aeration. A chase protocol was used to elevate oxygen consumption to the maximum level (Clark et al., 2013; Norin and Clark, 2016; Killen et al., 2017). Briefly, each fish was individually chased by hand for 2 min, before being immediately placed back in the respirometer for the measurement of the maximum oxygen consumption rate as a proxy for MMR. All fish reached exhaustion after 2 min of vigorous chasing in a preliminary set of experiments ($n=6$). Exhaustion was recorded when the fish no longer swam away when being chased and tapped on the tail. Oxygen consumption measurements were continued for 30 min to ensure that the maximum rate of decline in oxygen concentration was captured. The fish were returned to their respective holding tanks after the respirometry protocol. Notably, a pump failure in one of the 14°C acclimation tanks, between weeks 3 and 7, halved the sample size at this acclimation temperature and resulted in $n=6$ diploids and $n=8$ triploids being measured at 14°C thereafter.

Critical thermal maxima

At 9 weeks of acclimation, fish were placed in respirometers, as described above, for measurement of RMR, but a CT_{max} protocol, rather than an exercise protocol, was conducted the following morning. The temperature of the water bath containing the respirometers was increased stepwise by 2°C every 75 min (ramped increase over 15 min, then held stable for 60 min to ensure stable oxygen measurements), and routine metabolic rate was measured over two 15 min periods during each 60 min interval once temperature had plateaued (similar to Brijs et al., 2015). The protocol was ceased when the fish displayed loss of equilibrium (LOE), which was defined as the inability to right themselves after 10 s. The time and temperature at LOE were recorded and each fish was immediately killed via anaesthetic overdose. Blood was sampled from the caudal vasculature in a 1 ml heparinised syringe, transferred to a 2 ml Eppendorf tube and immediately placed on ice until processing (< 1 h).

Dissections and ploidy verification

Following LOE and blood sampling, the fish were dissected, and the ventricle, liver and spleen masses were recorded (Ohaus Scout Pro Portable Electronic Balance, Parsippany, NJ, USA). A sample (2 μl) of blood was smeared on a glass slide, dried and stained in Dif Quik (Sigma-Aldrich, Castle Hill, NSW, Australia) for ploidy verification. The blood smears were examined using light

microscopy (Leitz Wetzlar, Germany) and photographed using a Leica DFC310 FX microscope camera connected to a PC with Leica Application Suite Version 4.0.0 software (Leica Microsystems Limited, Heerbrugg, Switzerland). The photographs were processed using ImageJ 1.48v (Wayne Rasband, National Institutes of Health, Bethesda, MD, USA). The nucleus major axis was measured for at least 50 randomly chosen erythrocytes per individual, and triploids and diploids were statistically separated through cluster analysis on the mean nuclear length (Benfey et al., 1984).

Data analyses

Specific growth rate (SGR) was calculated using Eqn 1:

$$\text{SGR} = \{[\ln(M_f) - \ln(M_i)] \times t^{-1}\} \times 100, \quad (1)$$

where M_f and M_i are the final and initial masses, respectively, of an individual and t is the elapsed time between mass measurements in days.

Oxygen consumption rates ($\text{mg O}_2 \text{ kg}^{-1} \text{ h}^{-1}$) were calculated using slopes derived from linear regressions between oxygen concentration and time during each sealed event in the chamber and accounting for the volume of the respirometer, as in Eqn 2:

$$\dot{M}_{\text{O}_2} = \frac{(\Delta\text{O}_2/\Delta t)(P_B - P_V) \times C_{\text{O}_2} \times V \times 0.2093}{M_b}, \quad (2)$$

where ΔO_2 is the change in oxygen concentration within the respirometer over the change in time in hours (Δt), P_B is the barometric pressure in kPa, P_V is the water vapour pressure (kPa, Antoine equation), C_{O_2} is the calculated oxygen capacitance of freshwater at the acclimation temperature ($\text{mg l}^{-1} \text{ kPa}^{-1}$; Dejours, 1981), V is the volume of the respirometer minus that of the fish (assuming 1 kg wet mass=1 l) in l, 0.2093 is the fractional concentration of oxygen in well-aerated water and M_b is the body mass (kg). Note that most of the statistical analyses were conducted using oxygen consumption rate in $\text{mg O}_2 \text{ h}^{-1}$, and body mass was included as a covariate (details below).

RMR was determined as the mean of the lowest 10% of oxygen consumption values throughout the measuring period (17–20 h), excluding outliers [values ± 2 s.d. from the mean (Norin et al., 2014)]. MMR was calculated from the slope immediately after the exhaustive chase protocol, which was always found to be the highest. Absolute aerobic scope was calculated by subtracting RMR from MMR, and factorial aerobic scope was calculated by dividing MMR by RMR.

CT_{max} was calculated by modifying the critical swimming speed equation from Brett (1964) into Eqn 3:

$$\text{CT}_{\text{max}} = T_f + \left(\frac{t_f}{t_i} \times T_i \right), \quad (3)$$

where T_f is the highest temperature the fish endured for the full time period, t_f is the time the fish lasted at its final temperature step, t_i is the prescribed time for each temperature (i.e. 60 min) and T_i is the incremental temperature increase (i.e. 2°C).

Temperature coefficients (Q_{10}) were calculated to quantify the influence of acclimation temperature on the metabolism of diploids and triploids, using mean RMR values in Eqn 4:

$$Q_{10} = \left(\frac{R_2}{R_1} \right)^{10^\circ / (T_2 - T_1)}, \quad (4)$$

where R_2 and R_1 are the mean RMR values corresponding to two acclimation temperatures (T_2 and T_1 , respectively). Q_{10} values were

calculated between 10 and 18°C for each ploidy at each acclimation time point (3, 7 and 9 weeks).

All data are presented as means \pm 95% confidence intervals in the figures and the text. Organ weights were analysed using ANCOVAs (see below) and presented as body mass adjusted means from the ANCOVA outputs.

Statistical analyses

Mass and SGR data were analysed using a series of two-way ANOVAs and ANCOVAs, respectively. The two factors (ploidy and acclimation temperature) were analysed within each time point and α for significance was set at 0.017 (i.e. 0.05 divided by three tests). ANCOVAs for SGR used the initial mass as the covariate. Differences in metabolic rates were analysed using general linear mixed models, testing metabolic rate ($\text{mg O}_2 \text{ h}^{-1}$) against sampling time point, ploidy and acclimation temperature, with mass as a covariate and accounting for repeated measures on an individual. Results are reported from the repeated measures analyses using F tests (type III Wald F tests with Kenward–Roger degrees of freedom approximation). Where applicable, *post hoc* tests for pairwise comparisons using Bonferroni corrections were utilised to investigate the differences between ploidies within time points, and the differences across time points within ploidy. CT_{max} data were analysed using a two-way ANOVA because there was no time point variable. Measurements of routine metabolic rate ($\text{mg O}_2 \text{ kg}^{-1} \text{ h}^{-1}$) during the CT_{max} protocol were averaged for each 60 min interval, and then analysed with a linear mixed effects model to test the effect of temperature (covariate) increase on metabolic rate between ploidies (factor), with individual as a random factor to control for repeated measures. Organ weights were analysed using ANCOVAs testing the absolute values against ploidy and acclimation temperature, with mass as a covariate. Significance was accepted at $P < 0.05$ unless otherwise indicated (for multiple tests), and Bonferroni-corrected *post hoc* tests were conducted on covariate-adjusted means where applicable. All analyses were conducted using R Studio (Version 1.0.136, <https://www.rstudio.com/>), with R packages car (Fox and Weisberg, 2011), nlme (Pinheiro et al., 2016) and lsmeans (Lenth, 2016).

RESULTS

Ploidy verification

The erythrocyte nuclei measurements confirmed that the ploidy of all fish was correctly classified. In this study, the major axis of the nucleus in triploids was 1.48 times longer than that of diploids, which compares favourably with a value of 1.26 times longer reported in Benfey et al. (1984).

Survival, mass and growth

There were no natural mortalities in either ploidy at the 10 or 14°C acclimation temperatures once the experiment commenced. There were two triploid mortalities at 18°C between weeks 7 and 9, although the low mortality rate makes it difficult to attribute these deaths to a lower chronic thermal tolerance of triploids.

Triploid mass was ~ 1.5 times greater than that of diploids at week 0 across all acclimation temperatures (10°C: $F_{1,18}=9.01$, $P=0.008$; 14°C: $F_{1,21}=37.58$, $P<0.001$; 18°C: $F_{1,23}=11.91$, $P=0.002$; Fig. 1A). Triploids and diploids acclimated to 10 and 18°C had similar sizes at week 3, and remained similar at subsequent time points. However, triploid fish acclimated to 14°C continued to be larger than diploids at week 3 ($F_{1,21}=6.93$, $P=0.016$), but the masses of the ploidies converged by week 7. The temporal dynamics in fish mass across the ploidies and acclimation temperatures were consistent with SGR of the different groups (Fig. 1B).




Fig. 1. Growth dynamics in diploid and triploid Atlantic salmon. (A) Mass and (B) specific growth rate (SGR) for diploid (grey) and triploid (black) fish during 9 weeks of temperature acclimation to 10, 14 and 18°C. Sample sizes are in parentheses in A. All values are means±95% confidence intervals and are positioned side by side to reduce overlap for clarity. Asterisks denote significance between ploidy groups based on ANOVAs (mass) and ANCOVAs (SGR) with α reduction for multiple testing (see Materials and methods). Note that in B, values represent SGR between time points (e.g. from 0 to 3 weeks) and therefore could not be calculated for week 0 (n.a.).

Relative ventricle, liver and spleen masses were similar in diploids and triploids across all acclimation temperatures. Not surprisingly, gonad mass was significantly greater in diploids than in the inherently sterile triploids at all acclimation temperatures (10°C: $F_{1,17}=54.32$, $P<0.001$; 14°C: $F_{1,6}=17.68$, $P=0.006$; 18°C: $F_{1,13}=56.77$, $P<0.001$; Table 1).

Metabolic rates

As expected, there was a general increase in RMR with acclimation temperature from ~55 to ~125 mg kg⁻¹ h⁻¹ in both ploidies [Q_{10} between 10 and 18°C (week 3, week 7 and week 9, respectively)=1.75, 1.75 and 2.31 for diploids; 1.87, 1.73 and 1.99 for triploids]. Counter to our expectation that RMR would decrease as fish progressively acclimated to the higher temperatures, there were some significant increases in RMR between weeks 3 and 7 (~75 to ~115 mg kg⁻¹ h⁻¹) in both

ploidies at 14°C (diploids: $F_{2,79}=10.96$, $P=0.004$; triploids: $F_{2,79}=34.87$, $P<0.001$) and in triploids at 18°C (~96 to ~120 mg kg⁻¹ h⁻¹; $F_{2,79}=21.61$, $P<0.001$; Fig. 2A).

On average, RMR was 15% higher in diploids than in triploids across all acclimation temperatures during the week 3 measurements (10°C: $F_{1,62}=10.20$, $P=0.002$; 14°C: $F_{1,62}=7.84$, $P=0.007$; 18°C: $F_{1,62}=12.72$, $P=0.001$; Fig. 2A). Diploids also had a higher RMR at 10°C during the week 7 measurements (10°C: $F_{1,62}=6.39$, $P=0.014$), but RMR was similar in ploidy groups at 14 and 18°C. There were no differences in RMR between diploids and triploids during the week 9 measurements.

Absolute aerobic scope and MMR were generally stable across temperatures (~400 and ~500 mg kg⁻¹ h⁻¹, respectively) for both ploidies (Fig. 2B,C), whereas there was a tendency for a decline of ~25% in factorial aerobic scope across the range of acclimation temperatures (Fig. 2D). Within ploidies, there were small but

Table 1. Body mass adjusted means (from ANCOVA) for the organ masses of diploid and triploid Atlantic salmon acclimated to three different temperatures

Measurement	Acclimation temperature					
	10°C		14°C		18°C	
	Diploid	Triploid	Diploid	Triploid	Diploid	Triploid
Ventricle (%)	0.077±0.007	0.076±0.007	0.092±0.012	0.091±0.012	0.069±0.006	0.070±0.005
Liver (%)	0.858±0.110	0.943±0.110	1.054±0.221	1.053±0.204	0.636±0.101	0.788±0.093
Spleen (%)	0.084±0.018	0.101±0.019	0.184±0.068	0.093±0.049	0.073±0.018	0.069±0.017
Gonad (%)	0.193±0.023*	0.075±0.023	0.217±0.042*	0.093±0.059	0.239±0.034*	0.046±0.044

Values are presented as a percentage of body mass and are means±95% confidence intervals. Asterisks denote significant differences between ploidy groups within a temperature (* $P<0.05$).




Fig. 2. Metabolic rates and aerobic scope in Atlantic salmon.

(A) Minimum oxygen consumption [as a proxy for resting metabolic rate (RMR)], (B) maximum oxygen consumption [as a proxy for maximum metabolic rate (MMR)], (C) absolute aerobic scope and (D) factorial aerobic scope for diploid (grey) and triploid (black) fish measured during acclimation to 10, 14 and 18°C. Values are means±95% confidence intervals. Significance between ploidies is denoted by asterisks (* $P<0.05$), and differences between measuring time points (weeks) within a ploidy are signified by different lowercase letters (Bonferroni-adjusted P -values for pairwise comparisons). See Fig. 1A for sample sizes.

significant increases in MMR between weeks 3 and 7 at 14 and 18°C, and an increase in absolute aerobic scope between weeks 3 and 7 at 18°C (MMR at 14°C, diploids: $F_{1,40}=4.63$, $P=0.038$; triploids: $F_{1,40}=7.85$, $P=0.008$; 18°C, diploids: $F_{1,40}=17.04$, $P<0.001$; triploids: $F_{1,40}=10.83$, $P=0.002$; aerobic scope at 18°C, diploids: $F_{1,40}=15.85$, $P<0.01$; triploids: $F_{1,40}=4.46$, $P=0.041$; Fig. 2B,C).

MMR and absolute aerobic scope were similar in ploidies at all acclimation temperatures within both measuring time points (weeks 3 and 7) (Fig. 2B,C). Factorial aerobic scope, however, was higher in the triploids than in the diploids within the 10°C acclimation group at week 3 ($F_{1,18}=11.29$, $P=0.047$; Fig. 2D) and within the 14°C acclimation group at week 7 ($F_{1,11}=6.62$, $P=0.026$). There was some evidence of factorial aerobic scope decreasing between weeks 3 and 7 in triploids at 10°C (10°C: $F_{1,13.4}=11.45$, $P=0.001$; Fig. 2D).

Critical thermal maxima

CT_{max} did not differ between ploidies ($F_{1,51}=1.76$, $P=0.190$; Fig. 3A), but increased with acclimation temperature from ~26 to ~29°C ($F_{2,51}=66.27$, $P<0.001$). Thus, there was generally an improvement of ~0.4°C in CT_{max} for every 1°C increase in acclimation temperature. Within diploids, CT_{max} was similar in fish acclimated to 10 and 14°C, but higher for fish acclimated to 18°C ($P_{10-18°C}<0.001$, $P_{14-18°C}<0.001$). Thermal effects were more consistent for triploids, in which CT_{max} increased significantly between the 10 and 14°C acclimation groups and between the 14 and 18°C acclimation groups ($P_{10-14°C}=0.016$, $P_{14-18°C}=0.001$).

Consistent with the findings for RMR described above, diploids had a higher routine metabolic rate than that of triploids during the CT_{max} protocol, when commencing at the acclimation temperatures of 14 and 18°C. In contrast, there were no differences in routine




Fig. 3. Critical thermal maxima (CT_{max}) and metabolic measurements in Atlantic salmon. (A) CT_{max} for diploid (grey) and triploid (black) fish across acclimation temperatures. Values are means±95% confidence intervals. Lower case letters denote differences within a ploidy across acclimation temperatures. (B) Oxygen consumption rate during the CT_{max} protocol. Values are means±95% confidence intervals, fitted with exponential regressions with the following equations. Diploids at 10°C: $y=42.385(0.088) \times e^{0.094(0.004)x}$ ($R^2=0.84$); triploids at 10°C: $y=50.135(0.099) \times e^{0.085(0.004)x}$ ($R^2=0.81$); diploids at 14°C: $y=42.571(0.114) \times e^{0.099(0.005)x}$ ($R^2=0.92$); triploids at 14°C: $y=30.291(0.133) \times e^{0.107(0.006)x}$ ($R^2=0.86$); diploids at 18°C: $y=63.089(0.186) \times e^{0.070(0.007)x}$ ($R^2=0.60$) and triploids at 18°C: $y=45.139(0.157) \times e^{0.077(0.006)x}$ ($R^2=0.68$). P -values represent significance between the two regressions. Numbers in parentheses indicate when sample sizes decreased.

metabolic rate between the ploidy groups acclimated to 10°C (10°C: $F_{1,20}=0.002$, $P=0.969$; 14°C: $F_{1,11}=5.115$, $P=0.045$; 18°C: $F_{1,20}=6.676$, $P=0.018$; Fig. 3B).

DISCUSSION

In contrast with our initial hypothesis, triploids performed similarly to their diploid counterparts when held chronically at three acclimation temperatures, and during the CT_{max} challenge. Our finding of similar thermal performance across ploidies is consistent with some previously published studies on Atlantic salmon (Ojolic et al., 1995; McGeachy et al., 1995), although there is some evidence for species differences (Ojolic et al., 1995; McGeachy

et al., 1995; O'Flynn et al., 1997; Altimiras et al., 2002). Ploidy differences in the present study were observed for mass, SGR and RMR, but not for MMR and aerobic scope (Figs 1, 2). We discuss below how these parameters may interact to result in subtle but important differences between ploidies.

Growth and metabolism

As expected, RMR increased with acclimation temperature in both ploidies, a trend that is well documented for fishes (Fry, 1971; Clarke and Johnston, 1999; Gillooly et al., 2001). Interestingly, there was a difference between the ploidies for RMR within an acclimation temperature in the first periods of the study (weeks 3 and 7), but not at the end (week 9). This is similar to the temporal trends in mass and SGR (Figs 1, 2A), indicating that there may be some common mechanisms between the measured parameters. Indeed, food consumption, growth and RMR are thought to be intrinsically linked (Metcalf et al., 1995; Pedersen, 1997; Yamamoto et al., 1998; Norin et al., 2016; Van Leeuwen et al., 2012).

The lower RMR of triploids indicates a lower maintenance cost per unit body mass, which should theoretically translate to higher growth rates for a given energy intake, assuming equivalent assimilation efficiencies. However, diploids rather than triploids displayed a higher SGR in the earlier weeks of the experiment, when the RMR of triploids was lower. It is possible that these counterintuitive findings are a consequence of fish size, as growth rates are known to be negatively related to body size in fish (Iwama, 1996). Indeed, when diploids caught up to triploids in mass by week 7, SGR and RMR also converged, supporting the idea that it was fish size rather than ploidy that drove the differences in the earlier weeks of the experiment.

Nevertheless, it is also possible that behavioural differences between the ploidies played some role in the counterintuitive findings. Specifically, diploid salmonids have been reported to outperform their triploid counterparts when raised in mixed ploidy populations, resulting in improved growth of diploids (Carter et al., 1994; Galbreath et al., 1994). This potential behavioural difference between ploidies may explain why investigations into growth in triploids and their diploid conspecifics have been largely inconclusive, with triploid Atlantic salmon being reported to grow faster, slower and similarly compared with their diploid counterparts (Galbreath et al., 1994; McGeachy et al., 1995; McCarthy et al., 1996). That is, the disparate findings may be largely attributed to husbandry practices such as keeping the ploidies separate versus mixing them in a common tank (Galbreath et al., 1994; Maxime, 2008). For example, Atlantic salmon triploids grew at a faster rate and had a higher mass when kept in a tank separate from diploids, whereas diploids outperformed the triploids when the two groups were mixed (Galbreath et al., 1994). In instances where diploids outperformed triploids in mixed tanks, it has been attributed to a more aggressive nature of diploids (Galbreath et al., 1994). In this context, intraspecific competition and dominance status have been correlated with standard metabolic rate, such that individuals with an inherently higher standard metabolic rate (such as the diploids in the present study) are stimulated to eat more food, and thus be more competitive/aggressive (Metcalf et al., 1995; Cutts et al., 1998; Norin et al., 2016). In any event, similar growth rates have also been observed in ploidies of Atlantic salmon and coho salmon (*Oncorhynchus kisutch*) when the ploidies were mixed in tanks, highlighting that further controlled experiments are required on this topic to tease apart the roles of ploidy, body size, RMR and behaviour (Johnson et al., 1986; Carter et al., 1994).

Acute thermal tolerance and aerobic capacity

Ploidy did not influence the CT_{max} of Atlantic salmon in this study, despite triploids from the 14 and 18°C acclimated groups maintaining a slightly lower routine metabolic rate during the temperature ramp (Fig. 3). The lower routine metabolic rate was consistent with the lower RMR observed for triploids in the earlier weeks of acclimation (Fig. 2A). Despite the common observation that triploids suffer higher mortality at elevated temperatures, few studies have found ploidy differences in CT_{max} protocols with a variety of heating rates (e.g. 2°C h⁻¹, 15°C h⁻¹, 2°C day⁻¹) and species (e.g. rainbow trout, brook charr, brook trout) (Benfey et al., 1997; Galbreath et al., 2006; Maxime, 2008; Scott, 2012; Scott et al., 2015). However, the CT_{max} experiments conducted here and in previous studies used juvenile fish, and it is known that smaller individuals can be more thermally tolerant than their adult conspecifics (Clark et al., 2012; Messmer et al., 2016; Clark et al., 2017). Furthermore, adult salmonids face other energetically costly physiological processes, such as sexual maturation and ion regulation in a hypertonic environment upon their movement from freshwater to saltwater during later life stages (Bœuf and Payan, 2001). Although the enlarged erythrocytes of triploids do not seem to impair oxygen transport capacity, they could hinder other critical functions like ion regulation. Therefore, it is possible that experimental results from juvenile triploids in freshwater may not be directly applicable to adult triploids in marine environments.

Another consideration is that the present study investigated metabolism and CT_{max} exclusively under normoxic conditions, whereas salmon in wild or cultured environments may be subjected to periods of hypoxia. In this context, upper thermal limits have been proposed to be oxygen limited in normoxic environments (Pörtner and Knust, 2007). Nevertheless, CT_{max} of red drum (*Sciaenops ocellatus*) and marine lumpfish (*Cyclopterus lumpus*) were found to be independent of oxygen availability over a wide range of oxygen tensions (Ern et al., 2016). Indeed, CT_{max} was not affected by oxygen availability until close to (in lumpfish) or below (in red drum) the species-specific critical oxygen tensions (P_{crit}) despite significant decreases in aerobic scope (72 and 89% reductions, respectively) (Ern et al., 2016). Given that it would be rare for Atlantic salmon in natural or cultured environments to experience oxygen levels below their P_{crit} (~35% air saturation for the size used here; Barnes et al., 2011), our findings should be applicable across the range of oxygen levels Atlantic salmon typically experience throughout their lifecycle.

Interestingly, a larger relative ventricle mass has been correlated with higher temperature tolerance in fish (Ozolina et al., 2016). It has been suggested that a deterioration in cardiovascular performance may be causal to the upper thermal tolerance limits of fishes (Lannig et al., 2004; Clark et al., 2008; Farrell, 2009). Indeed, enhancing oxygen availability to the heart of European perch has been shown to play a role in maintaining stroke volume at critically high temperatures. Nevertheless, a concurrent decline in heart rate was likely to reflect direct temperature effects rather than an oxygen limitation (Ekström et al., 2016). As ventricle mass is positively correlated with stroke volume (Graham and Farrell, 1989), and we did not see any differences in ventricle mass between ploidies in the present study (Table 1), we suggest that stroke volume was similar in triploids and diploids at the three acclimation temperatures. As such, we did not find any evidence that enhanced cardiovascular performance played any role in increasing CT_{max} with acclimation temperature (Fig. 3). Contrasting results for relative ventricle mass between ploidies have been reported, ranging from significantly heavier in Atlantic salmon triploids, to

similar among ploidies for rainbow trout and Atlantic salmon (Verhille et al., 2013; Fraser et al., 2013, 2015). The heart is a highly plastic organ, so the contrasting observations could be due to environmental conditions such as rearing temperature, salinity and natural versus controlled environmental challenges (Fraser et al., 2015).

Values of CT_{max} increased by approximately 0.4°C with every 1°C increase in acclimation temperature in both triploids and diploids, which is consistent with the changes in CT_{max} observed with acclimation temperature in previous studies of salmonids (Lutterschmidt and Hutchison, 1997; Currie et al., 1998). Nevertheless, aerobic scope remained stable across acclimation temperatures in both ploidies, suggesting that increased aerobic capacity was not responsible for the increase in CT_{max} of the fish acclimated to the higher temperatures (Fig. 2C). This contrasts with the idea of oxygen- and capacity-limited thermal tolerance (OCLTT), which assumes that temperature-dependent performance and thermal tolerance are governed by aerobic scope (Pörtner and Knust, 2007; Pörtner and Farrell, 2008). The lack of linkage between aerobic scope and CT_{max} adds to a growing database indicating that thermally dependent fitness is primarily driven by factors other than oxygen supply capacity (Clark et al., 2013; Lefevre, 2016; Sandblom et al., 2016).

Conclusions and future directions

We have shown that the thermal tolerance of juvenile Atlantic salmon in freshwater is similar in diploids and triploids, and it does not appear to be influenced by aerobic capacity. We have also shown that differential growth rates between ploidies can emerge when ploidies are mixed within the same tanks. Based on this and previous studies, we recommend that future efforts to understand the reported differences in thermal tolerance between ploidies should utilise large size ranges, freshwater versus saltwater and mixed-ploidy versus single-ploidy tank arrangements. Moreover, investigations should examine different physiological attributes in concert with the oxygen transport cascade. For example, Sambraus et al. (2017) found lower ion (Cl⁻, Na⁺ and K⁺) concentrations and higher glucose levels in blood plasma of seawater-acclimated triploid compared with diploid post-smolt Atlantic salmon at warm temperatures, suggesting lower physiological tolerance in triploids. Therefore, osmotic challenges (e.g. parr-smolt transformation, spawning migration of adults) could be investigated in conjunction with thermal stress to illuminate any role of ion regulation in determining differential ploidy survival at high temperatures. Such multi-stressor occurrences also exist during different phases of salmon aquaculture (e.g. freshwater influx in coastal aquaculture facilities), where triploids are increasingly favoured because of the advantages gained by their inherent sterility (Benfey, 1999, 2001). The findings and suggestions we have highlighted here should pave the way for future studies to further ascertain the extent to which triploids represent a useful model for elucidating the mechanisms underlying environmental tolerance in fish.

Acknowledgements

The authors would like to thank staff at the Salmon Enterprises of Tasmania Pty Ltd (SALTAS) for assistance with sourcing and transporting fish.

Competing interests

The authors declare no competing or financial interests.

Author contributions

Conceptualization: A.J.B., S.A., T.C.; Methodology: A.J.B., S.A., T.C.; Validation: A.J.B.; Formal analysis: A.J.B.; Investigation: A.J.B.; Data curation: A.J.B., T.C.;

Writing - original draft: A.J.B.; Writing - review & editing: A.J.B., S.A., N.E., P.F., T.C.; Visualization: S.A., N.E., P.F., T.C.; Supervision: S.A., N.E., P.F., T.C.; Funding acquisition: N.E., P.F.

Funding

Funding for this study was provided by the Commonwealth Scientific and Industrial Research Organisation (CSIRO) and the University of Tasmania. Support for A.J.B. was also provided by the Australian Research Training Program.

References

- Altimiras, J., Axelsoon, M., Claireaux, G., Lefrancois, C., Mercier, C. and Farrell, A. P. (2002). Cardiorespiratory status of triploid brown trout during swimming at two acclimation temperatures. *J. Fish. Biol.* **60**, 102-116.
- Atkins, M. E. and Benfey, T. J. (2008). Effect of acclimation temperature on routine metabolic rate in triploid salmonids. *Comp. Biochem. Physiol. A* **149**, 157-161.
- Barnes, R. (Katersky), King, H. and Carter, C. G. (2011). Hypoxia tolerance and oxygen regulation in Atlantic salmon, *Salmo salar*, from a Tasmanian population. *Aquaculture* **318**, 397-401.
- Benfey, T. J. (1999). The physiology and behavior of triploid fishes. *Rev. Fish. Sci.* **7**, 39-67.
- Benfey, T. (2001). Use of sterile triploid Atlantic salmon (*Salmo salar* L.) for aquaculture in New Brunswick, Canada. *ICES J. Mar. Sci.* **58**, 525-529.
- Benfey, T. J. and Sutterlin, A. M. (1984a). The haematology of triploid landlocked Atlantic salmon, *Salmo solar* L. *J. Fish. Biol.* **24**, 333-338.
- Benfey, T. J. and Sutterlin, A. M. (1984b). Oxygen utilization by triploid landlocked Atlantic salmon (*Salmo salar* L.). *Aquaculture* **42**, 69-73.
- Benfey, T. J., Sutterlin, A. M. and Thompson, R. J. (1984). Use of erythrocyte measurements to identify triploid salmonids. *Can. J. Fish. Aquat. Sci.* **41**, 980-984.
- Benfey, T. J., McCabe, L. E., Pepin, P., Benfey, T., McCabe, L. and Pepin, P. (1997). Critical thermal maxima of diploid and triploid brook charr, *Salvelinus fontinalis*. *Environ. Biol. Fish.* **49**, 259-264.
- Bœuf, G. and Payan, P. (2001). How should salinity influence fish growth? *Comp. Biochem. Phys. C* **130**, 411-423.
- Brett, J. R. (1964). The respiratory metabolism and swimming performance of young sockeye salmon. *J. Fish. Res. Bd. Can.* **21**, 1183-1226.
- Brijs, J., Jutfelt, F., Clark, T. D., Gräns, A., Ekström, A. and Sandblom, E. (2015). Experimental manipulations of tissue oxygen supply do not affect warming tolerance of European perch. *J. Exp. Biol.* **218**, 2448-2454.
- Carter, C. G., McCarthy, I. D., Houlihan, D. F., Johnstone, R., Walsingham, M. V. and Mitchell, A. I. (1994). Food consumption, feeding behaviour, and growth of triploid and diploid Atlantic salmon, *Salmo salar* L., parr. *Can. J. Zool.* **72**, 609-617.
- Clark, T. D., Sandblom, E., Cox, G. K., Hinch, S. G. and Farrell, A. P. (2008). Circulatory limits to oxygen supply during an acute temperature increase in the Chinook salmon (*Oncorhynchus tshawytscha*). *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **295**, R1631-R1639.
- Clark, T. D., Donaldson, M. R., Pieperhoff, S., Drenner, S. M., Lotto, A., Cooke, S. J., Hinch, S. G., Patterson, D. A. and Farrell, A. P. (2012). Physiological benefits of being small in a changing world: responses of Coho salmon (*Oncorhynchus kisutch*) to an acute thermal challenge and a simulated capture event. *PLoS ONE* **7**, e39079.
- Clark, T. D., Sandblom, E. and Jutfelt, F. (2013). Aerobic scope measurements of fishes in an era of climate change: respirometry, relevance and recommendations. *J. Exp. Biol.* **216**, 2771-2782.
- Clark, T. D., Roche, D. G., Binning, S. A., Speers-Roesch, B. and Sundin, J. (2017). Maximum thermal limits of coral reef damselfishes are size-dependent and resilient to near-future ocean acidification. *J. Exp. Biol.* **220**, jeb162529.
- Clarke, A. and Johnston, N. M. (1999). Scaling of metabolic rate with body mass and temperature in teleost fish. *J. Anim. Ecol.* **68**, 893-905.
- Currie, R. J., Bennett, W. A. and Beiting, T. L. (1998). Critical thermal minima and maxima of three freshwater game-fish species acclimated to constant temperatures. *Environ. Biol. Fish.* **51**, 187-200.
- Cutts, C. J., Betcalfe, N. B. and Caylor, A. C. (1998). Aggression and growth depression in juvenile Atlantic salmon: the consequences of individual variation in standard metabolic rate. *J. Fish. Biol.* **52**, 1026-1037.
- Dejours, P. (1981). Principles of comparative respiratory physiology.
- Ekström, A., Brijs, J., Clark, T. D., Gräns, A., Jutfelt, F. and Sandblom, E. (2016). Cardiac oxygen limitation during an acute thermal challenge in the European perch: effects of chronic environmental warming and experimental hyperoxia. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **311**, R440-R449.
- Ern, R., Norin, T., Gampel, A. K. and Esbaugh, A. J. (2016). Oxygen dependence of upper thermal limits in fishes. *J. Exp. Biol.* **219**, 3376-3383.
- Farrell, A. P. (2009). Environment, antecedents and climate change: lessons from the study of temperature physiology and river migration of salmonids. *J. Exp. Biol.* **212**, 3771-3780.
- Fox, J. and Weisberg, S. (2011). *An R Companion to Applied Regression*. 2nd edn. Thousand Oaks, California: Sage.
- Fraser, T. W. K., Hansen, T., Skjæraasen, J. E., Mayer, I., Sambras, F. and Fjellidal, P. G. (2013). The effect of triploidy on the culture performance, deformity prevalence, and heart morphology in Atlantic salmon. *Aquaculture* **416-417**, 255-264.
- Fraser, T. W. K., Mayer, I., Hansen, T., Poppe, T. T., Skjæraasen, J. E., Koppang, E. O. and Fjellidal, P. G. (2015). Vaccination and triploidy increase relative heart weight in farmed Atlantic salmon, *Salmo salar* L. *J. Fish. Dis.* **38**, 151-160.
- Fry, F. E. J. (1971). The effect of environmental factors on the physiology of fish. In *Environmental Relations and Behavior* (ed. W. S. Hoar and D. J. Randall), pp. 1-98. Amsterdam: Elsevier.
- Galbreath, P. F., Jean, W. S., Anderson, V. and Thorgaard, G. H. (1994). Freshwater performance of all-female diploid and triploid Atlantic salmon. *Aquaculture* **128**, 41-49.
- Galbreath, P. F., Adams, N. D., Sherrill, L. W. and Martin, T. H. (2006). Thermal tolerance of diploid versus triploid rainbow trout and brook trout assessed by time to chronic lethal maximum. *Environ. Biol. Fish.* **75**, 183-193.
- Gillooly, J. F., Brown, J. H., West, G. B., Savage, V. M. and Charnov, E. L. (2001). Effects of size and temperature on metabolic rate. *Science* **293**, 2248-2251.
- Graham, M. S. and Farrell, A. P. (1989). The effect of temperature acclimation and adrenaline on the performance of a perfused trout heart. *Physiol. Zool.* **62**, 38-61.
- Graham, M. S., Fletcher, G. L. and Benfey, T. J. (1985). Effect of triploidy on blood oxygen content of Atlantic salmon. *Aquaculture* **50**, 133-139.
- Iwama, G. K. (1996). Growth of salmonids. In *Principles of Salmonid Culture* (ed. W. Pennell and B. A. Barton), pp. 467-515. Amsterdam: Elsevier.
- Johnson, O. W., Dickhoff, W. W. and Utter, F. M. (1986). Comparative growth and development of diploid and triploid coho salmon, *Oncorhynchus kisutch*. *Aquaculture* **57**, 329-336.
- Killen, S. S., Norin, T. and Halsey, L. G. (2017). Do method and species lifestyle affect measures of maximum metabolic rate in fishes? *J. Fish. Biol.* **90**, 1037-1046.
- Lannig, G., Bock, C., Sartoris, F. J. and Pörtner, H. O. (2004). Oxygen limitation of thermal tolerance in cod, *Gadus morhua* L., studied by magnetic resonance imaging and on-line venous oxygen monitoring. *Am. J. Physiol. Regul. Integr. Comp. Physiol.* **287**, R902-R910.
- Lefevre, S. (2016). Are global warming and ocean acidification conspiring against marine ectotherms? A meta-analysis of the respiratory effects of elevated temperature, high CO₂ and their interaction. *Conserv. Physiol.* **4**, cow009.
- Lenth, R. V. (2016). Least-squares means: the R package lsmeans. *J. Stat. Softw.* **69**, 1-33.
- Lijalad, M. and Powell, M. D. (2009). Effects of lower jaw deformity on swimming performance and recovery from exhaustive exercise in triploid and diploid Atlantic salmon *Salmo salar* L. *Aquaculture* **290**, 145-154.
- Lutterschmidt, W. I. and Hutchison, V. H. (1997). The critical thermal maximum: history and critique. *Can. J. Zool.* **75**, 1561-1574.
- Maxime, V. (2008). The physiology of triploid fish: current knowledge and comparisons with diploid fish. *Fish Fish.* **9**, 67-78.
- McCarthy, I. D., Carter, C. G., Houlihan, D. F., Johnstone, R. and Mitchell, A. I. (1996). The performance of all-female diploid and triploid Atlantic salmon smolts on transfer together to sea water. *J. Fish. Biol.* **48**, 545-548.
- McGeachy, S. A., Benfey, T. J. and Friars, G. W. (1995). Freshwater performance of triploid Atlantic salmon (*Salmo salar*) in New Brunswick aquaculture. *Aquaculture* **137**, 333-341.
- Messmer, V., Pratchett, M. S., Hoey, A. S., Tobin, A. J., Coker, D. J., Cooke, S. J. and Clark, T. D. (2016). Global warming may disproportionately affect larger adults in a predatory coral reef fish. *Glob. Chang. Biol.* **23**, 2230-2240.
- Metcalfe, N. B., Taylor, A. C. and Thorpe, J. E. (1995). Metabolic rate, social status and life-history strategies in Atlantic salmon. *Anim. Behav.* **49**, 431-436.
- Norin, T. and Clark, T. D. (2016). Measurement and relevance of maximum metabolic rate in fishes. *J. Fish. Biol.* **88**, 122-151.
- Norin, T., Malte, H. and Clark, T. D. (2014). Aerobic scope does not predict the performance of a tropical eurythermal fish at elevated temperatures. *J. Exp. Biol.* **217**, 244-251.
- Norin, T., Malte, H. and Clark, T. D. (2016). Differential plasticity of metabolic rate phenotypes in a tropical fish facing environmental change. *Funct. Ecol.* **30**, 369-378.
- O'Flynn, F. M., McGeachy, S. A., Friars, G. W., Benfey, T. J. and Bailey, J. K. (1997). Comparisons of cultured triploid and diploid Atlantic salmon (*Salmo salar* L.). *ICES J. Mar. Sci.* **54**, 1160-1165.
- Ojolic, E. J., Cusack, R., Benfey, T. J. and Kerr, S. R. (1995). Survival and growth of all-female diploid and triploid rainbow trout (*Oncorhynchus mykiss*) reared at chronic high temperature. *Aquaculture* **131**, 177-187.
- Oliva-Teles, A. and Kaushik, S. J. (1990). Growth and nutrient utilization by 0+ and 1+ triploid rainbow trout, *Oncorhynchus mykiss*. *J. Fish. Biol.* **37**, 125-133.
- Ozolina, K., Shiels, H. A., Ollivier, H. and Claireaux, G. (2016). Intraspecific individual variation of temperature tolerance associated with oxygen demand in the European sea bass (*Dicentrarchus labrax*). *Conserv. Physiol.* **4**, cov060.
- Pedersen, B. H. (1997). The cost of growth in young fish larvae, a review of new hypotheses. *Aquaculture* **155**, 259-269.
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. and R Core Team (2018). nlme: Linear and Nonlinear Mixed Effects Models. R package version 3.1-131.1. <https://CRAN.R-project.org/package=nlme>

- Pörtner, H. O. and Farrell, A. P.** (2008). Ecology. Physiology and climate change. *Science* **322**, 690-692.
- Pörtner, H. O. and Knust, R.** (2007). Climate change affects marine fishes through the oxygen limitation of thermal tolerance. *Science* **315**, 95-97.
- Sadler, J., Wells, R. M. G., Pankhurst, P. M. and Pankhurst, N. W.** (2000). Blood oxygen transport, rheology and haematological responses to confinement stress in diploid and triploid Atlantic salmon, *Salmo salar*. *Aquaculture* **184**, 349-361.
- Sambraus, F., Olsen, R. E., Remen, M., Hansen, T. J., Torgersen, T. and Fjellidal, P. G.** (2017). Water temperature and oxygen: The effect of triploidy on performance and metabolism in farmed Atlantic salmon (*Salmo salar* L.) post-smolts. *Aquaculture* **473**, 1-12.
- Sandblom, E., Clark, T. D., Gräns, A., Ekström, A., Brijs, J., Sundström, L. F., Odelström, A., Adill, A., Aho, T. and Jutfelt, F.** (2016). Physiological constraints to climate warming in fish follow principles of plastic floors and concrete ceilings. *Nat. Commun.* **7**, 11447.
- Scott, M. A.** (2012). Performance of wild and domestic strains of diploid and triploid rainbow trout (*Oncorhynchus mykiss*) in response to environmental challenges. *MSc thesis*, University of British Columbia, Vancouver, BC, Canada.
- Scott, M. A., Dhillon, R. S., Schulte, P. M., Richards, J. G. and Magnan, P.** (2015). Physiology and performance of wild and domestic strains of diploid and triploid rainbow trout (*Oncorhynchus mykiss*) in response to environmental challenges. *Can. J. Fish. Aquat. Sci.* **72**, 125-134.
- Small, S. A. and Benfey, T. J.** (1987). Cell size in triploid salmon. *J. Exp. Zool.* **241**, 339-342.
- Swarup, H.** (1959). Effect of triploidy on the body size, general organization and cellular structure in *Gasterosteus aculeatus* (L.). *J. Genet.* **56**, 143-155.
- Teskeredžić, E., Donaldson, E. M., Teskeredžić, Z., Solar, I. I. and McLean, E.** (1993). Comparison of hydrostatic pressure and thermal shocks to induce triploidy in coho salmon (*Oncorhynchus kisutch*). *Aquaculture* **117**, 47-55.
- Van Leeuwen, T. E., Rosenfeld, J. S. and Richards, J. G.** (2012). Effects of food ration on SMR: influence of food consumption on individual variation in metabolic rate in juvenile coho salmon (*Oncorhynchus kisutch*). *J. Anim. Ecol.* **81**, 395-402.
- Verhille, C., Anttila, K. and Farrell, A. P.** (2013). A heart to heart on temperature: impaired temperature tolerance of triploid rainbow trout (*Oncorhynchus mykiss*) due to early onset of cardiac arrhythmia. *Comp. Biochem. Physiol. A* **164**, 653-657.
- Yamamoto, T., Ueda, H. and Higashi, S.** (1998). Correlation among dominance status, metabolic rate and otolith size in masu salmon. *J. Fish. Biol.* **52**, 281-290.