

**Title:** Dietary fibre intake and its association with inflammatory markers in adolescents

**Authors:**

Olivia G. Swann<sup>1</sup>, Monique Breslin<sup>1</sup>, Michelle Kilpatrick<sup>1</sup>, Therese A. O'Sullivan<sup>2</sup>, Trevor A. Mori<sup>3</sup>, Lawrence J. Beilin<sup>3</sup>, Wendy H. Oddy<sup>1</sup>

**Institutions:**

1 - Menzies Institute for Medical Research, University of Tasmania, Hobart, Tasmania, Australia

2 - School of Medical and Health Science, Edith Cowan University, Joondalup, Western Australia, Australia

3 - Medical School, Royal Perth Hospital Unit, University of Western Australia, Perth, Western Australia, Australia

**Corresponding author:**

Professor Wendy H. Oddy

Menzies Institute for Medical Research, University of Tasmania

Private Bag 23, Hobart TAS 7000,

[Wendy.Oddy@utas.edu.au](mailto:Wendy.Oddy@utas.edu.au)

+61 3 6226 7514



This peer-reviewed article has been accepted for publication but not yet copyedited or typeset, and so may be subject to change during the production process. The article is considered published and may be cited using its DOI

10.1017/S0007114520001609

The British Journal of Nutrition is published by Cambridge University Press on behalf of The Nutrition Society

**Running head:** Dietary fibre and inflammation in adolescents

**Keywords:** dietary fibre, dietary fiber, inflammation, C-reactive protein, adolescence, Raine Study

**Abstract**

A high dietary fibre intake has been associated with improvements in inflammatory conditions in adults. However, little is known on whether associations between dietary fibre and inflammation are evident during adolescence. We examined the relationship between dietary fibre intake measured by food frequency questionnaire and the inflammatory marker high-sensitivity C-reactive protein (hs-CRP) and the adipokines leptin and adiponectin cross-sectionally in 17-year-olds participating in the Raine Study (n=621). In weighted analysis using tobit and linear regression, and after excluding participants with hs-CRP>10mg/L, higher total dietary fibre intake (per 5g/day) was significantly associated with lower leptin ( $\beta=-0.13$ , 95% CI -0.17, -0.09) and adiponectin ( $\beta=-0.28$ , 95% CI -0.49, -0.07), but not hs-CRP, in unadjusted analyses. These associations were no longer significant after adjustment for gender, anthropometry and a number of lifestyle factors. However, higher cereal and grain fibre intake was significantly associated with lower leptin ( $\beta=-0.06$ , 95% CI -0.10, -0.01) in fully adjusted analysis. Our findings suggest that a higher intake of cereal and grain fibre may contribute to lower leptin in adolescents. This may contribute to reductions in low-grade chronic inflammation and improved health outcomes.

**Acronyms:**

hs-CRP: high sensitivity C Reactive Protein

CSIRO: Commonwealth Scientific and Industrial Research Organisation

FFQ: Food Frequency Questionnaire

NUTTAB: Nutrient Tables for use in Australia

BMI: Body Mass Index

WHtR: Waist Height Ratio

## ***Introduction***

Inflammation is an intrinsic part of the immune system, playing a role in the prevention of infection<sup>(1)</sup>. However, prolonged inflammation in the absence of an acute inflammatory stimulus, referred to as chronic inflammation, has been implicated in the progression of conditions such as cardiovascular disease, type 2 diabetes, non-alcoholic fatty liver disease, metabolic syndrome, and depression<sup>(2-4)</sup>. While mechanisms by which dietary fibre may decrease disease risk are not fully understood, a higher fibre intake has been associated with lower risk of cardiovascular disease, type 2 diabetes, and metabolic syndrome<sup>(5-7)</sup>. Reducing or preventing chronic inflammation is a plausible mechanism, supported by observed associations between higher dietary fibre intake and lower inflammatory markers<sup>(8, 9)</sup>.

Some non-digestible dietary fibres are fermented in the intestine by the gut microbiota, creating by-products with the ability to inhibit and suppress inflammation and oxidative stress<sup>(10, 11)</sup>. Potential mechanisms include reducing the permeability of the intestinal membrane<sup>(12-14)</sup>, lowering release of lipopolysaccharide from the gut and consequent production of inflammatory cytokines such as interleukin (IL) 6<sup>(15)</sup>. In turn, IL-6 stimulates the production of C-reactive protein (CRP), an acute-phase inflammatory reactant frequently used to measure chronic inflammation<sup>(16, 17)</sup>.

A higher dietary fibre intake may alternatively impact inflammation through a reduction in adipose tissue<sup>(18)</sup>, as excess adiposity has been associated with elevated inflammation<sup>(19-21)</sup>. Leptin is a pro-inflammatory adipokine and immune modulator produced in adipose tissue and associated with increased inflammation<sup>(22, 23)</sup>. In contrast, adiponectin is anti-inflammatory and high concentrations have been associated with reduced inflammation in the metabolic syndrome and other related conditions<sup>(22, 24)</sup>. Although associations between dietary fibre and inflammation have been studied in adult populations, only one study has previously investigated the concept in healthy adolescents, to our knowledge. The study of American adolescents found that a higher dietary fibre intake was linked to greater levels of adiponectin and lower levels of leptin<sup>(9)</sup>. Higher dietary fibre intake was also associated with lower leptin concentrations in adults and female young adults<sup>(25, 26)</sup>.

Inadequate dietary fibre intakes are not exclusive to adults, with adolescent fibre intakes being below recommendations in much of the western world<sup>(27, 28)</sup>. More research to determine the effect of low dietary fibre intake on biomarkers of health during adolescence is warranted as adolescent health may have consequences in adulthood<sup>(29, 30)</sup>. Additionally, the

impact of dietary fibre sourced from different food groups (e.g. cereals and grains, fruits, and vegetables) on CRP, leptin, and adiponectin concentrations has been insufficiently studied in healthy populations. The analysis of different sources of dietary fibre is supported by two studies of adults with diabetes which found a significant association between higher cereal fibre intake and lower CRP in women<sup>(31)</sup> and with higher cereal fibre intake and higher adiponectin in men<sup>(32)</sup>. Another study of women found associations between higher total fibre and cereal fibre intake and higher adiponectin, and between higher cereal fibre and lower hs-CRP<sup>(33)</sup>. We aimed to investigate the association between dietary fibre intake, both overall and from different sources, and hs-CRP, leptin and adiponectin in adolescents participating in the 17-year follow-up of the Raine Study.

## ***Methods***

### *Participants*

The Raine Study is a prospective study which recruited 2,900 pregnant mothers (generation 1) through the major maternity hospital in Perth, Western Australia between 1989 and 1991, resulting in 2,868 live births (generation 2). Multiple follow-ups have been performed from birth through to adulthood and cross-sectional data from the 17-year follow-up is used in this study based on availability of inflammation variables. Detailed cohort information and characteristics have been published elsewhere<sup>(34, 35)</sup>. Briefly, the 17-year cohort included 1,621 adolescents, with 748 eligible for this study after exclusion of those without complete data for diet and inflammatory markers. Exclusion of those without anthropomorphic and lifestyle data resulted in 621 adolescents with all data (Figure 1).

### *Dietary variables*

Dietary fibre and energy intake data were obtained from the 212-item Commonwealth Scientific and Industrial Research Organisation (CSIRO) semi-quantitative food frequency questionnaire (FFQ), completed by the adolescent. The FFQ measures usual diet over the previous year with a series of questions about consumption frequency and serving size of various foods, including mixed dishes and beverages. Estimates of daily intake of dietary fibre and energy were provided by the CSIRO, calculated from serving size and nutrient content of each food item. The CSIRO FFQ has been validated for use in the Australian population<sup>(36)</sup> and within the Raine Study cohort<sup>(37)</sup>. Dietary fibre, sourced from cereals and

grains, fruits and vegetables, and nuts and legumes was determined by assigning a fibre concentration (%) to each food item (Supplementary table 1, based on the Nutrient Tables for use in Australia (NUTTAB) dietary composition database) and combining it with each adolescent's intake of those foods (g/day) to obtain a daily fibre intake from each food (g/day). These intakes from individual foods were combined to give a total fibre intake per food group. Dietary fibre intake from nuts and legumes was considered but not analysed separately due to extremely low intakes (mean intake of 0.60g/day). Dietary misreporting was calculated from energy intake and basal metabolic rate using the Goldberg method to classify participants as under-, over-, or plausible reporters<sup>(38)</sup> as described elsewhere<sup>(39)</sup>.

### *Sociodemographic and lifestyle variables*

Information on lifestyle factors including screen time, leisure-time physical activity, smoking status, alcohol intake, and oral contraceptive use in females was obtained from self-report questionnaires completed by the adolescent at the 17-year follow-up. Screen time reported the average number of hours spent either using a computer or watching television (whichever was highest) on a weekend day. Physical activity refers to hours per week spent in vigorous physical activity outside of school or work. Smoking was categorised as the average number of cigarettes smoked per day in the previous week. Alcohol intake was measured in standard drinks per week, calculated from reports of the number and variety of alcoholic beverages consumed in the previous week. Females were categorised into those currently taking oral contraceptives or not. Inflammation linked medications considered in the study included antihistamines, oral steroids, immunosuppressants, anti-cholinergics, and non-steroidal anti-inflammatory drugs and was recorded as yes or no. Adolescents who were taking these medications (n=5) were excluded from analysis.

Parental ethnicity was recorded during pregnancy and used to categorise the adolescents as Caucasian (both parents Caucasian), mixed (one parent Caucasian and one parent non-Caucasian) or non-Caucasian (both parents non-Caucasian). Height (to the nearest 0.1cm using a Holtain Stadiometer), weight (to the nearest 100g using a Wedderburn Digital Chair Scale), and waist circumference (to the nearest 0.1cm) were measured by trained researchers. Body mass index (BMI) was calculated from weight (kg) divided by height squared (m<sup>2</sup>). Waist height ratio (WHtR), a measure of central adiposity, was calculated by dividing waist circumference (cm) by height (cm). WHtR and BMI were both chosen as they measure different aspects of adiposity (general adiposity for BMI and central adiposity for

WHtR) and are independently associated with fibre intake and inflammation (data not shown). Adolescents with a WHtR of 0.5 or higher were considered to have central obesity<sup>(40)</sup>.

### *Inflammatory markers*

Fasting blood samples (no food or beverage other than water consumed in the previous 12 hours) were taken by a phlebotomist in the home of the participant. Blood samples were analysed at the PathWest Laboratory at the Royal Perth Hospital. hs-CRP was measured on an Architect c16000 Analyser with an intra-assay coefficient of variation of 15.96% and a lower detection threshold of 0.14mg/L. Leptin was measured by the ACTIVE® Human Leptin ELISA kit (DSL-10-23100, Diagnostic Systems Laboratories, Webster, TX, USA) with a lower detection threshold of 1.5µg/L. Adiponectin was measured by Quantikine® Human Total Adiponectin/Acrp30 Immunoassay (R&D Systems, Minneapolis, MN, USA). Based on the recommendations of the American Heart Association, hs-CRP values  $\geq 10$  mg/L were excluded as they are considered to represent acute inflammation due to injury or sickness, rather than chronic inflammation<sup>(41)</sup>.

### *Ethics*

This study was conducted according to the guidelines laid down in the Declaration of Helsinki. Ethics approval for the Raine study was granted by the committees of King Edward Memorial Hospital for Women and Princess Margaret Hospital for Children, Perth, Western Australia. Informed written consent was obtained from the adolescent and their primary caregiver.

### *Analytical methods*

After removing participants who were taking inflammation-linked medications or had hs-CRP  $>10$ mg/L, 602 participants were included in the analysis. Associations between dietary fibre intake (per 5g/day increases) and hs-CRP and leptin were estimated using Tobit regression. Tobit regression was developed for analysis of datasets where insufficient range in the measurement scale results in a large number of observations clustering at one end of the scale<sup>(42)</sup>. In this case, the lower detection threshold of the leptin and hs-CRP measurement assays led to partial information loss (censoring) where the true levels below that threshold were unknown. Tobit regression uses a modified maximum likelihood estimation method

which considers whether the data is censored when calculating estimates to give an unbiased outcome<sup>(42)</sup>. Both hs-CRP and leptin were log transformed due to skewed distribution. Associations between dietary fibre intake (per 5g/day increases) and adiponectin concentrations were estimated with linear regression. Confounders were chosen based on biological plausibility and a significant ( $p < 0.1$ ) association with hs-CRP, leptin, or adiponectin, determined by univariable Tobit and linear regression (Supplementary table 2). Inverse probability weighting was applied to analysis due to the large numbers of participants who did not have the required data. Weighting variables were measured for the full population at birth and consisted of maternal age at birth, family income, and sex of the participant. There were 3 multivariable models tested in addition to the unadjusted analysis (model 1). Model 2 included sex only, while model 3 further adjusted for WHtR, BMI, energy intake, and dietary misreporting. Model 4 adjusted for the factors included in model 3 with the addition of ethnicity and lifestyle factors (screen time, physical activity, smoking, and alcohol intake). Sensitivity analysis including only participants categorised as plausible reporters was performed with results consistent with the main analysis (Supplementary table 3). Sex stratified analysis was also performed due to a significant interaction between dietary fibre intake and sex. In sex stratified analysis, models 1, 3, and 4 were used (minus the sex variable), and oral contraceptive usage was added to model 4 in females only. Statistical analysis was performed with R version 3.5.0<sup>(43)</sup> and R studio version 1.2.1335<sup>(44)</sup>.

## **Results**

Table 1 shows the characteristics of adolescents included in the analysis and those excluded due to incomplete data. Proportions in the excluded adolescents were calculated for those without missing data for the relevant variable (valid %). The excluded population had a higher proportion of males (51%) than the included population (47%). Excluded adolescents were less active, smoked more, and drank more alcohol than those with complete data. A higher proportion of excluded participants had elevated hs-CRP ( $>3\text{mg/L}$ ) compared to those included in the study (16% and 11% respectively). Of the female adolescents included, 34% were taking oral contraceptives compared to 25% in those who were excluded (Table 1). Excluded adolescents also had higher hs-CRP and leptin concentrations (Table 2).

Mean concentrations of hs-CRP were significantly higher in females (1.33mg/L (SD 1.75mg/L)) than in males (0.92mg/L (SD 1.43mg/L),  $p=0.002$ ). Concentrations of leptin differed substantially, with a mean concentration of 27.77 $\mu\text{g/L}$  (SD 19.07 $\mu\text{g/L}$ ) in females

and 5.40 $\mu$ g/L (SD 7.37 $\mu$ g/L) in males ( $p < 0.001$ ). Likewise, adiponectin levels were higher in females (11.88mg/L (SD 5.19mg/L)) than in males (7.98mg/L (SD 3.86mg/L),  $p < 0.001$ ). Conversely, daily dietary fibre intake was significantly lower in females (20.7g/day (SD 7.5g/day)) than in males (24.1g/day (SD 9.8g/day),  $p < 0.001$ ) (Figure 2).

In unadjusted analysis (model 1), a 5g/day higher dietary fibre intake was significantly associated with lower levels of leptin and adiponectin (Table 3). The association between dietary fibre intake and lower leptin remained after adjustment for sex (model 2) but was no longer significant after further adjustment (models 3 and 4). In analyses that examined males and females separately there was a significant association between higher dietary fibre intake and lower leptin in males (Supplementary table 4) and between dietary fibre intake and lower hs-CRP in females (Supplementary table 5), but neither persisted after adjustment for confounders.

When dietary fibre from different food groups were examined separately, a 5g/day higher cereal and grain fibre intake was significantly associated with lower hs-CRP, leptin, and adiponectin in unadjusted analysis (model 1). The association of cereal and grain fibre with leptin remained significant after adjusting for all confounders (model 4), although the effect was small (Table 3). Similar results were seen in males, with a significant association between higher cereal and grain fibre and lower leptin in all models (Supplementary table 4). There were no associations between cereal and grain fibre and hs-CRP, leptin, or adiponectin in females (Supplementary table 5). Likewise, fibre sourced from fruits and vegetables was not associated with hs-CRP, leptin, or adiponectin overall (Table 3) or in females (Supplementary table 5). In males, fruit and vegetable fibre was associated with lower adiponectin in unadjusted analysis only (Supplementary table 4).

## ***Discussion***

In our population cohort of adolescents at 17 years of age, there were no associations between a higher dietary fibre intake and hs-CRP, leptin or adiponectin after adjustment for gender, anthropometry and a number of lifestyle factors. However, a 5g/day higher fibre intake from cereals and grains had a small but significant association with lower levels of leptin in fully adjusted analysis. In sex stratified analysis, the same association between cereal and grain fibre and leptin was seen in males but not females. There were no associations between hs-CRP, leptin or adiponectin with fruit and vegetable fibre intake.

The association between cereal and grain fibre and lower leptin concentrations concurs with existing research showing lower levels of leptin in mice fed cereal fibres<sup>(45)</sup>. A potential explanation for this association is that a high wholegrain intake has been linked to lower overweight and obesity<sup>(46)</sup> which may in turn reduce leptin<sup>(47)</sup>. In our study, however, the observed association of cereal and grain fibre with lower leptin persisted after adjustment for WHtR and BMI. Similarly, no significant changes in weight or BMI were seen in a study of overweight and obese children where an intervention consisting of increasing wholegrain intake resulted in lower leptin concentrations compared to controls who refrained from consuming wholegrains<sup>(48)</sup>. However, it may be that a more direct measure of fat mass such as body composition measured by dual-energy X-ray absorptiometry (DXA) is required to determine the impact of adiposity, as dietary fibre may impact visceral adipose tissue independent of subcutaneous adipose tissue<sup>(49)</sup>. BMI and WHtR may not accurately represent the effect of different types of adipose tissue on inflammation. Beyond overweight and obesity, nutrients such as polyphenols which are found in foods high in cereal and grain fibre may also play a role in the association between cereal and grain fibre and leptin<sup>(50)</sup>.

The importance of leptin to health is highlighted by the regulatory role it plays in many processes in the body including the immune system<sup>(51)</sup>. Leptin can stimulate immune cells and has been shown to upregulate production of inflammatory compounds such as tumour necrosis factor alpha (TNF- $\alpha$ )<sup>(52)</sup>. Therefore, through lowering leptin levels a higher cereal and grain fibre intake may help to reduce chronic inflammation and improve health.

In our study we showed that at age 17, males had higher intakes of dietary fibre than females, and that females had higher concentrations of hs-CRP, leptin, and adiponectin than males, consistent with previous work in this cohort at 14<sup>(53)</sup> and 17 years<sup>(54, 55)</sup>. Similar higher levels of hs-CRP in women than men were also observed in a national study of young American adults<sup>(56)</sup>. Concentrations of hs-CRP, leptin, and adiponectin are well known to be linked with adiposity<sup>(57, 58)</sup>. It is speculated that the sex differences in leptin and adiponectin may be due to higher body fat percentages in females<sup>(59)</sup> or through differences in sex hormones<sup>(60)</sup>. The association between male sex and lower hs-CRP, leptin, and adiponectin remained significant after adjustment for BMI and WHtR (data not shown).

The lack of an association between total dietary fibre intake and hs-CRP in our study is in contrast to previous studies in healthy adult populations<sup>(8, 61-65)</sup> and in adolescents<sup>(9)</sup>. However, it is not without precedent, with studies in female young adult and female postmenopausal populations finding no association between dietary fibre intake and hs-CRP<sup>(66, 67)</sup>. A possible explanation for the discrepancy in results relates to how dietary fibre

intake was determined. Most studies have used a 24-hour dietary recall method to measure diet, in contrast to the FFQ used in our study. Studies using FFQs in healthy populations are scarce, and only one known study reported a significant association between hs-CRP and dietary fibre intake<sup>(62)</sup>. FFQs may not accurately represent diet at the time that inflammatory markers were measured, and therefore may not capture the effects of short term dietary changes on hs-CRP concentrations. Additionally, studies on the impact of dietary fibre intake on hs-CRP are primarily conducted in adult populations. There is insufficient research in adolescents to determine whether age may play a role in the relationship between dietary fibre and inflammation. Adolescents in our study had lower concentrations of hs-CRP than are reported in most studies of adults<sup>(8, 61, 63-65)</sup>. Although an association between higher dietary fibre intake and lower hs-CRP was found in another adolescent population with similar concentrations of hs-CRP, that study used a 24hr dietary recall method to measure dietary fibre intake<sup>(9)</sup>.

### *Strengths and limitations*

Strengths of our study include the use of a large population-based pregnancy cohort with comprehensive participant data, allowing for testing of a wide range of confounders. Our dietary data were taken from the CSIRO FFQ, which includes questions on a wide range of foods, and reflects usual intake over the past 12 months. However, a limitation of the FFQ is that it can be affected by recall bias and does not reflect transient changes in diet which may impact inflammation in the short term. One of the major proposed mechanisms for an effect of dietary fibre on inflammation is through the gut microbiota<sup>(10)</sup>. The composition of the microbiota responds rapidly to changes in diet, with differences in diversity seen within days of dietary intervention<sup>(68)</sup>. Therefore, dietary fibre intake as measured over the previous year with an FFQ may not reflect the current state of the microbiota and its impact on hs-CRP concentrations.

Another limitation is that due to our cross-sectional study design we could not determine causality. However, we judged cross-sectional analysis to be more appropriate than longitudinal analysis to examine the impact of dietary fibre on inflammation as current diet is a more likely modifier of inflammation than diet from several years earlier<sup>(68)</sup>. Additionally, data on leptin and adiponectin concentrations were not available at the previous adolescent follow-up (year 14). Other limitations lie in the availability of adiposity variables; we were

unable to adjust for fat mass, which is a more direct measure of overweight and obesity and has been correlated with leptin concentrations<sup>(69)</sup>.

The study may have been subject to exclusion bias. The number of 17-year participants with complete data was relatively low (621 of 1621 people), with those excluded having less healthy lifestyles and higher levels of hs-CRP and leptin. The exclusion of adolescents with higher inflammation may contribute to the relatively low levels of hs-CRP in our population, potentially impacting the strength of the study to identify associations between dietary fibre intake and inflammatory markers. To address this potential exclusion bias we applied weighting to the analysis of associations.

### *Conclusion*

An understanding of the effect of low dietary fibre intake on inflammatory markers during adolescence is of benefit in designing dietary interventions to prevent inflammatory diseases in adulthood. Our study does not provide evidence of an independent effect of dietary fibre intake on plasma hs-CRP, leptin, and adiponectin in adolescents. However, our results suggest that a higher intake of cereal and grain fibre specifically may contribute to lower leptin concentrations. This may improve health by reducing production of inflammatory cytokines thereby reducing inflammation. Randomised controlled trials in adolescent age groups are required to expand on and confirm our findings.

**Table 1:** Characteristics of those included and excluded from the study

|                                 | Included |    | Excluded |         |
|---------------------------------|----------|----|----------|---------|
|                                 | n        | %  | n        | Valid % |
| n                               | 621      |    | 1000     |         |
| Sex                             |          |    |          |         |
| Female                          | 331      | 53 | 489      | 49      |
| Male                            | 290      | 47 | 511      | 51      |
| Race                            |          |    |          |         |
| Caucasian                       | 525      | 85 | 842      | 87      |
| Mixed                           | 58       | 9  | 88       | 9       |
| Non-Caucasian*                  | 38       | 6  | 38       | 4       |
| Screening time (hr/weekend day) |          |    |          |         |
| <2                              | 231      | 37 | 236      | 36      |
| 2-4                             | 217      | 35 | 219      | 34      |
| >4                              | 173      | 28 | 195      | 30      |
| Physical activity (hr/week)     |          |    |          |         |
| ≤1                              | 247      | 40 | 290      | 45      |
| 2-3                             | 163      | 26 | 178      | 27      |
| 4-6                             | 131      | 21 | 94       | 15      |
| >7                              | 80       | 13 | 86       | 13      |
| Smoking (cigarettes/day)        |          |    |          |         |
| <1                              | 553      | 89 | 463      | 79      |
| 1-5                             | 38       | 6  | 40       | 7       |
| >5                              | 30       | 5  | 84       | 14      |
| Alcohol (standard drinks/week)  |          |    |          |         |
| 0                               | 331      | 53 | 268      | 44      |
| 0.1-5                           | 105      | 17 | 76       | 12      |
| 5.1-10                          | 79       | 13 | 105      | 17      |
| >10                             | 106      | 17 | 160      | 26      |
| Anti-inflammatory medications   |          |    |          |         |
| No                              | 616      | 99 | 770      | 99      |
| Yes                             | 5        | 1  | 6        | 1       |
| Contraceptives (female only)    |          |    |          |         |
| No                              | 220      | 66 | 295      | 75      |
| Yes                             | 111      | 34 | 98       | 25      |
| Elevated hs-CRP (3-10mg/L)      |          |    |          |         |
| No                              | 543      | 89 | 526      | 84      |
| Yes                             | 67       | 11 | 103      | 16      |
| WHtR                            |          |    |          |         |
| Non-obese (<0.5)                | 511      | 82 | 453      | 78      |
| Obese (>0.5)                    | 110      | 18 | 130      | 22      |

Hs-CRP, high sensitivity C-reactive protein; WHtR, waist height ratio

\*Non-Caucasian ethnicity includes Aboriginal, Asian, and Polynesian

Valid % is the proportion calculated excluding those with missing data for the relevant variable.

Percentages were rounded to the nearest 1%

**Table 2:** Diet variables, physical measurements, and markers of inflammation in those included and excluded from the study

|                                   | Included (n=621) |       |               | Excluded (n=1000) |       |               |
|-----------------------------------|------------------|-------|---------------|-------------------|-------|---------------|
|                                   | Mean             | SD    | Range         | Mean              | SD    | Range         |
| Fibre (g/day)                     | 22.3             | 8.9   | 5.9 - 68.3    | 23.1              | 10.4  | 4.2 - 65.7    |
| Cereal and grain fibre (g/day)    | 9.7              | 5.2   | 1.1 - 31.1    | 9.9               | 6.8   | 0.0 - 60.6    |
| Fruit and vegetable fibre (g/day) | 9.0              | 5.3   | 0.5 - 36.6    | 9.6               | 5.8   | 0.0 - 41.1    |
| Energy (MJ)                       | 9.26             | 3.32  | 3.01 - 20.14  | 9.51              | 3.49  | 3.24 - 20.07  |
| BMI (kg/m <sup>2</sup> )          | 22.50            | 3.79  | 15.03 - 38.47 | 23.31             | 4.90  | 14.33 - 50.16 |
| WHtR                              | 0.455            | 0.06  | 0.23 - 0.78   | 0.465             | 0.07  | 0.28 - 0.84   |
| hs-CRP (mg/L)                     | 1.13             | 1.61  | 0.14 - 9.39   | 1.52              | 2.05  | 0.14 - 9.69   |
| Log hs-CRP                        | 0.59             | 0.52  | 0.13 - 2.34   | 0.71              | 0.60  | 0.13 - 2.37   |
| Leptin (µg/L)                     | 17.20            | 18.40 | 1.40 - 153.60 | 21.51             | 26.11 | 1.40 - 203.00 |
| Log leptin                        | 2.38             | 1.08  | 0.88 - 5.04   | 2.47              | 1.18  | 0.88 - 5.32   |
| Adiponectin (mg/L)                | 9.51             | 4.81  | 1.00 - 42.80  | 9.53              | 6.13  | 1.70 - 74.00  |

BMI, body mass index; WHtR, waist height ratio; hs-CRP, high sensitivity C-reactive protein  
 Participants with hs-CRP>10mg/L are excluded from mean hs-CRP

**Table 3:** Weighted associations between dietary fibre intake and inflammatory markers measured at 17 years of age in the Raine Study

|                           | Log hs-CRP    |                      |              | Log leptin    |                        |                  | Adiponectin  |                      |                  |
|---------------------------|---------------|----------------------|--------------|---------------|------------------------|------------------|--------------|----------------------|------------------|
|                           | $\beta$       | 95% CI               | <i>P</i>     | $\beta$       | 95% CI                 | <i>P</i>         | $\beta$      | 95% CI               | <i>P</i>         |
| Dietary fibre             |               |                      |              |               |                        |                  |              |                      |                  |
| Model 1                   | -0.025        | -0.050, 0.00018      | 0.052        | <b>-0.13</b>  | <b>-0.17, -0.085</b>   | <b>&lt;0.001</b> | <b>-0.28</b> | <b>-0.49, -0.070</b> | <b>0.009</b>     |
| Model 2                   | -0.014        | -0.040, 0.011        | 0.273        | <b>-0.034</b> | <b>-0.062, -0.0059</b> | <b>0.018</b>     | -0.12        | -0.32, 0.085         | 0.250            |
| Model 3                   | -0.0052       | -0.036, 0.025        | 0.735        | -0.023        | -0.049, 0.0027         | 0.079            | -0.12        | -0.38, 0.14          | 0.361            |
| Model 4                   | 0.0083        | -0.023, 0.039        | 0.600        | -0.014        | -0.041, 0.013          | 0.324            | -0.13        | -0.40, 0.14          | 0.330            |
| Cereal and grain fibre    |               |                      |              |               |                        |                  |              |                      |                  |
| Model 1                   | <b>-0.064</b> | <b>-0.11, -0.021</b> | <b>0.004</b> | <b>-0.33</b>  | <b>-0.40, -0.27</b>    | <b>&lt;0.001</b> | <b>-0.73</b> | <b>-1.1, -0.37</b>   | <b>&lt;0.001</b> |
| Model 2                   | -0.034        | -0.080, 0.011        | 0.140        | <b>-0.067</b> | <b>-0.12, -0.018</b>   | <b>0.008</b>     | -0.28        | -0.64, 0.086         | 0.135            |
| Model 3                   | -0.029        | -0.077, 0.020        | 0.251        | <b>-0.062</b> | <b>-0.10, -0.020</b>   | <b>0.004</b>     | -0.24        | -0.66, 0.18          | 0.258            |
| Model 4                   | -0.014        | -0.063, 0.036        | 0.587        | <b>-0.057</b> | <b>-0.10, -0.014</b>   | <b>0.010</b>     | -0.21        | -0.64, 0.21          | 0.325            |
| Fruit and vegetable fibre |               |                      |              |               |                        |                  |              |                      |                  |
| Model 1                   | 0.0085        | -0.034, 0.051        | 0.697        | -0.0018       | -0.073, 0.070          | 0.960            | -0.25        | -0.61, 0.11          | 0.171            |
| Model 2                   | 0.0071        | -0.035, 0.049        | 0.741        | -0.014        | -0.060, 0.032          | 0.556            | -0.27        | -0.61, 0.069         | 0.118            |
| Model 3                   | 0.013         | -0.029, 0.055        | 0.547        | -0.012        | -0.048, 0.024          | 0.513            | -0.24        | -0.60, 0.12          | 0.191            |
| Model 4                   | 0.027         | -0.015, 0.068        | 0.211        | -0.0014       | -0.038, 0.035          | 0.941            | -0.23        | -0.59, 0.14          | 0.223            |

hs-CRP, high sensitivity C-reactive protein


Model 1 - dietary fibre (per 5g/day higher intake)

Model 2 - Model 1 + sex


Model 3 - Model 2 + waist height ratio, body mass index, energy intake, and dietary misreporting

Model 4 - Model 3 + ethnicity, screentime, physical activity, smoking, and alcohol intake

Weighted by inverse probability of having complete data and being included in the study



**Figure 1:** Number of participants in the Raine Study 17-year follow-up and number with complete variable data. Dietary data includes dietary fibre and energy intake. \*Some participants are missing data from multiple variables and therefore numbers do not add up to the total number of people excluded for missing data.



**Figure 2:** Mean daily dietary fibre intakes and concentrations of high sensitivity C-reactive protein (hs-CRP), leptin, and adiponectin in 17-year-olds from the Raine Study separated by sex.

## References

1. Medzhitov R (2008) Origin and physiological roles of inflammation. *Nature* 454, 428-435.
2. Minihane AM, Vinoy S, Russell WR *et al.* (2015) Low-grade inflammation, diet composition and health: current research evidence and its translation. *The British journal of nutrition* 114, 999-1012.
3. Miller AH, Maletic V Raison CL (2009) Inflammation and its discontents: the role of cytokines in the pathophysiology of major depression. *Biological psychiatry* 65, 732-741.
4. Hotamisligil GS (2006) Inflammation and metabolic disorders. *Nature* 444, 860-867.
5. Meyer KA, Kushi LH, Jacobs DR, Jr. *et al.* (2000) Carbohydrates, dietary fiber, and incident type 2 diabetes in older women. *The American journal of clinical nutrition* 71, 921-930.
6. Threapleton DE, Greenwood DC, Evans CEL *et al.* (2013) Dietary fibre intake and risk of cardiovascular disease: systematic review and meta-analysis. *BMJ (Clinical research ed)* 347, f6879.
7. Chen J-P, Chen G-C, Wang X-P *et al.* (2018) Dietary Fiber and Metabolic Syndrome: A Meta-Analysis and Review of Related Mechanisms. *Nutrients* 10, 24.
8. Lie L, Brown L, Forrester T *et al.* (2018) The Association of Dietary Fiber Intake with Cardiometabolic Risk in Four Countries across the Epidemiologic Transition. *Nutrients* 10, E628.
9. Parikh S, Pollock NK, Bhagatwala J *et al.* (2012) Adolescent fiber consumption is associated with visceral fat and inflammatory markers. *The Journal of clinical endocrinology and metabolism* 97, E1451-1457.
10. Cummings JH (1981) Short chain fatty acids in the human colon. *Gut* 22, 763-779.
11. Tong L-C, Wang Y, Wang Z-B *et al.* (2016) Propionate Ameliorates Dextran Sodium Sulfate-Induced Colitis by Improving Intestinal Barrier Function and Reducing Inflammation and Oxidative Stress. *Front Pharmacol* 7, 253.
12. Macfarlane GT & Macfarlane S (2012) Bacteria, colonic fermentation, and gastrointestinal health. *Journal of AOAC International* 95, 50-60.
13. den Besten G, van Eunen K, Groen AK *et al.* (2013) The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. *Journal of lipid research* 54, 2325-2340.
14. Tremaroli V & Backhed F (2012) Functional interactions between the gut microbiota and host metabolism. *Nature* 489, 242-249.
15. Schedlowski M, Engler H Grigoleit JS (2014) Endotoxin-induced experimental systemic inflammation in humans: a model to disentangle immune-to-brain communication. *Brain, behavior, and immunity* 35, 1-8.
16. Pepys MB & Hirschfield GM (2003) C-reactive protein: a critical update. *The Journal of clinical investigation* 111, 1805-1812.
17. Eklund CM (2009) Proinflammatory cytokines in CRP baseline regulation. *Advances in clinical chemistry* 48, 111-136.
18. Slavin JL (2008) Position of the American Dietetic Association: health implications of dietary fiber. *Journal of the American Dietetic Association* 108, 1716-1731.
19. Cox AJ, West NP Cripps AW (2015) Obesity, inflammation, and the gut microbiota. *The lancet Diabetes & endocrinology* 3, 207-215.

20. Miller GE, Freedland KE, Carney RM *et al.* (2003) Pathways linking depression, adiposity, and inflammatory markers in healthy young adults. *Brain, behavior, and immunity* 17, 276-285.
21. Shelton RC & Miller AH (2010) Eating ourselves to death (and despair): the contribution of adiposity and inflammation to depression. *Progress in neurobiology* 91, 275-299.
22. Scotece M, Conde J, Lopez V *et al.* (2014) Adiponectin and leptin: new targets in inflammation. *Basic & clinical pharmacology & toxicology* 114, 97-102.
23. Procaccini C, La Rocca C, Carbone F *et al.* (2017) Leptin as immune mediator: Interaction between neuroendocrine and immune system. *Dev Comp Immunol* 66, 120-129.
24. Nedvidkova J, Smitka K, Kopsky V *et al.* (2005) Adiponectin, an adipocyte-derived protein. *Physiological research* 54, 133-140.
25. Murakami K, Sasaki S, Takahashi Y *et al.* (2007) Nutrient and food intake in relation to serum leptin concentration among young Japanese women. *Nutrition (Burbank, Los Angeles County, Calif)* 23, 461-468.
26. Kuroda M, Ohta M, Okufuji T *et al.* (2010) Frequency of soup intake and amount of dietary fiber intake are inversely associated with plasma leptin concentrations in Japanese adults. *Appetite* 54, 538-543.
27. Edwards CA, Xie C Garcia AL (2015) Dietary fibre and health in children and adolescents. *The Proceedings of the Nutrition Society* 74, 292-302.
28. Fayet-Moore F, Cassettari T, Tuck K *et al.* (2018) Dietary Fibre Intake in Australia. Paper I: Associations with Demographic, Socio-Economic, and Anthropometric Factors. *Nutrients* 10, 599.
29. Metcalf SA, Jones PB, Nordstrom T *et al.* (2017) Serum C-reactive protein in adolescence and risk of schizophrenia in adulthood: A prospective birth cohort study. *Brain, behavior, and immunity* 59, 253-259.
30. Biro FM & Wien M (2010) Childhood obesity and adult morbidities. *The American journal of clinical nutrition* 91, 1499S-1505S.
31. Qi L, van Dam RM, Liu S *et al.* (2006) Whole-grain, bran, and cereal fiber intakes and markers of systemic inflammation in diabetic women. *Diabetes care* 29, 207-211.
32. Qi L, Rimm E, Liu S *et al.* (2005) Dietary glycemic index, glycemic load, cereal fiber, and plasma adiponectin concentration in diabetic men. *Diabetes care* 28, 1022-1028.
33. AlEsa HB, Ley SH, Rosner B *et al.* (2016) High Fiber and Low Starch Intakes Are Associated with Circulating Intermediate Biomarkers of Type 2 Diabetes among Women. *The Journal of nutrition* 146, 306-317.
34. Straker L, Mountain J, Jacques A *et al.* (2017) Cohort profile: the Western Australian Pregnancy Cohort (Raine) Study-Generation 2. *International journal of epidemiology* 46, 1384-1385j.
35. White SW, Eastwood PR, Straker LM *et al.* (2017) The Raine study had no evidence of significant perinatal selection bias after two decades of follow up: a longitudinal pregnancy cohort study. *BMC pregnancy and childbirth* 17, 207.
36. Lassale C, Guilbert C, Keogh J *et al.* (2009) Estimating food intakes in Australia: validation of the Commonwealth Scientific and Industrial Research Organisation (CSIRO) food frequency questionnaire against weighed dietary intakes. *Journal of human nutrition and dietetics : the official journal of the British Dietetic Association* 22, 559-566.
37. Ambrosini GL, de Klerk NH, O'Sullivan TA *et al.* (2009) The reliability of a food frequency questionnaire for use among adolescents. *European journal of clinical nutrition* 63, 1251-1259.

38. Goldberg GR, Black AE, Jebb SA *et al.* (1991) Critical evaluation of energy intake data using fundamental principles of energy physiology: 1. Derivation of cut-off limits to identify under-recording. *European journal of clinical nutrition* 45, 569-581.
39. Appannah G, Pot GK, O'Sullivan TA *et al.* (2014) The reliability of an adolescent dietary pattern identified using reduced-rank regression: comparison of a FFQ and 3 d food record. *The British journal of nutrition* 112, 609-615.
40. Maffei C, Banzato C, Talamini G (2008) Waist-to-Height Ratio, a Useful Index to Identify High Metabolic Risk in Overweight Children. *J Pediatr* 152, 207-213.e202.
41. Pearson TA, Mensah GA, Alexander RW *et al.* (2003) Markers of inflammation and cardiovascular disease: application to clinical and public health practice: A statement for healthcare professionals from the Centers for Disease Control and Prevention and the American Heart Association. *Circulation* 107, 499-511.
42. McBee M (2010) Modeling Outcomes With Floor or Ceiling Effects: An Introduction to the Tobit Model. *Gift Child Q* 54, 314-320.
43. R Core Team (2018) R: A language and environment for statistical computing, 3.5.0 ed. Vienna, Austria: R Foundation for Statistical Computing.
44. RStudio (2018) RStudio, 1.1.453 ed. Boston RStudio.
45. Zhang R, Jiao J, Zhang W *et al.* (2016) Effects of cereal fiber on leptin resistance and sensitivity in C57BL/6J mice fed a high-fat/cholesterol diet. *Food Nutr Res* 60, 31690.
46. Liu S, Willett WC, Manson JE *et al.* (2003) Relation between changes in intakes of dietary fiber and grain products and changes in weight and development of obesity among middle-aged women. *The American journal of clinical nutrition* 78, 920-927.
47. Lopez-Jaramillo P, Gomez-Arbelaes D, Lopez-Lopez J *et al.* (2014) The role of leptin/adiponectin ratio in metabolic syndrome and diabetes. *Horm Mol Biol Clin Invest* 18, 37-45.
48. Hajihashemi P, Azadbakht L, Hashemipour M *et al.* (2014) Whole-grain intake favorably affects markers of systemic inflammation in obese children: A randomized controlled crossover clinical trial. *Molecular Nutrition & Food Research* 58, 1301-1308.
49. Davis JN, Alexander KE, Ventura EE *et al.* (2009) Inverse relation between dietary fiber intake and visceral adiposity in overweight Latino youth. *The American journal of clinical nutrition* 90, 1160-1166.
50. Fardet A (2010) New hypotheses for the health-protective mechanisms of whole-grain cereals: what is beyond fibre? *Nutr Res Rev* 23, 65-134.
51. Iikuni N, Lam QLK, Lu L *et al.* (2008) Leptin and inflammation. *Curr Immunol Rev* 4, 70-79.
52. Shen J, Sakaida I, Uchida K *et al.* (2005) Leptin enhances TNF-alpha production via p38 and JNK MAPK in LPS-stimulated Kupffer cells. *Life sciences* 77, 1502-1515.
53. Huang R-C, Mori TA, Burke V *et al.* (2009) Synergy Between Adiposity, Insulin Resistance, Metabolic Risk Factors, and Inflammation in Adolescents. *Diabetes care* 32, 695.
54. Le-Ha C, Beilin LJ, Burrows S *et al.* (2014) Gender and the active smoking and high-sensitivity C-reactive protein relation in late adolescence. *Journal of lipid research* 55, 758-764.
55. Ayonrinde OT, Olynyk JK, Beilin LJ *et al.* (2011) Gender-specific differences in adipose distribution and adipocytokines influence adolescent nonalcoholic fatty liver disease. *Hepatology* 53, 800-809.
56. Ford ES, Giles WH, Myers GL *et al.* (2003) C-reactive protein concentration distribution among US children and young adults: findings from the National Health and Nutrition Examination Survey, 1999-2000. *Clinical chemistry* 49, 1353-1357.

57. Galic S, Oakhill JS Steinberg GR (2010) Adipose tissue as an endocrine organ. *Mol Cell Endocrinol* 316, 129-139.
58. Ebrahimi M, Heidari-Bakavoli AR, Shoeibi S *et al.* (2016) Association of Serum hs-CRP Levels With the Presence of Obesity, Diabetes Mellitus, and Other Cardiovascular Risk Factors. *J Clin Lab Anal* 30, 672-676.
59. Christen T, Trompet S, Noordam R *et al.* (2018) Sex differences in body fat distribution are related to sex differences in serum leptin and adiponectin. *Peptides* 107, 25-31.
60. Blum WF, Englaro P, Hanitsch S *et al.* (1997) Plasma leptin levels in healthy children and adolescents: dependence on body mass index, body fat mass, gender, pubertal stage, and testosterone. *The Journal of clinical endocrinology and metabolism* 82, 2904-2910.
61. Mazidi M, Kengne AP, Mikhailidis DP *et al.* (2018) Effects of selected dietary constituents on high-sensitivity C-reactive protein levels in U.S. adults. *Ann Med* 50, 1-6.
62. Bo S, Durazzo M, Guidi S *et al.* (2006) Dietary magnesium and fiber intakes and inflammatory and metabolic indicators in middle-aged subjects from a population-based cohort. *The American journal of clinical nutrition* 84, 1062-1069.
63. Ajani UA, Ford ES Mokdad AH (2004) Dietary fiber and C-reactive protein: findings from national health and nutrition examination survey data. *The Journal of nutrition* 134, 1181-1185.
64. King DE, Egan BM Geesey ME (2003) Relation of dietary fat and fiber to elevation of C-reactive protein. *The American journal of cardiology* 92, 1335-1339.
65. Ma Y, Griffith JA, Chasan-Taber L *et al.* (2006) Association between dietary fiber and serum C-reactive protein. *The American journal of clinical nutrition* 83, 760-766.
66. Murakami K, Sasaki S, Takahashi Y *et al.* (2008) Total n-3 polyunsaturated fatty acid intake is inversely associated with serum C-reactive protein in young Japanese women. *Nutrition research (New York, NY)* 28, 309-314.
67. Ma Y, Hebert JR, Li W *et al.* (2008) Association between dietary fiber and markers of systemic inflammation in the Women's Health Initiative Observational Study. *Nutrition (Burbank, Los Angeles County, Calif)* 24, 941-949.
68. David LA, Maurice CF, Carmody RN *et al.* (2013) Diet rapidly and reproducibly alters the human gut microbiome. *Nature* 505, 559.
69. Considine RV, Sinha MK, Heiman ML *et al.* (1996) Serum immunoreactive-leptin concentrations in normal-weight and obese humans. *The New England journal of medicine* 334, 292-295.

### ***Acknowledgements***

We are grateful to the Raine Study participants and their families and we thank the Raine Study research staff for cohort coordination and data collection. The core management of the Raine Study is funded by the University of Western Australia, Curtin University, Telethon Kids Institute, Women and Infants Research Foundation, Edith Cowan University, Murdoch University, The University of Notre Dame Australia and the Raine Medical Research Foundation. We thank the National Health and Medical Research Council of Australia (NHMRC) for their long-term contribution to funding the study over the last 30 years and the Telethon Kids Institute for long term support of the Raine Study.

### ***Funding***

This work was supported by the Heart Foundation Beyond Blue Strategic Research Program (Oddy et al ID G08P4036 2009-2012). The Raine Study received funding from the Raine Medical Research Foundation at The University of Western Australia, the NHMRC, the Telstra Research Foundation, the Western Australian Health Promotion Foundation and Australian Rotary Health Research Fund. Data collection and biological specimens at the 17 years follow-up were funded by the NHMRC Program Grant ID 353514 and Project Grant #403981. TAM is supported by an NHMRC Research Fellowship (ID 1136046).

### ***Conflict of interest***

All authors declare that there are no potential or existing conflicts of interest.

### ***Authorship contributions***

Olivia G Swann - Formulating the research question, study design, analysis, interpretation, drafting and preparation of final manuscript

Monique Breslin - Formulating the research question, study design, analysis, interpretation, drafting and review of final manuscript

Michelle Kilpatrick - Formulating the research question, study design, interpretation, drafting and review of final manuscript

Therese A O'Sullivan - Study design, interpretation, review of final manuscript

Trevor A Mori - Study design, interpretation, review of final manuscript

Lawrence J Beilin - Study design, interpretation, review of final manuscript

Wendy H Oddy - Formulating the research question, study design, interpretation, drafting and review of final manuscript