

BMJ Open Acceptance commitment therapy (ACT) for psychological distress associated with inflammatory bowel disease (IBD): protocol for a feasibility trial of the ACTforIBD programme

Subhadra Evans ¹, Lisa Olive,² Madeleine Dober,³ Simon Knowles,⁴ Matthew Fuller-Tyszkiewicz,³ Eric O,⁵ Peter Gibson,⁶ Leanne Raven,⁷ Richard Geary,⁸ Andrew McCombie,⁹ Leesa van Niekerk,¹⁰ Susan Chesterman,² Daniel Romano,² Antonina Mikocka-Walus ¹

To cite: Evans S, Olive L, Dober M, *et al.* Acceptance commitment therapy (ACT) for psychological distress associated with inflammatory bowel disease (IBD): protocol for a feasibility trial of the ACTforIBD programme. *BMJ Open* 2022;**12**:e060272. doi:10.1136/bmjopen-2021-060272

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-060272>).

Received 19 December 2021
Accepted 20 May 2022



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Subhadra Evans;
subhadra.evans@deakin.edu.au

ABSTRACT

Introduction Inflammatory bowel disease (IBD) involves an abnormal immune response to healthy gut bacteria. When a person develops IBD, their susceptibility to anxiety and/or depression increases. The ACTforIBD programme, specifically designed for people with IBD and comorbid psychological distress, draws on acceptance and commitment therapy (ACT), which promotes acceptance of situations that cannot be solved such as persistent physical symptoms. There are no ACT trials for IBD using an active control group or a telemedicine approach, which is important to improve accessibility, particularly in the context of the ongoing COVID-19 pandemic. The ACTforIBD programme is administered online with a 4-hour therapist involvement per participant only; if successful it can be widely implemented to improve the well-being of many individuals with IBD.

Methods and analysis Our team have codesigned with consumers the ACTforIBD programme, an 8-week intervention of 1-hour sessions, with the first three sessions and the last session delivered one-to-one by a psychologist, and the other sessions self-directed online. This study aims to evaluate the feasibility and preliminary efficacy of ACTforIBD to reduce psychological distress in patients with IBD. Using a randomised controlled trial, 25 participants will be randomised to ACTforIBD, and 25 patients to an active control condition.

Ethics and dissemination This protocol has been approved by Deakin University Research Ethics Committee in September 2021 (Ref. 2021-263) and the New Zealand Central Health and Disability Ethics Committee in December 2021 (Ref. 2021 EXP 11384). The results of this research will be published in peer-reviewed journals and shared with various stakeholders, including community members, policy-makers and researchers, through local and international conferences.

Trial registration number ACTRN12621001316897.

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ This feasibility randomised controlled trial of acceptance and commitment therapy (ACT) in people with inflammatory bowel disease (IBD) will use an active control group (a psychoeducation-focused programme adapted to IBD).
- ⇒ ACTforIBD is an 8-week blended-delivery intervention, with the first three sessions and the last session delivered one-to-one by a psychologist, and the other sessions self-directed online.
- ⇒ Participants will be randomised to ACTforIBD and an active control condition in equal numbers (1:1), but the study is not powered to examine efficacy as its focus is on feasibility of ACTforIBD.
- ⇒ This internet-delivered study may be slow to recruit and associated with high attrition rates due to COVID-19-related internet fatigue.

INTRODUCTION

Inflammatory bowel disease (IBD) is an umbrella term for several relapsing-remitting inflammatory conditions of the gastrointestinal tract involving an abnormal immune response to healthy gut bacteria in those with a genetic predisposition.¹ The most common forms of IBD are Crohn's disease and ulcerative colitis. Key IBD-related symptoms include chronic pain, bloody diarrhoea with frequent bowel movements, weight loss, anaemia and fatigue, causing significant impairments in daily functioning. The disease affects as many as 3 million people in the USA,² with the prevalence expected to reach 1% in the Western world by 2030.³

Bidirectional gut-brain links are considered part of the pathophysiology of IBD; when a person develops IBD, their susceptibility to anxiety and/or depression increases.⁴ Up

to 66% of individuals report symptoms during an active disease phase.⁵ Our meta-analyses found that people with IBD experience a consistently poor quality of life (QoL) both during disease flare-ups and remission.^{6,7} Further, IBD is negatively affected by mental disorders, resulting in more frequent flare-ups,⁸ a more aggressive presentation,⁹ increased rates of hospital readmissions and increased risk of surgery.¹⁰ In addition, a registry-based observational study has shown that the top 5% of patients with highest medical charges had high prevalence of depression, anxiety and chronic pain, suggesting that these comorbidities need to be managed adequately to reduce the financial burden of IBD.¹¹

Our meta-analysis¹² demonstrates the efficacy of psychotherapy in improving QoL and depression in IBD. However, the findings also highlight the urgent need for trials of psychotherapy interventions targeting the groups at risk of suboptimal outcomes (eg, people with psychological distress), to improve both mental and physical health management in IBD. This is because most studies conducted to date have focused on people without psychological comorbidities and this could have contributed to low efficacy of psychotherapy in IBD on psychological outcomes such as anxiety or coping.

The ACTforIBD programme, specifically designed for people with IBD and comorbid psychological distress, is an 8-week intervention that draws on acceptance and commitment therapy (ACT). ACT promotes acceptance of situations that cannot be solved such as persistent physical symptoms, and is taught as an alternative to experiential avoidance. The application of acceptance processes, along with commitment and value-directed behaviour change processes, is aimed at increasing psychological flexibility (eg, being fully present in the present moment, without trying to avoid/fight unwanted thoughts/feelings).¹³ The psychological flexibility afforded by ACT is associated with reduced psychological distress in debilitating chronic diseases.^{14,15} ACT is derived from cognitive-behavioural therapy (CBT). Both these therapies support adaptive emotion regulation but while CBT focuses on antecedent emotion regulation, ACT targets maladaptive regulation strategies such as suppression.¹⁶ CBT has been found to be effective in reducing symptoms of depression and improving QoL in IBD.¹² ACT demonstrates similar efficacy to CBT for improving psychological distress in the general population^{17,18} and those with pain-related chronic illness.¹⁸

To date, only one previous trial examined ACT in the treatment of IBD,¹⁹ demonstrating its preliminary efficacy in improving levels of depression and QoL. Another trial is currently in progress.²⁰ Both these trials use an inactive control group (ie, treatment as usual). There are no ACT trials utilising an active control group or combining therapist-led sessions with self-directed online sessions (a telemedicine approach). These modalities are an important advancement to improve accessibility, particularly in the context of the ongoing COVID-19 pandemic. To address this gap, our team have codesigned with consumers the ACTforIBD program.²¹

This study aims to evaluate the feasibility and efficacy potential of ACTforIBD to reduce psychological distress in patients with IBD and inform the subsequent design of a definitive large-scale efficacy trial. Using a randomised controlled trial (RCT), our primary objectives are:

- ▶ The feasibility of recruitment to a full trial, reasons for withdrawal and session attendance in the ACTforIBD group versus an active control group at 8 weeks since baseline (primary outcomes).
- ▶ Consumer satisfaction with ACTforIBD at 8 weeks since baseline and 3 months postintervention (secondary outcomes).

Noting both uncertainties in effect estimates that may arise from small pilot studies²² and absence of suitable prior studies to guide sample size guidelines for a definitive RCT for this population, our secondary objectives are to:

- ▶ Obtain estimates (rate of completion, mean differences at post-treatment timepoints and associated 95% CIs and SD on outcome measures) that may both enable evaluation of efficacy potential of ACTforIBD in improving patient psychological distress, disease activity, QoL, fatigue, pain, resilience and self-efficacy at 8 weeks since baseline and 3 months postintervention (secondary outcomes) and facilitate sample size calculations for a large-scale RCT.
- ▶ Derive preliminary indication of cost-effectiveness based on healthcare utilisation (secondary outcomes).

METHODS AND ANALYSIS

Ethical approval

This protocol has been approved by Deakin University Research Ethics Committee in September 2021 (Ref. 2021–263) and the New Zealand Central Health and Disability Ethics Committee in December 2021 (Ref. 2021 EXP 11384). The trial will adhere to the Consolidated Standards of Reporting Trials (CONSORT) statement.

Design

A parallel, mixed-methods, randomised, controlled feasibility trial will be conducted. Since psychological interventions for IBD tested to date have not been codesigned with consumers and offer only limited efficacy,¹² we will incorporate a comprehensive qualitative evaluation alongside the quantitative data collection. Participants will be randomly allocated with a 1:1 ratio to one of two groups: the ACTforIBD programme or active control. Randomisation will be stratified by IBD activity measured on the Manitoba Index.²³ Those with no to low activity (occasionally active, giving me symptoms 1–2 days/month; rarely active, giving me symptoms on a few days in the past 3 months; and I was well in the past 3 months, what I consider a remission or absence of symptoms) will be grouped into one category and those with moderate to high activity into another (Constantly active, giving me symptoms every day; Often active, giving me symptoms most days; and Sometimes active, giving me symptoms on some days). Computer-generated random sequences will

be embedded in the Qualtrics platform ensuring allocation concealment. Participants and all the investigators including statisticians will be blinded to group allocation. However, the trial managers enrolling participants and psychologists delivering therapy will not be blinded, for practical reasons. Participants in both treatment arms will remain on their treatment as usual. All participants will complete questionnaires before and after the intervention (8 weeks since baseline and 3 months postintervention) and a subsample will also participate in qualitative interviews at 8 weeks since baseline and 3 months postintervention. The trial will follow the intention-to-treat paradigm.

Participants

Inclusion criteria

1. Diagnosis of IBD: Crohn's disease, ulcerative colitis or indeterminate colitis established by a health professional using standard criteria (participants will need to provide a letter from their general practitioner or gastroenterologist or provide other proof of diagnosis such as endoscopy reports).
2. Psychological distress: scores 16–29 (mild to moderate distress) on Kessler Psychological Distress Scale (K10).²⁴
3. Age: 18 years and older.
4. People with IBD residing in Australia or New Zealand, able to read and write in English, with access to the internet to ensure participation in the online intervention.

Exclusion criteria

1. No psychological distress (scores <16 on K10) or severe distress (scores 30–50 on K10).

Participants with active IBD can participate in the study and a flare will not prevent them from continuing the intervention should they wish to remain in the study.

Recruitment

We will recruit via consumer organisations and IBD-related social media, for example, Facebook, Twitter, Instagram of Crohn's & Colitis Australia (a social media following of 30 000 across all platforms). Crohn's and Colitis New Zealand will assist with recruitment in a similar way. We will ask participants for confirmation of diagnosis. It is anticipated that recruitment will take 1 year, with 4–5 participants recruited per month.

Sample size

Feasibility studies do not require power calculations as they are not focused on efficacy. However, the selected sample size needs to be adequate to estimate clinical parameters with a degree of precision²⁵ and, where possible, should factor in a plausible target effect size for a subsequent, definitive RCT.²⁶ In the absence of prior findings to accurately inform a likely effect size, we used a recommended, reasonably conservative standardised mean difference of 0.30.²⁷ Based on this value, and Whitehead *et al*'s²⁶

Box 1 The ACTforIBD programme modules

- ⇒ Module 1: Commitment to therapy and assessment (therapist led).
 - ⇒ Module 2: Creative hopelessness (therapist led).
 - ⇒ Module 3: Personal values (therapist led).
 - ⇒ Module 4: Mindfulness (self-directed).
 - ⇒ Module 5: Fusion and defusion (self-directed).
 - ⇒ Module 6: Acceptance (self-directed).
 - ⇒ Module 7: Values and goals (self-directed)
 - ⇒ Module 8: Commitment and overcoming barriers moving forward (therapist led).
- ACT, acceptance and commitment therap; IBD, Inflammatory bowel disease.

calculations for the optimal pilot study sample size, we aim to recruit 50 participants (25 per group).

Intervention and control condition

The ACTforIBD programme

People with IBD experience unique worries related to QoL (eg, toilet use frequency, travelling, embarrassment), unpredictability (of flares in symptoms), the symptoms themselves and treatment concerns (eg, medication side effects).²⁸ Our intervention is consistent with the core principles of ACT (eg, mindfulness, value-based decision making, defusion²⁹ and tailored to people with IBD to relate to some of these key concerns).²⁸ The intervention includes eight fully-scripted modules (each session is broken down into activities which need to be delivered), each addressing a different ACT process, with each session building on the previous process (see [box 1](#)). It has a blended design: the first three sessions and the last session are therapist-led and delivered one-to-one (via a Zoom/Skype appointment scheduled by the trial managers with the same therapist working with a given participant throughout the programme), and the between sessions are self-led (using a video/podcast available on our website without restrictions plus self-assessment activities). The protocol for the intervention was codesigned with consumers.²¹ The sessions occur weekly for 8 weeks and take 1 hour (duration established as effective.³⁰ The therapist-led individual sessions will be delivered by provisionally registered psychologists under the supervision of a fully registered clinical psychologist, with extensive training and experience in the provision of ACT. Quality assurance will include regular therapist supervision and session content audits conducted by Dr van Niekerk (senior psychologist). While we will not be formally recording and comparing the delivery of sessions by various therapists, fidelity of the intervention is monitored through training of the therapists (all therapists will be provided with a manual explaining IBD, its treatment, psychological comorbidities, etc), use of detailed scripts for each session (ie, instructions for therapists of what needs to be discussed with the participant with any relevant worksheets provided to both the participant and therapist) and group supervision where adherence to scripts is discussed.

Box 2 The active control group programme

- ⇒ Module 1: Stress management (therapist led).
- ⇒ Module 2: Progressive muscle relaxation (therapist led).
- ⇒ Module 3: Problem solving (therapist led).
- ⇒ Module 4: Coping strategies (self-directed).
- ⇒ Module 5: How to think more assertively (self-directed).
- ⇒ Module 6: How to behave more assertively (self-directed).
- ⇒ Module 7: Sleeping for better well-being (self-directed).
- ⇒ Module 8: Towards a healthy self-esteem (therapist led).

The goals of treatment with this programme broadly include: assisting clients to define valued directions and take committed action in the direction of their identified values; increasing understanding and insight into the client's personal experiences by promoting presence and increasing cognitive defusion; improved coping with physical symptoms through the process of acceptance and cognitive defusion; and increasing psychological flexibility around psychological distress associated with IBD. The specific goals of treatment with each person will vary depending on their symptoms, life experiences and importantly, what they want to gain from therapy.

The active control group will be asked to use a psychoeducation-focused programme adapted to IBD based on the Centre for Clinical Interventions materials (<https://www.cci.health.wa.gov.au/Resources/Looking-After-Yourself>) (see [box 2](#)).

The duration of the control condition will mirror the experimental group with 8 weeks of 1-hour involvement with the website recommended. Participant demands across both groups will be the same including allocated time for homework activities. This group will also be facilitated in a similar fashion to the ACT group, that is, the first three sessions and the last session are therapist-led one-to-one (via a Zoom/Skype appointment). The therapists will be advised to focus on the content of the sessions rather than therapeutic processes. Experience with this and potential challenges will be monitored via supervision.

Withdrawal criteria

Participants are free to withdraw at any time. We will ask about their reasons for withdrawal for statistical and reporting purposes but answering this question will not be compulsory. No aspect of participant IBD care will be affected by their decision to withdraw from the study. We will monitor dropout/attrition closely to be able to establish satisfaction with the intervention.

Measures

Quantitative measures

Measures of feasibility for progression to a full RCT

- ▶ Feasibility of recruitment assessed based on data from study database (n=50 recruited during a 1-year period of recruitment).

- ▶ Feasibility of retention in the study based on data from study database (at least 80% of recruited participants retained at 8 weeks follow-up).

Consumer satisfaction for progression to a full RCT: see [table 1](#) for detail re measures of satisfaction. In addition, reasons for withdrawal and session attendance (at least 50% of sessions attended) will be examined.

Measures for evaluating efficacy potential: patient psychological distress, disease activity, QoL, fatigue, pain, resilience and self-efficacy. See [table 1](#) for detail of the scales including their scoring and assessment time.

Preliminary cost-effectiveness: see online supplemental appendix 1 for detail of treatment utilisation scale.

Qualitative interviews

A rich sample capturing key demographics (representatives of different genders, age groups, urban/rural location, IBD types, group allocation) of 20 participants will be invited to participate in a longitudinal qualitative study consisting of 2 semistructured interviews conducted via phone or zoom/Teams (~30–45 min) immediately postintervention (8 weeks)^{31 32} and at 3 months postintervention. The longitudinal design of this qualitative study follows qualitative principles of change across time,³³ and will help us monitor the use of ACT strategies over time, to inform ongoing support. The proposed sample size is consistent with recommended guidelines for thematic analysis.³¹ The aim of the interviews is to understand: (1) perceived barriers and enablers to undertaking the programme, including elements of the programme that are of continued use and (2) meaningful change experienced by participants. To understand meaningful change, we will use most significant change (MSC) principles, an interviewing technique developed for the purpose of evaluating complex interventions.³² Interview questions begin with inquiry about the individual's circumstances, or 'story' about what led them to consider being involved in the study, and narrow in focus to ask participants what the MSCs were for them during the intervention, and postintervention. Interviews aim to understand what significant changes were experienced, why these changes were significant for the individual, and what meaning the individual attributed to these changes. The changes reported by participants in interviews will triangulate quantitative outcomes.

Procedure

Participants will largely be recruited online via social media. They will be asked to email the researchers or complete an online expression of interest form if they are interested in participating. Participants will then be asked to read and consider the study's plain language statement which tells them that they will be allocated randomly to one of the two study groups and, if still interested, to sign a consent form and return to the investigators along with evidence of their IBD diagnosis via email. After consenting, participants will be asked to complete the K10, and the trial managers will

Table 1 Quantitative measures

		No of items	Assessment time
Demographics	Age, sex, level of education, marital status, employment, language spoken at home, postcode, private insurance	8	Baseline
Health-related questions	IBD subtype (CD, UC, IC)	1	Baseline
	When was your IBD diagnosed?	1	Baseline
	Do you currently have any of the following (click all that apply): Stoma (bag); Fistula; Perianal disease; Unsure	1	Baseline
	Do you suffer from other chronic illnesses (eg, diabetes, arthritis, depression)? If so, please list.	1	Baseline
	Please list all your prescribed, over the counter and complementary therapies	1	Baseline
	What treatment do you currently take for IBD?	1	Baseline
	Do you regularly use opioid medication such as oxycontin, codeine, tramadol, fentanyl or similar painkillers? If yes, please list.	1	Baseline
	Do you take antidepressants or anti-anxiety medication?	1	Baseline
	Are you currently seeing a mental health professional?	1	Baseline
	Smoking habits	1	Baseline
	Alcohol	1	Baseline
	BMI (weight and height)	2	Baseline
	IBD activity	IBD Control Scale ⁵⁵	14
Manitoba index ²³		1	Baseline and postintervention, follow-up
PRO3 for Crohn's disease ⁵⁶		3	Baseline and postintervention, follow-up
1-item re usual bowel function		1	Baseline and postintervention, follow-up
PRO2 for ulcerative colitis ⁵⁷		2	Baseline and postintervention, follow-up
Pain	A Numeric Rating Pain Scale (on a scale 0–10, with 0 meaning no pain and 10 meaning the worst possible pain, how much pain are you having now?)	1	Baseline and postintervention, follow-up
	Gastrointestinal Unhelpful Thinking Scale ⁵⁸	15	Baseline and postintervention, follow-up
Fatigue	Fatigue Symptom Inventory ⁵⁹	14	Baseline and postintervention, follow-up
Measure of mental health	Kessler Psychological Distress Scale ²⁴	10	Baseline to confirm eligibility
	DASS-21 (depression, anxiety, somatisation symptoms) ⁶⁰	21	Baseline and postintervention, follow-up
	EQ5D5L ⁶¹	6	Baseline and postintervention, follow-up
	Brief Resilience Scale ⁶²	6	Baseline and postintervention, follow-up
	General Self-efficacy Scale ⁶³	10	Baseline and postintervention, follow-up
	Acceptance and Action Scale ⁶⁴	7	Baseline and postintervention, follow-up
Satisfaction	A 0–10 satisfaction rating scale: On a scale from 0 to 10 how satisfied are you with the intervention you participated in? Open-ended questions: For example: What did you find most helpful about the programme? What would you say to a friend interested in taking part? The Credibility/Expectancy Questionnaire ⁶⁵	10	Postintervention
Safety	Did the intervention produce any undesired effects? If yes, please specify	1	
Treatment utilisation	A series of questions on outpatient and emergency department visits; hospitalisations; diagnostic and investigation use, etc	8–9	Baseline and postintervention, follow-up

BMI, body mass index; IBD, inflammatory bowel disease.

confirm their eligibility. Those eligible will complete the baseline quantitative measures via a Qualtrics link. Participants will then be randomised to one of the two groups by the randomisation software built into Qualtrics, informed about the starting date and sent the Zoom link with any necessary instructions. Participants

will meet weekly with the psychologist for the first 3 weeks, will then switch to self-directed learning in weeks 4–7, and have their final session with the psychologist again. After the final day of the intervention, participants will be asked to complete the quantitative post-intervention measures via Qualtrics. They will then be

contacted after 3 months postintervention to complete the follow-up measures via Qualtrics. An \$A30 gift card will be offered to participants who complete all three surveys (baseline, 8 weeks and 3 months postintervention). Participants invited to take part in the longitudinal qualitative aspect of the study will also receive an \$A30 gift card when they complete both interviews.

The researchers will use a range of evidence-based cohort retention strategies to minimise attrition, including having a facilitator and a short intervention duration, sending reminders and minimising time spent completing surveys.³⁴ At the trial completion, participants will be debriefed about their group allocation. All participants who express interest in receiving a summary data leaflet will be provided with one via email. In addition, the researchers will post this document on the consumer organisations' websites and associated social media platforms.

Data analysis

Quantitative analysis

Analyses will be carried out on an intention-to-treat basis. As feasibility trials cannot establish efficacy, the analysis will be largely descriptive (including measures of uncertainty, such as 95% CIs). However, we will attempt to run a group comparison on the main outcome measures with linear mixed effects models, which enable retention of participants with missing waves of data, under the assumption that data are missing at random. In these models, the dependent variable is the outcome measure, with predictors being Time, Group and the Interaction between Time and Group. The coefficient for interaction and its 95% CI will provide a range of estimates for change attributable to the target intervention. Following Bell *et al*,²² estimates of the point estimate and upper CI values (at 80%) for SD in the outcome measures, as well as estimates of drop-out and the time \times group intervention effect will be used to calculate a range of possible sample sizes required for a definitive RCT. Consistent with CONSORT guidelines³⁵ for pilot and feasibility studies, these sample size calculations will be offered tentatively and with caveats concerning potential instability in estimates from small sampled pilots.

A trial-based cost–utility analysis will evaluate the incremental resource use and cost of ACTforIBD programme compared with control group at 8 weeks and 3 months follow-up. Mean costs will be estimated using: costs of the ACTforIBD programme; outpatient and emergency department visits; hospitalisations; medication use; diagnostic and investigation use; general practitioner visits; and quality-adjusted life-years .

Qualitative analysis

Longitudinal qualitative data collected during semi-structured interviews with participants will be transcribed and analysed thematically, following the main procedural steps of reflexive thematic analysis,^{36–38} which

include: (1) becoming familiar with the data; (2) generating initial codes; (3) constructing initial themes; (4) reviewing potential themes; (5) defining and naming themes and (6) producing the report. All steps will be completed by the first author, with the wider qualitative research team discussing the content of codes and candidate themes, to agree on final themes. We will examine the data inductively, allowing participants' reflections to speak for themselves. In reporting results, we will adhere to Consolidated Criteria for Reporting Qualitative research.³⁹

ETHICS AND DISSEMINATION

Participants will be provided with the Plain Language Statement detailing the participant commitment, potential risks and benefits of the study, and the investigators' contact details. After perusing the plain language statement and before participants are randomised to the groups, they will sign an informed consent form. Participation in the trial is voluntary and the participants can withdraw at any time without providing a reason. Those with severe psychological distress (scores 30–50 on K10) will not be eligible for the study as it is anticipated that these participants would require a more intensive therapeutic approach before benefiting from the current intervention. Similarly, participants whose mental health worsens during the study will be referred back to their treating doctors and provided with publicly available mental health helplines. Only deidentified data will be published. Our article will include a discussion section that details deviations from protocol, and trial limitations related to imprecision in estimates, sources of bias and generalisability threats (as per Consolidated Standards of Reporting Trials - CONSORT). Findings will be disseminated via peer-reviewed publications, conference presentations and briefs on the consumer organisations' websites.

Patient and public involvement

People with IBD were engaged to help develop the ACTforIBD program.²¹ We will also incorporate a comprehensive qualitative evaluation alongside the quantitative data collection to help further inform the programme development to test in future large-scale RCTs.

DISCUSSION

IBD is increasingly considered a disease of brain–gut interaction, with emerging evidence of the bidirectional brain–gut and gut–brain links.⁴ These links frequently translate into a high prevalence of anxiety and depression in people with IBD.⁵ For example, as many as 28% and 20% of people report symptoms of anxiety and depression, respectively, during remission, while these rates rise to 66% and 35% during IBD relapses.⁵ Consistent with brain–gut links, these episodes of anxiety and depression then have a significant impact on IBD disease outcomes, including IBD flares,⁴⁰ and hospital readmissions and surgery.¹⁰ Consequently,

international IBD guidelines⁴¹ and the Australian IBD Standards⁴² recommend not only regular screening for mental illness in people with IBD, but also incorporation of mental healthcare in IBD management. Unfortunately, very few patients actually receive psychological treatment. In our recent Australia-wide study, we have shown that psychological support for people with IBD is very poor, with close to 70% of those with severe psychological distress not seeing a mental health practitioner.^{43 44} This study has the potential to reach many more people with IBD as it is delivered online. However, ACTforIBD relies on therapist support rather than being exclusively self-directed and that means that, while it might reduce the time spent by the therapist with a patient and thus create time for psychologist to see more clients, the programme will likely be administered by hospitals/IBD centres rather than being publicly available.

Psychological treatment to improve adaptation to IBD typically involves CBT, and there is some support for its efficacy on QoL and symptoms of depression.¹² However, the findings of this meta-analysis also highlight the need for trials of psychotherapy interventions specifically targeting the groups at risk of suboptimal outcomes (eg, people with psychological distress), to improve both mental and physical health management in IBD. The proposed intervention will further contribute to understanding the utility of interventions that specially target those with IBD and psychological distress. Other limitations of existing research on psychological interventions include a focus on traditional models of CBT, developed primarily for non-chronic disease populations and without consultation with patients with IBD. This has often resulted in interventions that (1) are not informed by consumer preferences, which is associated with poor treatment retention and low efficacy^{45 46}; and (2) omit important targets for psychological treatment relevant to this population, including psychological strategies to address negative disease-specific beliefs and disease-related thoughts and feelings, such as fear or shame.⁴⁷ It is likely that the core processes of ACT, which include acceptance and cognitive defusion (which can be described as changing the relationship with one's thoughts), are highly suited to chronic, incurable health conditions such as IBD. In addition, existing CBT interventions for IBD offer only short-term benefits for mental health and QoL in IBD,¹² with a dearth of studies focused on psychological distress as the outcome, and little telemedicine used.

Minimal contact therapies have not received adequate attention in gastroenterology, although the preliminary evidence is promising in terms of reducing healthcare seeking behaviour.⁴⁸ In IBD specifically, little research on low intensity or online psychotherapy is available.¹² Online interventions importantly increase equity and access to therapy for people living in rural and remote communities, as well as convenience for all by removing barriers such as time, transportation cost and inconvenience, and motivation to be physically present. As stated above, this intervention can be potentially implemented via existing hospital psychology clinics and made available throughout Australia contributing to integrated biopsychosocial healthcare in

IBD. Given poor access to psychological care reported by people with IBD,⁴⁴ minimal contact online and low-cost therapies, such as the proposed intervention, have the potential to fill a current gap in services. ACT is considered one of the most promising psychological interventions in serious chronic illness.⁴⁹ Due to the efficacy of ACT in other chronic disease populations,^{50 51} it is likely that our intervention may be feasible and improve outcomes in IBD, including psychological distress, disease activity, QoL, fatigue and pain. ACT delivered by a mental health practitioner is a safe, non-invasive, non-drug intervention worth investigating in the context of IBD.

LIMITATIONS

Many psychological interventions conducted to date have produced small-to-moderate effect sizes (though clinically meaningful), such as 0.2 for psychotherapy.⁵² It is possible that ACTforIBD will produce a similarly small effect. However, since the present intervention is accessible to those in rural and remote communities via the hospitals utilising telemedicine appointments and reduces the reliance on therapists in comparison to psychotherapy facilitated by a face-to-face therapist each week, we believe even a small effect is meaningful for the future availability of psychotherapy in hospitals supporting people with IBD in Australia. It is also possible that given the current COVID-19 pandemic, there may be internet fatigue and coupled with ACT being a potentially lesser-known psychotherapy, recruitment might be slow and attrition high. However, we will employ strategies to enhance retention, including a short follow-up, brief surveys and small incentives throughout the study. We decided not to screen participants using a full-length psychological interview. This was dictated by the practicality of this once the intervention is implemented into practice but also by the fact that K10, we will rely on while screening, is a highly sensitive measure which outperforms other commonly used scales and correlates highly with diagnostic measures such as the *Mini* International Neuropsychiatric Interview (MINI).^{53 54}

Author affiliations

¹Psychology, Deakin University Faculty of Health, Burwood, Victoria, Australia

²School of Psychology, Deakin, Geelong, Victoria, Australia

³School of Psychology, Deakin University, Burwood, Victoria, Australia

⁴School of Health Sciences, Swinburne University of Technology, Hawthorn, Victoria, Australia

⁵Faculty of Health, Deakin University, Burwood, Victoria, Australia

⁶Gastroenterology, Monash University Faculty of Medicine Nursing and Health Sciences, Clayton, Victoria, Australia

⁷Crohn's and Colitis Australia, Camberwell, Victoria, Australia

⁸Department of Medicine, Christchurch School of Medicine and Health Sciences, Medicine, Christchurch, New Zealand

⁹Medicine, Dunedin School of Medicine, University of Otago, Dunedin, New Zealand

¹⁰School of Psychological Sciences, University of Tasmania, Hobart, Tasmania, Australia

Twitter Antonina Mikocka-Walus @AntMikocka

Contributors SE contributed to the trial design, drafted the protocol paper, approved its final version and is the guarantor of the submission. LO, MD, SRK, MF-T, EO, PG, LR, RG, AM-W, LVN, SC and DR contributed to the trial design, commented

on drafts of the protocol paper, and approved the final version. AM-W designed the trial, contributed to drafts of the protocol paper and approved the final version.

Funding This work was supported by Crohn's & Colitis Foundation of America Litwin IBD Pioneers Award 2021 (Award Number: 828660).

Competing interests We do not have any conflicts of interest in relation to the present study, however, outside this work, SK has served as an educational speaker for Janssen, Ferring and Takeda. PG has served as a speaker, a consultant and an advisory board member for Allergan, Janssen, MSD, Pfizer, Anantara, Atmo Biosciences, Immunic Therapeutics, Novozymes and Takeda and Bristol-Meyers Squibb, and has received research funding from from MSD and Atmo Biosciences; and RG owns stocks and shares in Atmo Biosciences. LR served on the Roche International Patient Advisory Council and the Takeda IBD Patient Expert Council. RG has served on advisory boards for AbbVie New Zealand and Australia, Zespri New Zealand and Janssen New Zealand and has received research funding from AbbVie. AM-W has served as an educational speaker for Janssen and Ferring.

Patient and public involvement Patients and/or the public were involved in the design, or conduct, or reporting, or dissemination plans of this research. Refer to the Methods section for further details.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Subhadra Evans <http://orcid.org/0000-0002-1898-0030>

Antonina Mikocka-Walus <http://orcid.org/0000-0003-4864-3956>

REFERENCES

- Chang JT. Pathophysiology of inflammatory bowel diseases. *N Engl J Med* 2020;383:2652–64.
- CentreforDiseaseControlandPrevention. Inflammatory bowel disease prevalence (IBD) in the United States: centre for disease control and prevention, 2019. Available: <https://www.cdc.gov/ibd/data-statistics.htm>
- Kaplan GG, Windsor JW. The four epidemiological stages in the global evolution of inflammatory bowel disease. *Nat Rev Gastroenterol Hepatol* 2021;18:56–66.
- Gracie DJ, Guthrie EA, Hamlin PJ, et al. Bi-Directionality of brain-gut interactions in patients with inflammatory bowel disease. *Gastroenterology* 2018;154:1635–46.
- Mikocka-Walus A, Knowles SR, Keefer L, et al. Controversies revisited: a systematic review of the comorbidity of depression and anxiety with inflammatory bowel diseases. *Inflamm Bowel Dis* 2016;22:752–62.
- Knowles SR, Keefer L, Wilding H, et al. Quality of life in inflammatory bowel disease: a systematic review and Meta-analyses-Part II. *Inflamm Bowel Dis* 2018;24:966–76.
- Knowles SR, Graff LA, Wilding H, et al. Quality of life in inflammatory bowel disease: a systematic review and Meta-analyses-Part I. *Inflamm Bowel Dis* 2018;24:742–51.
- Bernstein CN, Singh S, Graff LA, et al. A prospective population-based study of triggers of symptomatic flares in IBD. *Am J Gastroenterol* 2010;105:1994–2002.
- Kochar B, Barnes EL, Long MD, et al. Depression is associated with more aggressive inflammatory bowel disease. *Am J Gastroenterol* 2018;113:80–5.
- Ananthakrishnan AN, Gainer VS, Perez RG, et al. Psychiatric comorbidity is associated with increased risk of surgery in Crohn's disease. *Aliment Pharmacol Ther* 2013;37:445–54.
- Click B, Ramos Rivers C, Koutroubakis IE, et al. Demographic and clinical predictors of high healthcare use in patients with inflammatory bowel disease. *Inflamm Bowel Dis* 2016;22:1442–9.
- Gracie DJ, Irvine AJ, Sood R, et al. Effect of psychological therapy on disease activity, psychological comorbidity, and quality of life in inflammatory bowel disease: a systematic review and meta-analysis. *Lancet Gastroenterol Hepatol* 2017;2:189–99.
- Hayes SC, Strosahl K, Wilson KG. *Acceptance and commitment therapy: an experiential approach to behavior change*. New York: Guilford Press, 1999.
- Ciarrochi J, Fisher D, Lane L. The link between value motives, value success, and well-being among people diagnosed with cancer. *Psychooncology* 2011;20:1184–92.
- Hulbert-Williams NJ, Storey L. Psychological flexibility correlates with patient-reported outcomes independent of clinical or sociodemographic characteristics. *Support Care Cancer* 2016;24:2513–21.
- Hofmann SG, Asmundson GJG. Acceptance and mindfulness-based therapy: new wave or old HAT? *Clin Psychol Rev* 2008;28:1–16.
- Hofmann SG, Sawyer AT, Witt AA, et al. The effect of mindfulness-based therapy on anxiety and depression: a meta-analytic review. *J Consult Clin Psychol* 2010;78:169–83.
- Ost L-G. The efficacy of acceptance and commitment therapy: an updated systematic review and meta-analysis. *Behav Res Ther* 2014;61:105–21.
- Wynne B, McHugh L, Gao W, et al. Acceptance and commitment therapy reduces psychological stress in patients with inflammatory bowel diseases. *Gastroenterology* 2019;156:935–45.
- Trindade IA, Pereira J, Galhardo A, et al. The LIFEwithIBD intervention: study protocol for a randomized controlled trial of a face-to-face acceptance and commitment therapy and Compassion-Based intervention tailored to people with inflammatory bowel disease. *Front Psychiatry* 2021;12:699367.
- Dober M, Mikocka-Walus A, Evans S, et al. Perspectives on an acceptance and commitment therapy (act) based program for patients with inflammatory bowel disease and comorbid anxiety and/or depressive symptoms. *Psychother Res* 2021;31:668–81.
- Bell ML, Whitehead AL, Julious SA. Guidance for using pilot studies to inform the design of intervention trials with continuous outcomes. *Clin Epidemiol* 2018;10:153–7.
- Clara I, Lix LM, Walker JR, et al. The Manitoba IBD index: evidence for a new and simple indicator of IBD activity. *Am J Gastroenterol* 2009;104:1754–63.
- Andrews G, Slade T. Interpreting scores on the Kessler psychological distress scale (K10). *Aust N Z J Public Health* 2001;25:494–7.
- Guidance on applying for feasibility studies London: National Institute for health research, 2019. Available: <https://www.nihr.ac.uk/documents/nihr-research-for-patient-benefit-rfpb-programme-guidance-on-applying-for-feasibility-studies/20474>
- Whitehead AL, Julious SA, Cooper CL, et al. Estimating the sample size for a pilot randomised trial to minimise the overall trial sample size for the external pilot and main trial for a continuous outcome variable. *Stat Methods Med Res* 2016;25:1057–73.
- Norman GR, Sloan JA, Wyrwich KW. The truly remarkable universality of half a standard deviation: confirmation through another look. *Expert Rev Pharmacoecon Outcomes Res* 2004;4:581–5.
- Keeton RL, Mikocka-Walus A, Andrews JM. Concerns and worries in people living with inflammatory bowel disease (IBD): a mixed methods study. *J Psychosom Res* 2015;78:573–8.
- Hayes SC, Levin ME, Plumb-Villardaga J, et al. Acceptance and commitment therapy and contextual behavioral science: examining the progress of a distinctive model of behavioral and cognitive therapy. *Behav Ther* 2013;44:180–98.
- McCombie A, Gearry R, Andrews J, et al. Does computerized cognitive behavioral therapy help people with inflammatory bowel disease? A randomized controlled trial. *Inflamm Bowel Dis* 2016;22:171–81.
- Braun V, Clarke V. *Successful qualitative research*. London: Sage, 2013.
- Davies R, Dart J. *The 'Most Significant Change' Technique - A Guide to Its Use*. CARE International, 2005.
- Tutthill EL, Maltby AE, DiClemente K, et al. Longitudinal qualitative methods in health behavior and nursing research: assumptions, design, analysis and lessons learned. *Int J Qual Methods* 2020;19. doi:10.1177/1609406920965799. [Epub ahead of print: 26 10 2020].
- Teague S, Youssef GJ, Macdonald JA, et al. Retention strategies in longitudinal cohort studies: a systematic review and meta-analysis. *BMC Med Res Methodol* 2018;18:151.
- Eldridge SM, Chan CL, Campbell MJ, et al. Consort 2010 statement: extension to randomised pilot and feasibility trials. *Pilot Feasibility Stud* 2016;2:64.

- 36 Braun V, Clarke V. What can "thematic analysis" offer health and wellbeing researchers? *Int J Qual Stud Health Well-being* 2014;9:26152.
- 37 Braun V, Clarke V. Reflecting on reflexive thematic analysis. *Qual Res Sport Exerc Health* 2019;11:589–97.
- 38 Braun V, Clarke V, Hayfield N, et al. Thematic Analysis. In: Liamputtong P, ed. *Handbook of research methods in health social sciences*. Singapore: Springer Nature, 2018.
- 39 Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care* 2007;19:349–57.
- 40 Mikocka-Walus A, Pittet V, Rossel J-B, et al. Symptoms of depression and anxiety are independently associated with clinical recurrence of inflammatory bowel disease. *Clin Gastroenterol Hepatol* 2016;14:829–35.
- 41 Häuser W, Moser G, Klose P, et al. Psychosocial issues in evidence-based guidelines on inflammatory bowel diseases: a review. *World J Gastroenterol* 2014;20:3663–71.
- 42 CCA. *Australian IBD standards: standards of healthcare for people with inflammatory bowel disease in Australia*. Crohn's & Colitis Australia: Camberwell, VIC, 2016.
- 43 Mikocka-Walus A, Massuger W, Knowles SR, et al. Quality of care in inflammatory bowel disease: actual health service experiences fall short of the Standards. *Intern Med J* 2020;50:1216–25.
- 44 Mikocka-Walus A, Massuger W, Knowles SR, et al. Psychological distress is highly prevalent in inflammatory bowel disease: a survey of psychological needs and attitudes. *JGH Open* 2020;4:166–71.
- 45 Baumeister H, Hutter N, Bengel J. Psychological and pharmacological interventions for depression in patients with coronary artery disease. *Cochrane Database Syst Rev* 2011;9:Cd008012.
- 46 Liu SS, Ziegelstein RC. Depression in patients with heart disease: the case for more trials. *Future Cardiol* 2010;6:547–56.
- 47 Hadlandsmyth K, White KS, Nesin AE, et al. Proposing an acceptance and commitment therapy intervention to promote improved diabetes management in adolescents: a treatment conceptualization. *International Journal of Behavioral Consultation and Therapy* 2013;7:12–15.
- 48 Ahl A, Mikocka-Walus A, Gordon A, et al. Are self-administered or minimal therapist contact psychotherapies an effective treatment for irritable bowel syndrome (IBS): a systematic review. *J Psychosom Res* 2013;75:113–20.
- 49 Hulbert-Williams NJ, Storey L, Wilson KG. Psychological interventions for patients with cancer: psychological flexibility and the potential utility of acceptance and commitment therapy. *Eur J Cancer Care* 2015;24:15–27.
- 50 Kelson J, Rollin A, Ridout B, et al. Internet-Delivered acceptance and commitment therapy for anxiety treatment: systematic review. *J Med Internet Res* 2019;21:e12530.
- 51 Brown M, Glendenning A, Hoon AE, et al. Effectiveness of Web-Delivered acceptance and commitment therapy in relation to mental health and well-being: a systematic review and meta-analysis. *J Med Internet Res* 2016;18:e221.
- 52 Cuijpers P, van Straten A, Bohlmeijer E, et al. The effects of psychotherapy for adult depression are overestimated: a meta-analysis of study quality and effect size. *Psychol Med* 2010;40:211–23.
- 53 Kessler RC, Barker PR, Colpe LJ, et al. Screening for serious mental illness in the general population. *Arch Gen Psychiatry* 2003;60:184–9.
- 54 Spies G, Kader K, Kidd M, et al. Validity of the K-10 in detecting DSM-IV-defined depression and anxiety disorders among HIV-infected individuals. *AIDS Care* 2009;21:1163–8.
- 55 Bodger K, Ormerod C, Shackcloth D, et al. Development and validation of a rapid, generic measure of disease control from the patient's perspective: the IBD-control questionnaire. *Gut* 2014;63:1092–102.
- 56 Khanna R, Zou G, D'Haens G, et al. A retrospective analysis: the development of patient reported outcome measures for the assessment of Crohn's disease activity. *Aliment Pharmacol Ther* 2015;41:77–86.
- 57 Jairath V, Khanna R, Zou GY, et al. Development of interim patient-reported outcome measures for the assessment of ulcerative colitis disease activity in clinical trials. *Aliment Pharmacol Ther* 2015;42:1200–10.
- 58 Knowles SR, Apputhurai P, Burgell R. Development and validation of the gastrointestinal unhelpful thinking scale (guts): a brief self-report measure for clinical and research settings. *Gastrointestinal nursing* 2021.
- 59 Hann DM, Jacobsen PB, Azzarello LM, et al. Measurement of fatigue in cancer patients: development and validation of the fatigue symptom inventory. *Qual Life Res* 1998;7:301–10.
- 60 Osman A, Wong JL, Bagge CL, et al. The depression anxiety stress Scales-21 (DASS-21): further examination of dimensions, scale reliability, and correlates. *J Clin Psychol* 2012;68:1322–38.
- 61 Herdman M, Gudex C, Lloyd A, et al. Development and preliminary testing of the new five-level version of EQ-5D (EQ-5D-5L). *Qual Life Res* 2011;20:1727–36.
- 62 Smith BW, Dalen J, Wiggins K, et al. The brief resilience scale: assessing the ability to bounce back. *Int J Behav Med* 2008;15:194–200.
- 63 Schwarzer R, Jerusalem M. Generalized Self-Efficacy scale. In: Weinman J, Wright S, Johnston M, eds. *Measures in health psychology: A user's portfolio Causal and control beliefs*. Windsor, UK: NFER-NELSON, 1995. : 35–7p..
- 64 Bond FW, Hayes SC, Baer RA, et al. Preliminary psychometric properties of the acceptance and action Questionnaire-II: a revised measure of psychological inflexibility and experiential avoidance. *Behav Ther* 2011;42:676–88.
- 65 Devilly GJ, Borkovec TD. Psychometric properties of the credibility/expectancy questionnaire. *J Behav Ther Exp Psychiatry* 2000;31:73–86.